



# Plagiarism Checker X Originality Report

**Similarity Found: 0%**

Date: Sunday, August 26, 2018

Statistics: 0 words Plagiarized / 3206 Total words

Remarks: No Plagiarism Detected - Your Document is Healthy.

---

Clob.I Vaccrinei! I0 (6): 597-701, 2013 ISSN 1992{t97 O IDo\$ Public..ior8" 2013 DOI: 10.5t /idr!i.9"2013.10.6.1 136 rH&ssnuddin University, 90245 IndorEsit '1D€poftnsl ofBiolos/, Facutiy of Science Universitas Neg6i Mskaasar,90222 Indonesia rDcpa.tDa[ ofAnimal NutritioG Fecuky ofAnimsl Husb€r&y, Hassn|lddh Univ.rriy, Indoncsia 'DeportrE[ ofAnirnal Husbsldry, Provinc. of Sordr Sulawesi \$143. Indonesi rUIM Pcrd.hr School ofscicncr, T.cllmlo5/ ad Innovdbn Polky (-IIM perde. School). t2vcl 6, M€llara R zak. UDive.siti Tcknologi Malaysir Kuala Lumpur, Jalsn Sanaralq 54100 Kuala Lumpur, MahFia 6Deporhc ofPasr and Plsm Disca.c, Fa.uhy ofAgricuhlrc, Hrsanuddh Universiry, Irdooesia AtOr.t Tltc obicdiv. of rhi!p.p6 wB ro cvrlulr. tIE poitibility of \$httuning cl€rhdn gts (Pc,,ir.tt6 p.,rpr"rr) with ricc drerv fid hlr b..m fdmcnt d wirh whiG ,oa filtlgi in 8o.!. A aotd of t2 go@ \*c.! r!"donly eriSrEd ro ft.d dl oli. of !ta|.tn di€8 m.dir8 !o co.rplctdy raftnrir.d bloct d.dsr Th. atimd! wct dividcd i.io lhlc blo.t! .ccoding to rh.n initid body \*cight Tfr. t,cert. t ! w.r:  $RI = I (I(,6 Eldlatr 8rrts, R2 - 706 Elcp||rn lrlss + 3(7. ftrman d ie rtaw, RI - 30\% Elcphrt C,rsi! + Tt\%,fl.dtlld ric. st !w rd R4 = 1007. f.rDc .d ric. !irw.$

Pr[ltd.'3 m.alurEd wcl! &y man.r ard or8ftric ,ln&r di86tibility, liEd col|3unpth! av.nsl daily 8.in md t€.d cfici('rcy of 8066. D6.a analy3i! indici.d that lhlc DM .nd o4rnic Ind6 diggtibility ligniffc.ndy d..rrrs.d lrlhcn th. l.vcl offdDar.d ricc slw irEr.!d ir ih. rdiort. lochlion of fctaall.d ri. irtt} rp to ?'0% in thc raix d iot 3ipifcany s DM od Ot{ disBtibily hn \$bslitutinS grils wiih lura fqtn dr.d ric. rrlsw ortcdly d.cr!.6.d DM ad OM di8dibilitica.

Ttt€ b.n ft.d ctrcicrcy wE foutrd in d'c sod 8iv€.t RI (0.1, follow.d hy R2 (O ! o), Rr (0.09) md Ra (0,04). In cooclurio& ric, rir.w dd h6 b..n f.r|ctltd Eing whi& mt tungi car b. us.d to subdun cl.p||lt Srrar up !o 707. in lhlc rdior ofgoe. Futurc work is io optimiz. ih. la€ ofasrlindu\$ial lra{. ioto quality fc.d with f.rmerrcd by whire mr fingi. Th€ mair pioblco hEcd in rumin prodltion sr{.In h lvril0bility oa f..dstufr, crpocially in ey s..ron.

Therapeutic use of rice straw as a feed supplement for ruminants  
The use of rice straw as a feed supplement for ruminants is a well-established practice. Rice straw is a by-product of rice production and is a rich source of fiber. It is commonly used as a roughage supplement in the diets of cattle, sheep, and goats. The use of rice straw as a feed supplement can help improve the digestibility of the diet and reduce the cost of feed. However, it is important to ensure that the rice straw is properly processed and stored to avoid mold and other quality issues. The use of rice straw as a feed supplement is a sustainable and cost-effective way to utilize a by-product of rice production.

Rice straw is a by-product of rice production. It is a rich source of fiber and is commonly used as a feed supplement for ruminants. The use of rice straw as a feed supplement can help improve the digestibility of the diet and reduce the cost of feed. However, it is important to ensure that the rice straw is properly processed and stored to avoid mold and other quality issues. The use of rice straw as a feed supplement is a sustainable and cost-effective way to utilize a by-product of rice production.

Therapeutic use of rice straw as a feed supplement for ruminants  
The use of rice straw as a feed supplement for ruminants is a well-established practice. Rice straw is a by-product of rice production and is a rich source of fiber. It is commonly used as a roughage supplement in the diets of cattle, sheep, and goats. The use of rice straw as a feed supplement can help improve the digestibility of the diet and reduce the cost of feed. However, it is important to ensure that the rice straw is properly processed and stored to avoid mold and other quality issues. The use of rice straw as a feed supplement is a sustainable and cost-effective way to utilize a by-product of rice production.

Therapeutic use of rice straw as a feed supplement for ruminants  
The use of rice straw as a feed supplement for ruminants is a well-established practice. Rice straw is a by-product of rice production and is a rich source of fiber. It is commonly used as a roughage supplement in the diets of cattle, sheep, and goats. The use of rice straw as a feed supplement can help improve the digestibility of the diet and reduce the cost of feed. However, it is important to ensure that the rice straw is properly processed and stored to avoid mold and other quality issues. The use of rice straw as a feed supplement is a sustainable and cost-effective way to utilize a by-product of rice production.

Therapeutic use of rice straw as a feed supplement for ruminants  
The use of rice straw as a feed supplement for ruminants is a well-established practice. Rice straw is a by-product of rice production and is a rich source of fiber. It is commonly used as a roughage supplement in the diets of cattle, sheep, and goats. The use of rice straw as a feed supplement can help improve the digestibility of the diet and reduce the cost of feed. However, it is important to ensure that the rice straw is properly processed and stored to avoid mold and other quality issues. The use of rice straw as a feed supplement is a sustainable and cost-effective way to utilize a by-product of rice production.

Therapeutic use of rice straw as a feed supplement for ruminants  
The use of rice straw as a feed supplement for ruminants is a well-established practice. Rice straw is a by-product of rice production and is a rich source of fiber. It is commonly used as a roughage supplement in the diets of cattle, sheep, and goats. The use of rice straw as a feed supplement can help improve the digestibility of the diet and reduce the cost of feed. However, it is important to ensure that the rice straw is properly processed and stored to avoid mold and other quality issues. The use of rice straw as a feed supplement is a sustainable and cost-effective way to utilize a by-product of rice production.

qi,dit utd Choro,jir," b\$vn cflici.rn in dcoomposing stra\* and orh.r qop rcsiducs [lal. The obj€ctivc of d'is study va ro l.n thc pclibility of sub\$ituiin8 .leDherl rr!!j (Pennit tun proplre@) with ricc anw dlst hG bcar f.nn.rn d wirh whit iot tungi a3 fc.ding fo. eoor A.b.! .r. r..alrt: 'lrE .\*p.riEcd va cmdu.r.d @.dine b corpldclly rrdmitcd Uo.l d.siSrl Twclvc 8o.t! werE dividcd i'tro \$r€. Sroups mrdior to initirl tody wcighr (four onimlvgmup). E&t mimal i,l cv.'y grou, wa! r&d.mly Sivco or. of fou. E alDd[ di.rs, nrrncly: R1= 10096 Elcphaft grals (Co ml), R2= ?0'l. Elcphrm gr8 + 3091 fcrD.nEd dcc srw. ttl " lt7. ElcdE rllJ. + n !4 f.rm.dd rh. sls, id RF lo(X fcrn.n d !ic. ira\$. In nd&lio!|. !!dr lrlimrl \$B tllc givc. conc.iEi. at dE rlic of 1.5% of body wcisrL Eeh snilul \*rs kepl in hdividud mcrrbolim crd. .nd hE a fi€. a.ccli ro drinling watcr. The study wai carri.d otn for 30 d.ys which 1ras divid.d

ino fir. p.riodr: prtlimit|ay Fio4 d@i6 paiod &d sdnplng Fiod. Himirtly tinc vds irr.nd.d io ob6.rvc E rv6.8p aEount offc.d cq|3llm.d by .rdt drimd. Ttc scoord pcrrl wa in rd.d ro 8iv. dlmc. io ani sl ad@rirtg io crD..imcrnil dia and \$c ls\$ p6iod wa3 for .latu coll.crión \*hich wa! laslinS for icrl dsF. AniE l w.5 fd m lwic. t &y; O7..}O d l5.m md dh nmiml |jr. Siv€n ."Fim.r\*d dia aoud ,9 to !0a.9/ Bw'E [,sb aDimal w0r fd &ily, hr r[ lcnovcr fcd froot Fsrioll' d.y w!. rst'ovcd tul dt fccd h"\* bcforc addirS thc tlsh fccd" Thc ,roomit of f€.d

md rctuscd fccd wa! r.coidcd ro c3lcul6€ f..d MAT!|RIAI-S AND [II]THODS Glotul v.t ritetb, l0 (6): 697-701 , 2013 condiriór Bcforc fldina ro mimsl, E fcnn ri.d ric. !t t{ w6 cxpo.cd b entiqt lcmp6a!8. Thc cqprlrrB conot r ilohr wss p.!9.rEd edilg io pro..dlm of t151. l,roi.tory tu.ttr. TIE fa.d lrnplcs (f.fln€nt d ric! rnlw dd co.rc.rdc) dd fcc6 w.r! ,rdyz.d fo. dry Dtrd (DMI al! lo..l N n 'nri d.rf8ar fih! (NDF) rnd a.id d.t rgert fih! (AD]).

All sarnpl6 w€€ lround to F.s l-rnm lcrEar Fior lo &l'lysis. Pnp.nrio. ol f.ir.ra.d Ri.r Slrrr: In lh. pr€viou! \$dy, dE vEio6 hvcl! of Cqpriro coridrA i!oh. wcr! .dd.d to ric. iraw !ub.trd6 and fcrmcrncd d diffcftnt irrut ioD timc. h w6 fouql drt dE oFimum lcyd orn fcnn(r{ion tilDc d|ar cm h. sFlid io oplimrllly incrcal. thc quality of ricc alraw in t nn3 of irDplovIDg Duritiv. qlElity ard in viEo digBtibility ofric. strlw lub6ud wrs l0c6 (\*/\*) for l0 days. l}ftfG. in thi! dperin.in dd ldcl wE rlycd in r"t4.rioS Thc ric. \*!\* Da dqecd ino !6111 3iE (2-i oD lcrsd') tlr.n

mixcd holnoedo|uly with 1116 (w,rv) ol Coprina co"tda isrrld- Tk mixrrc th€. srs put into plasric b{s and fcrn.ntcd for 30 dsys und.r ,|erotic S.rd. C.aEtL.: Duing nE lirplirf, p..iod (10 d!F), lIE \$,a&ly of dict (f('rrct{.d ric. srg ari oqlccnEr.) oflir.d q Efrs.d (if Ey, e@ 'g'd.d dd) Eming srd ! sub{ampl. l{as bulk d for tuilx, an lysis. Th. d.ily fccal coll..lion sas pin in phrtic b.g! utd wcish.d- Thc blgs, sith f.c.rr w.r! lrhclD lrd ilorcd !! 5'C. Al th.

crlid of.ach colkdion. feccs colLc!8d drily from ...b dE p ov.. dE J.&y p.dd w.ft hdt d on m individd nillld bGn!, mix.d boroqhly utd . \$b- ianplc wa ovco &id (l00oc) io ddcnninc dry

matl! of th. fcc.& Anoth.r 3ub 3.mplc ( 1096) wa irL.rl ad kcpt fiDzar for 3ub6.Sr!'lt  
lrbordor, sn lysi!. 698 Alotul v.t riwr,, l0 (6):697-701.2013 DM cod.tn was dctlDit'cd by  
&yio8 d flfPC in rhc ovco for 24 IL TIE FrEcnt!8c of6h wa da.tmirEd by ignitiog thc  
samplc! for 6 h a. 55OoC. (hganic matcf (OM) \$,as cslculdcd a, loG%a\$ (D O. Fib.!  
oompoliliod (ADF srd NDF) \*.s alBlyzld erding !o [E p(lc.dult of l 16l. Prr.-...r! M..t r!d:  
Th..iy n rlc r (DIO ard ort,nic mat r (OtrO inirkc \*ct€ &&mi.od by 3ubfa.lir8 dtc arnou.t  
ofDM alld OM off.€d otrad wih tlt DM.rd OM digcstibilircs which w€tE dclaflnincd usirS  
formuld Fcod Etucy - AYltTe &it, gi(8y&iY DM Lrlt (s) T-L l Tlidt DM d oM ttrL. DM !d oM  
dttlrlti6 d f..d.lfio.ncy f6 ah tud DM disrnibiliry - (DM innlddFf€ DM GyDM iarcG)xlmTo  
rbl,b fdln d bt df.d ry.Ern r -!iL E Er !if.d, ditdIP{or) Rr ( lota E}rr-r g-l R2 (7ota E4a-r !E +  
lota frud rio slrrl a](l0ta Ek ndtB+ 70\*t Erql ic sr F.r'drd) d Ra (lota Fdn d Rir sd) OM  
Dis6riHlity = (OMi rrdS) - Fd.s OM (8)r'OM intak (g) x l0o'l. &ESI]LTS AND DISCUSSION  
In gal€rd.

coflnlio of dE cxFin€rt0i mimab sl3 in Sood condition lhtoughoul the sldy. DM tnd OM  
diSEsibilitid w€r! siSnificadly rficd.d by drc Mncnts (Tablc l). DM itn t d.(t!..s d e dE lcaclorf  
fsr€in d dc. stEri in thc rdion hErca!.4 ransins noln 57 | -6.1 8/d (Rl ) !o 275.54 S/d (R4).  
Simih.ly OM inta&e dcor!!!€! fiom 522.71 g/d (Rl) to 220.58 S/d (R1). DM and OM  
diScribilitica wcn atr..t d (P<0.05) by thc tEodtrcn! (Td. l). FurdEr rEsr indicdcd fnd DM ,d  
OM of eoat Ec€ivins R l.

R2, R3 wnE significrlily higIF th.n tttnl of R1. wh;lc Rl, R2. Rl !r.r sitrrilr (P{.Ct). Av.rleE d.ily  
gain ofenls dlct R l . R2 ri &1 wB sistrifcdly hishc' thm that of Elatn€ol R4. Similrrly. f.cd  
cfficicoc, for ftamcnB Rl, R2, ai Rl wll bcn rcompordto thsl of Fecd inaatc car b€ u&d a ,n  
indicaor of th. paldbiliry of rhc dicr& The avdlAc valu.

of 8od n civirts 10016 fqndt d ri..3tss (R4) b 275J35ll day in (Tabl. l). This value b low6  
Uran dlc 'lccomndcd lvlcl for fccd i \*r for go3r whici tu 89 - 104.9 g/ @\rr"). AvcnSc  
co'rflmption h th.s. cxFrimcdrt |',t.1a lolD "j6 9.51 0.t0 0.15 004 r&ucd tom 275.54 gday  
(IU)'. ric. strlw fcrncd.d) to 571.64 (100t6 clcphs gri!s). Fccd consunption ofthc 8od! in.hi!  
iody wc bvcr d|m dd ttpo.Ld by tltl who rcort d fc.d id.t€ rmging fiom 529.0 to J47.9  
/day by us;ne a fc.d

codlinins Pmbio! rumal microb.3 oft ftlo ad cralylic nrpplsn ,16. Ev.n rhongh t!E fc.d  
intrr. of Rjl is 3lighlly lowcr thtr thd ofRl, but both rtstistic€lly simulu indiedn& Bln fld  
irtdc of Rl ird R4 a lowc' fhe In dl.' word irElurbo SOta of frm.,r.d ricc ciw io .obdin t  
clcpt8rr Sral! do6 mt ncglivcly atr€.r dlc intikc of .nimlb b'x shituinS cl(?llan gta with l0o(  
f.rtr|cd€d ricc slw ripifi.&rily &t.'iordcs dt€ irnte of mimals. fttu ptlcnocnon miSht b.  
rElat d to low paldb.ility ofstaw s.r \*ell 13 low digeslibility (Tabl. l). Onc physicd ft.or  
ita.rminiDg lhc lvlcl of iniatc of eimd n dE vohmc of rumsL S.'av !l v.y hrky msLrials.

Tndtfo.r lhc dE a"im,l rcc.iving ooly fi!.nlcdd nc. sllrw rcquirB more roon in th€ rllm.n

fm dE !6nc enout of diar d|!o ailnd rccivin8 ooly glir o. .ombindin ofgrast ard f€nnccld ric. srrlw. AndlEr frdor .lclrnitr8 tlr. idd(! of a,imlb b wh.ll th€ n .d for cn is/ tt ! bc(tl mct Accordiry ro ll9l, cdtsrmPion ofdE.rio, b b.sk lly inilricd to m..t tLE ci.rE/ i..\* ofn rlr! 30 th. sdmds will n p c..in8 whlo dtc, hryc bccn fc.lin8 n! .rl. S/ n€cd! havc h..i fulfilled. B'n \$cy rls0 will ltop cdinS wh.n thc ndcn ir tull .v.i

Orcugh oE slctg/ rlquir! nctrt i! mr ma y.t Utilizdioo of t m.r!.d riac cElw lo subditrt ddl'd 8rB! It dE kvcl of3ory. or 70t6 itD mt..ll!. a d.(tr8c on thc diSrnibility of dE .dlto coopdd lo rt|d of RI (cLph.nt 8rr\$ withod fcrn.r&d ric. stlal}). This illplics that ulilization offcncitcd ric. strlw up 10 tfti6 levcl ha! no ,Esrlivc cfrcts on disenibility, Str.td..l Anb"n: All &ra scr. sbjccLd b arslisit of v0rilrEc for complct ly Ehdorhis.block d6i8n usirg dE C,a'esi LirEor Mod.l (GLM) Drocedu.e ofSPSS. 'n'e dift rlixc anong ft€ ltlensn m€.ru wls dcl.'r ndby Du'.Jl Muhiple ReBc Tci [ll.

hngivhg 10016 fcrmc Ed riE faw Datcdly d.s.ar. tl'c DM !,'d OM di8.nibilni6. oic po"ribl. r€{son for dJs i! dar fi.rE is a poritivc a-alociarive cftclE by mixins the grasr with fcrmcnted .ic. rltw up lo \$ar lvcl. Prcviding only fcrm.otcd ric. slraw miShr b. oot crloll8h lo sulpon optim fdm€ tion in th. rurrcn BullirS e significsd d€.rE id thc digcstibility. DM ,rd OM digeriuliti6 eE r dng fion 442lv.b 67.11% dfn,ii 19.65% ta 61.01v...!3p.divcly. DM n|d OM diScstibilhi6 of RI. R2 ud RI w.,! linilat. av.t:8in8 (65.96, 64.30, 6?.ll).% ,lid (67 .07,6.44,65.92Yo tut DM and OM rcsppccrvcly. According to [20], rhc disestibility of.ty m&.r in th. raSe of555J9r i! hish id euld bc or@d lo imFoyc livleo\* glot[th.

In viro di8€dibility of fcmEntld rioc atnw in dri3 ddy is r4.2 17., which b lowcr rhar ntd lhc'lwld h vitm ({t.09?6). This ir ir lgr.snqit wirh onc ftporr.d by [2ll who 'lid dE! dlc in vivo diS6ribility i3 highcr thar digdtlibility of in vilo sirdia Soln fc.ro6 th.t ca! coruihd. lo lhis diffcrtnc.. J! th. FtsaE of nEr.didB ddivcd f'om dE bod, 90 rt l lrialt3 cddlird in tLE fcc.s i! rn c"rymc thr! ir sc cra.d inio thc diScstilc trEt dld is Doa absort d boct, rs w.ll s3 maLriil, drd !..ping c.lls iom ftc walls ofrh. dig.siivc trrct! t221. Avc.lgc daily glir 3d f€d cf6ci.rrcy of th. tosls hayins RI, R2, R3 \*.rc similr cd siSrific!.\$b highcr dEl dxt6.

llc.iviltr RI.It DtfA. &ily ir for RI, Ril ar RI wrs ,.7t 8/d Sinilart f..d cffcisrcy ofd,l. 8od giva RI, Rjz, R3 wer! simiLr. DEspit ll|c d..rt !. tltncl! offccd irlt\*. as thc fcrmcnt d ric! 3taw inc1!a!. io dE rdior! body wci8tu is not .fic.lcd vhn suhit'lins gr8s up !o 7f?6 eilh fcnncdcd ricc a!a\*. This inplca th. drimil! c€tr w€ll diliz. dlc r|Uri.ri oonLn! of lIE did otulining fcrmcnt d ricc sEaw up to 70r( TIE vdrE offc.d cfEcitr i3 r&Uing norn 0.(X (R1) ro 0.15 (RJ). This sholr thd fo. ctEry l lg ofrdirn u!..!Dals cer incfta\* thc wcidt by 4 to lJ grsDs. Nun€ricdly, rfi. high.st f..d





Iqbal and A.G. Ahl, 20(x). Sedy of cdtulotdic ,oil F Irgi and Two Nov6 SFi6 and Mcdiun. JoufInl of ztEijisns univcBity. 7(6): 4511-66. 14. Gaur, A.C., 19tl. Improvinr sil f.itiliry th@8h org ic rccyoling: a m ual of rurrl c.nporin& TAOiUNDP, R.sixal Proj6t. Prcic.r Field Food Ind Agricultrc Orsrnization of rhc Unit€d N.riom. 15. AchrDa4 Mugiom, T.Arlitnti drh C"A-ai.20Il. Prndsr L..rBlrpJrnur.,C(,,IL P. .\$.orSwsd"y!- 16.

Godile, HJa &rd PJ. VII 506! lylo. Forasc ffb6 ady8i& Agricuhunl Hardboo\* No. 379. USDA 17. Stcel, RG.D. and J.H. Toric, 193. Prinsip dan Pros€dur sarilriL surn! P€d.latsn BiooaiL Jrknta. TcrjcmIhan PT Gran.dia (In trdo&ai"n). 18. Krisun, R-, B. Hrycto and K.G. Dm Wiryaw&t, 2m& Pcns@h Konbimsi PcogSlmen Pmbbiik Mikroba Rum.'t &ogsn Supl.rn.n Kshlitft drlsE Pste Tqt drp K.c.nu!o dan Kanl&rinit Rumcfl Domba- Ju.nal Ilmu Tch'olosi vct .inary. la(a): 262-269. (In Indoncsi.n). Erwrnt. 1995. Oflimalilssi 3in!m fdmcentasi .',rncn ,n.lalui s'lplana asi sultu , dcfam!3i, rEduki aisi mdrn dan stinulai pcttmbuhln mikoba pada r.'llak rumirsnsi. Dis.rrsasi. Prograrn Pscas.rjana INritul Peiaiar Bo8or, Bo8or.

Il7 Nm. On Pr6toa T-R- ald R,{. t.n& 1987. I,fttnins Rumina PEducrion systcm \*irh Availabl. Rmtl,6 ir thc Tmpics Ird Subnopica. P.dunbul Book Armi&I€, Arrrdio" Tongdilinti[ F.K., 192. Fccd DiS6libility Esimation in Rumin nls by In Viro M.thod BIPP Unhas, Uj'ms Pu'd.r's (In IDdor'6iu). TillD.,! AJ., H. Hrt di !'i S. RcfloladiFodi,, Prai/iroeu\$mo, S dj| L.bdo.ocroh, S. 199t. Il[u Mak&un Tcnit D!.rr. Cadah M.da Udvcrsity Pllss, Yo8i8l(ana. (In IndorEsi!"). Sir68ar.

S.B., 1994. R rlsn TcrEt Ruhi'lai& PlrEbar s\*.daya(tn IndorBim). Global vet iN,@. 1 0 (6) : 697-70 t, 201 3 t9. 20. 2t. 22. 23.

INTERNET SOURCES:

-----