



Plagiarism Checker X Originality Report

Similarity Found: 21%

Date: Jumat, September 13, 2019

Statistics: 786 words Plagiarized / 3814 Total words

Remarks: Medium Plagiarism Detected - Your Document needs Selective Improvement.

363 Proceedings of ICMSTEA 2016 : International Conference on Mathematics, Science, Technology, Education, and their Applications, Makassar, Indonesia, 3rd – 4th October 2016 A Simplified Model of *Aspergillus niger* Growth using *Nephrolepis biserrata* leaves for Exo-polygalacturonase Production Halifah Pagarra Department of Biology, Faculty Mathematic and Natural Science, Universitas Negeri Makassar Email: ifah_rani@yahoo.co.id Abstract.

A simple model to describe the growth of *Aspergillus niger* during solid state fermentation using *Nephrolepis biserrata* leaves is a substrate for the production of exo-polygalacturonase for 192 hours. The rate of kinetics growth of *Aspergillus niger* and the use of glucose as substrate using a Model Monod has shown that the maximum utilization of glucose 0.32 g/L of *Aspergillus niger* biomass of 0.496 mg/gdsf of glucose uptake 5.96 g/L to 0.32 g/L at 24 hours up to 120 hours of incubation.

Through alarrgrsion peoleai $\mu_{max} = 0.043$ per hour and $K_s = 0.473$ g/L for 120 hours of incubation. The values of kinetic parameters and the amount of biomass has a significant relationship with R^2 equal to 0.801 and exo-polygalacturonase optimum activity at 54.64 U/g. Data gathered for the kinetic parameters during evaluation of the production of exo-polygalacturonase in SSF by *Aspergillus niger*, a value $Y_{X/S}$ is 0.149 mg/g by using glucose from the medium. The values for the kinetic coefficient of exo-polygalacturonase income is $Y_{E/S} = 16.43$ U / g, $Y_{E/X} = 114.4$

U/mg and kinetic constants were $q_E = 4.79$ U/(mg.h) and $q_S = 0.289$ g/(mg.h). Keyword: Monod Model, *Nephrolepis biserrata* leaves, kinetics growth of *Aspergillus niger*, exo-polygalacturonase production. INTRODUCTION The role of microorganisms in the Income Pectinase Microorganisms play an important role in the production of pectinase

for large-scale industrial use.

Pectinase can be produced by solid state fermentation (SSF) or submerged fermentation (SmF). A variety of environmental factors for the fermentation process affects the production of pectinase. Among these factors are the concentration of nutrients, pH, temperature, humidity and the effect of inducer.

In addition, carbon and nitrogen sources indicate an overall effect on the productivity of pectinase (Catarina et al., 2003). Microorganisms which are a major source of industrial enzymes comprise 50% of fungi and yeast, 35% of the bacteria, while the remaining 15% are either from animals or plants.

Filamentous fungi described as microorganisms can secrete a variety of enzymes involved in the breakdown and recycling of complex biopolymers from both plant and animal tissues. The enzymes secreted these fungi are mostly hydrolytic and plays an important role in the growth of fungi, carbon and nitrogen. Potential fungal enzyme production and secretion of the attention of many researchers to study the production of various types of enzymes using filamentous fungi on large-scale industrial processes (El-Enshasy et al., 2008).

Among the most practical filamentous fungus in solid fermentation process is like genera *Mucor*, *Ascomycetes*, *Aspergillus niger*, *Fusarium* and *Penicillium decumbens oxyporum* (Mamma et al., 2007). For bacteria are used is *Bacillus* sp. (Akcan et al., 2012; Sukumaram et al., 2005; Kumar et al., 2012), *Streptomyces* sp. (Basha et al., 2009). As for the yeast used is *Saccharomyces* and *Candida* (Kumar et al., 2015).

Microorganisms filament is widely used in fermentation sink (SMF) and fermentation solid state (SSF) for the production of pectinase were *Aspergillus niger* (Akhter et al., 2011; Dartora et al., 2002; Okafor et al., 2010; 364 Proceedings of ICMSTEA 2016 : International Conference on Mathematics, Science, Technology, Education, and their Applications, Makassar, Indonesia, 3rd – 4th October 2016 Mamma et al. , 2007).

However, pectinase production by *Aspergillus* strain found to be more effective in solid-state fermentation process compared with submerged fermentation process (Thangaratham et al., 2014). *Aspergillus* sp. is one of the main agents of decomposition and decay of lignocellulosic waste and has the ability to produce various types of enzymes.

Glucosidase high level in *Aspergillus* sp. is important to complete the conversion of biomass into simple sugars lignocellulosa. In addition, *Aspergillus* sp. also known as

bioremediation agents effective (Verma et al., 2011). In the food industry, the *Aspergillus niger* has been exploited mainly in the processing of food (bread and starch) and drinks (wine, juice, beer and distilling industry).

In the industrial production of pectinase also between fungi *Aspergillus niger* is the most frequently used (Rashmi et al., 2008). The use of *Aspergillus niger* is generally recognized as safe (Generally Recognized as Safe) by the State Food and Drug Administration (FDA) US (Schuster et al., 2002).

Most of the kinetic model to describe the growth of microbes in bioprocessing is empirical and based on the Monod equation Monod or modifications which take account of microbial growth inhibition caused by the product or substrate concentration is high (Altiok, 2004). Specific growth rate achieved in the exponential phase is dependent on the concentration of nutrients.

If all the nutrients needed for growth is present in excess, the specific growth rate (μ) and the concentration of nutrients or substrate (S) can be expressed by Monod kinetics as shown in Equation 1. (1) K_s and concentrations of all other essential nutrients is constant. K_s remained constant known as saturated or semi-constant velocity is equal to the concentration of the rate-limited when certain growth rate is equal to half of the maximum (Shuler and Kargi, 2002). This equation has been used as a linear equation such as Equation 2.

(2) -Burk plot, gives a straight line with a gradient (Okpokwasili and Nweke, 2005). Next, the product is also defined as the weight of the product and substrate used in Leudeking Piret equation like Equation 3 and Equation 4. (3) which, $Y_{E/S}$ is the ratio of the enzyme to the substrate (X_0 to S) during incubation.

Kinetics of enzyme-based enzyme activity (U/g), estimated the equation Leudeking and Piret (Aquilar, et al., 2001), is the coefficient estimated from a linear correlation between the concentration of enzyme activity and substrate utilization. (4) which, $Y_{E/S}$ is the ratio of the enzyme to the substrate (X_0 to X).

$Y_{E/X}$ is the coefficient of correlation is estimated to linear line between the activities of exo-polygalacturonase and total biomass of *Aspergillus niger*. Specific enzyme production rate is q_E has been defined as Equation 5. $q_E = \mu \cdot E/X$ (5) 365 Proceedings of ICMSTEA 2016 : International Conference on Mathematics, Science, Technology, Education, and their Applications, Makassar, Indonesia, 3rd – 4th October 2016 Next, a specific substrate utilization rate is q_s was defined as Equation 6 METHODS AND MATERIALS Spores of *Aspergillus niger* ATCC120120 used in this study were collected

from the stock culture collection of Bioprocess and Applied Biology Laboratory, Faculty of Chemical Engineering and Energy, Universiti Teknologi Malaysia. Cultured stocks for the long term is provided in the form of a bead of culture and frozen at -80°C .

Solid State Fermentation The *Nephrolepis biserrata* leaves of the substrate pretreated. Each of the leaves of the pretreatment weighed as much as 10 g/flask into a 500 ml Erlenmeyer flask and added together with basal medium ((NH_4) SO_4 0.01% (w/v), $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ 0.05% (w/v), KH_2PO_4 0.05% (w/v), $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ 0.0005% (w/v), pectin of *Nephrolepis biserrata* leaves 1.0 % (w/v), yeast extract 0.1 % (w/v) and glucose 1.0 % (w/v).

The moisture adjusted to 70% of humidity at pH 5.5 by the addition of medium. The substrate was sterilized using an autoclave at a temperature of 121°C for 20 minutes and stored at room temperature before it was inoculated. After cooling, the inoculation was carried out with 10% of inoculums and incubated at 28°C . Enzyme Extraction Fermented samples were collected every 24 h.

In each sampling, 1 g of the sample was drawn out and mixed with 25 mL of citrate phosphate buffer (pH 4.0). The mixture was then placed on a vortex for 1 minute to ensure that enzyme or sugars on the surface were well-mixed with the buffer. Next, the suspension was centrifuged at 4000 rpm at 4°C for 20 minutes to separate the solid and liquid phase.

The supernatant was used as a crude enzyme for various assays. Analytical Procedures 1 g sample is removed and the fermentation is collected every 24 hours. For each sample, 1 g samples were removed then mixed with 10 mL of citrate phosphate buffer (pH 5.0). The mixture on a vortex for 1 minute to ensure all the enzyme or sugar on the surface of the media extracted into the buffer.

Next, suspended material at a speed of 4,000 rpm, 4°C for 30 minutes to separate the solids (pellets) and liquid (supernatant or crude enzyme). The supernatant was used to research the content of exo- polygalacturonase enzymes and proteins. Determination of enzyme activity is based on the DNS method kit (Dinitro Salicylic Acid Reagent) with some modifications.

This method seeks to determine the carbonyl group of reducing sugars (Miller, 1959). For exo- polygalacturonase activity can be identified by the color intensity by using a spectrophotometer at wavelengths of 540 nm (Kumar et al., 2010). One unit of enzyme exo- pygaauriaounpof educinsugar (reducing sugar), which is measured in terms of glucose, produced as a result of the action of enzymes in the extract 1.0 ml of 1 min at

35° C ± 1° C (Ali et al., 2010).

Determination of Biomass and Growth Kinetic of *Aspergillus niger* The determination of the biomass is carried out with 'Indirect Biomass Estimation' based of Ramachandran et al., (2005). Fungal biomass estimates made to determine the N-acetyl-Glucosamine is produced by acid hydrolysis of chitin contained in the fungal cell wall. A total of 0.5

g of fermentation is mixed with concentrated sulfuric acid (2 ml) and put in a 500 ml conical flask and the mixture is kept for 24 hours at a temperature of ± 30° C. The mixture is diluted with distilled water to make a solution of 1 N and autoclaved (15 psi for 1 hour at 121° C) and then filtered through filter paper. This solution is then neutralized with 4 N NaOH up to 100 ml with distilled water.

1 ml of solution mixed with 1 ml reagent acetyl acetone and incubated in a boiling water bath for 20 minutes. After cooling, ethanol (6 ml) was added followed by the addition of 1 ml of Ehrlich reagent and incubated at 65° C for 10 minutes. After cooling, the optical density of the reaction mixture was read with a spectrophotometer at 530 nm.

Results are expressed as mg glucosamine per gram of dried substrate fermentation $qS = X/S$ (6) 366 Proceedings of ICMSTEA 2016 : International Conference on Mathematics, Science, Technology, Education, and their Applications, Makassar, Indonesia, 3rd – 4th October 2016 (mg/gdsf). The determination of the growth kinetics of *Aspergillus niger* and use of substrate (glucose) is carried out using variables and data that generate exo-polygalacturonase in optimum condition.

Glucose is the total count of glucose from the medium (glucose, *Nephrolepis biserrata* leaves pretreated, *Nephrolepis biserrata* leaves pectin and nutrients) reducing sugar-based DNS method (Miller, 1959). The incubation time for this experiment is over 192 hours and every 24 hours samples were taken and analyzed. The rate of growth kinetics of *Aspergillus niger*, glucose utilization and *Aspergillus niger* biomass and exo-polygalacturonase activities .

Determination of kinetic constants calculated using the growth model of Monod and the product is also defined as the weight of the product (enzyme) with the substrate is calculated by the method Leudeking-Piret. Constant a ncoeffien kiniparets ered om s ae s ax, Y X/S, YE/S, YE/X, qS and qE. RESULT AND DISCUSSION Specic owt at(µmcrorganis e erlred o e miites atoccurdurth logarithmic growth phase.

To determine the pattern of interaction between the use of glucose as the substrate with the analysis of the growth of *Aspergillus niger* glucose utilization and growth of

microorganisms has been carried out during the production of the enzyme in this study. The *Aspergillus niger* growth pattern with an initial substrate concentration of 5.96 g/L under the optimum conditions of this study are shown in Figure 2. Figure 2 shows a time of adaptation to the environment metabolic *Aspergillus niger* found a very short time of incubation. This time is also the beginning of a phase of exponential (logarithmic phase) in which the initial glucose 5.96 g/L with the biomass 0.053 mg/gdsf reduced to 2.43 g/L with the biomass 0.18 mg/gdsf on 24 hour incubation time.

After entering the 24 hour incubation time, the exponential phase very quickly in the use of glucose as an energy source for the growth of *Aspergillus niger* and up to 120 hours of incubation time, the use of glucose is at 0.32 g/L with the biomass at 0.50 m/gdsf. In Figure 2, the utilization rate of glucose as substrate by *Aspergillus niger* showed glucose levels decreased significantly and indicates that the substrate has been used to the maximum.

high or low concentrations of glucose residues affected by *Aspergillus niger* ability to convert the carbon source contained in the substrate into biomass and products (Sharma, 2012). The *Aspergillus niger* optimal biomass production is dependent on the use of glucose that has been done in solid state fermentation. The concentration of glucose is used as substrate in these fermentations.

Microorganism growth pattern is followed logistics model as shown in Figure 2, where the maximum utilization of glucose 0.32 g/L with the growth of *Aspergillus niger* biomass is up to 0.496 mg/gdsf. FIGURE 2. Relationship between *Aspergillus niger* biomass growth rate and glucose consumption in Solid State Fermentation on a 24 hour incubation time up to 192 hours

Incubation time (hour)	Bio mass of <i>Aspergillus niger</i> (mg/gd sf)	The use of Glucose (g/L)
0	0	5.96
1	0.1	5.96
2	0.2	5.96
3	0.3	5.96
4	0.4	5.96
5	0.5	5.96
6	0.6	5.96
24	0.18	2.43
48	0.36	1.215
72	0.54	0.7425
96	0.72	0.5625
120	0.90	0.3825
144	1.08	0.2025
168	1.26	0.0225
192	1.44	0

Proceedings of ICMSTEA 2016 : International Conference on Mathematics, Science, Technology, Education, and their Applications, Makassar, Indonesia, 3rd – 4th October 2016

Umax and Ks value determination is made based on the curve 1/u versus 1/S as shown in Figure 3 from the uptake of glucose 5.96 g/L to 0.2 g/L at 24 hours and 120 hours of incubation. A linear regression equation obtained from the graph is equivalent to $Y = 10.89 + 23.02x$.

From the obtained μ_{max} our (maximum growth rate) was 0.473 g/L (saturation constant) from *Aspergillus niger*. The value of the growth is defined as the number of *Aspergillus niger* biomass produced per amount of substrate used, namely $Y_{X/S}$ is 0.149 mg/g. The activities of exo- pygaourse, μ h e atof *Aspergillus niger* biomass and the use of the substrate in the incubation time of 24 hours up to 192 hours.

Figure 4 (a), (b) and (c) show the results obtained coefficients of linear equations or a straight line from the Leudeking Piret equation. The coefficient YX/S is the coefficient estimated from the straight line correlation between the biomass of *Aspergillus niger* and the concentration of the substrate. FIGURE 3.

μ_{max} and K_s for the growth of *Aspergillus niger* and substrate utilization Coefficient YX/S is defined as the total biomass of *Aspergillus niger* income of 0.149 milligrams per gram of substrate used. YE/S of 16.43 U/L is the estimated coefficient of linear correlation between the activity and concentrations of exo-polygalacturonase glucose utilization. This shows that by 16.43 exo-activity unit of exo- polygalacturonase produced per gram of glucose consumed.

In Figure 4 (c) shows a linear correlation between the activity of exo-polygalacturonase and total biomass of *Aspergillus niger* YE/X of 111.4 U/mg. This value is defined as the total income of exo-polygalacturonase activity of 111.4 units per milligram amount of biomass produced by *Aspergillus niger*. Leudeking-Piret model was also used to obtain the specific content of exo-polygalacturonase income, ie q_E at 4.79 U/(mg.h) and the specific use of glucose as substrate, q_S at 0.096 g/(mg.h). The values of kinetic parameters of exo-polygalacturonase shown in Table 1.

$y = 10,893x + 23,029$ $R^2 = 0,8018$ 0 10 20 30 40 50 60 -3 -2,5 -2 -1,5 -1 -0,5 0 0,5 1 1,5 2 2,5 3 3,5 1/S 1/u 368 Proceedings of ICMSTEA 2016 : International Conference on Mathematics, Science, Technology, Education, and their Applications, Makassar, Indonesia, 3rd – 4th October 2016 a) b) c) FIGURE 4. (a) Graph the relation $(E - E_0)$ against $(S_0 - S)$ coefficients YE/S ,

(b) Graph the relation $(X - X_0)$ against $(S_0 - S)$ determine the coefficient YX/S and (c) relationship graph $((E - E_0)$ against $(X_0 - X)$ coefficients YE/X $y = 16,439x - 48,467$ $R^2 = 0,8261$ 0 10 20 30 40 50 60 0 1 2 3 4 5 6 $(E - E_0)$ $(S_0 - S)$ $y = 0,149x - 0,4341$ $R^2 = 0,845$ 0 0,1 0,2 0,3 0,4 0,5 0 2 4 6 $(X - X_0)$ $(S_0 - S)$ $y = 111,42x - 0,9111$ $R^2 = 0,9979$ 0 10 20 30 40 50 60 0 0,1 0,2 0,3 0,4 0,5 $(E_0 - E)$ $(X_0 - X)$ 369 Proceedings of ICMSTEA 2016 : International Conference on Mathematics, Science, Technology, Education, and their Applications, Makassar, Indonesia, 3rd – 4th October 2016 TABLE 1.

Value kinetic parameters of exo-polygalacturonase production by *Aspergillus niger*. Parameters Value of kinetics parameters μ_{max} 0.043 per jam K_s 0.473 g/L YE/S 16.43 U/g YX/S 0.149 mg/L YE/X 111.4 U/mg q_E 4.79 U/(mg.h) q_S 0.289 g/(mg.h) Monod model also clearly links the data findings (Rigon et al. 2009; Gerard et al., 2006). The specific content of exo-polygalacturonase, ie q_E at 4.79 U/(mg.h), showed that 4.79 units of exo-polygalacturonase activity per gram of biomass per hour *Aspergillus niger*. The

specific rate of glucose as substrate, namely q_s at 0.289 g/(mg.h), shows that the concentration of 0.289 grams of glucose as substrate used every milligram of *Aspergillus niger* biomass per hour.

CONCLUSION Therefore, it was concluded that the relationship between the use of the substrate, the growth rate of *Aspergillus niger* and the enzyme exo-polygalacturonase can be deduced from kinetic parameters were obtained. When there is a good relationship between the amount of biomass and the production of enzymes, the kinetic of glucose as a substrate for the growth of *Aspergillus niger* with enzyme activity even in direct proportion, namely by increasing the amount of biomass has increased the enzyme activity.

ACKNOWLEDGMENT The author are grateful to Bioprocess and Applied Biology Laboratory, Faculty of Chemical Engineering and Energy, Universiti Teknologi Malaysia for providing the research facilities. REFERENCE 1. Akcan, N., Serin, B and Uyarb, F. (2012). Production and Optimization Parameters of Amylases from *Bacillus subtilis* RSKK96 Under Solid State Fermentation. Chem. Biochem. Eng. Q., 26(3): 233-239. 2.

Akhter, N., Morshed, M.A., Uddin, A., Begum, F., Sultan, T. and Azad, K. A. (2011). Production of Pectinase by *Aspergillus niger* Cultured in Solid-State Media. International Journal of Boiscience, 1(1): 33-42. 3. Ali, J., Ahmed, J.S., Assad, Q., Hussain, A., Abid, H., and Gul, F. (2010). Optimization of Pectinase Enzyme Production using Sour Oranges Peel (*Citrus aurantium* L) as Substrate. Pak. J. Biochem. Mol. Biol., 43(3): 126-130. 4. Aquilar, C.N.,

Augur, C., Favela, E.T. and Viniegra, G.G. (2001). Production of Tannase by *Aspergillus niger* Aa-20 in Submerged and Solid State Fermentation: Influence of Glucose and Tannic Acid. Journal of Industrial Microbiology, 26: 296-302. 5. Basha, N.S., Rekha, R., Komala, M. and Ruby, S. (2009).

Production of Extracellular Anti-leukaemic Enzyme Lasparaginase from Marine Actinomycetes by Solid State and Submerged Fermentation: Purification and Characterisation. Tropical Journal of Pharmaceutical Research, 8(4), 353-360. 6. Carol, S.K.L., Chenyu Du., Koutinas, A., Wang, R. and Colinwebb. (2008). Substrate and Product Inhibition Kinetics in Succinic Acid Production by *Actinobacillus succinogenes*. Biochemical Engineering Journal, 41: 128 – 135. 7.

Catarina, omš, Pedr F. and José. (2003). Continuous production of *ctaby* Immobilized Yeast Cells on Spent Grains. Journal Of Boiscience And Bioengineering, 96(6): 513-518. 8. Dartora, A.B., Bertolin, T.E., Bilibio, D., Silveira, M.M. and Costa, A.V. (2002). Evaluation of Filamentous

Fungi and Inducers for the Production of Endo-Polygalacturonase by Solid State Fermentation. Z. Naturforsch.

57: 666-670 370 Proceedings of ICMSTEA 2016 : International Conference on Mathematics, Science, Technology, Education, and their Applications, Makassar, Indonesia, 3rd – 4th October 2016 9. El Azaly, Y.(2008). Optimization Of The Industrial Production Of Alkaline Protease By *Bacillus licheniformis* In Different Production Scales. Australian Journal of Basic and Applied Sciences, 2(3): 583- 593 10. Gerard, A. M., Paca, J., Kostekova, A. J., Stiborova, M. and Soccol, C. R.

(2006), Simple Models for the Continuous Aerobic Biodegradation of Phenol in a Packed Bed Reactor. Braz. Arch. Bioly. Technoly, 49(4): 669-676 11. Kumar, D.P., Thangabalan, B., Rao, P.V. and Yugandhar, N.M. (2010). Production of Pectinase Enzyme by *Aspergillus niger* using *Ficus religiosa* Leaves in Solid-State Fermentation. IJPT., 3(1): 1351-1359. 12. Kumar, M.D.J., Saranya, G.M., Suresh, K., Priyadharshini, A.D., Rajakumar and Kalaichelvan. P.T. (2012).

Production and Optimization of Pectinase from *Bacillus* sp. MFW7 using Cassava Waste. Asian Journal of Plant Science and Research, 2 (3):369-375. 13. Kumar, V.G., Maheswari, N. And Gupta, A. (2015). Isolation and Screening of Fungal Isolates for Multienzyme Production Through Submerged and Solid State Fermentations. Journal Bioprocess Biotech, 5(8): 1-7. 14. Mamma, D., Kourtoglou, E. And Christokopoulos, P. (2007).

Fungal Multienzyme Production on Industrial By-Products of the Citrus-Processing Industry, Bioresource Technology 99: 2373 – 2383. 15. Miller., G.L. (1959). Use of Dinitrosalicylic Acid Reagent for Determination of Reducing Sugar. Anal. Chemical, 1: 426-428. 16. Okafor, U.A., Okochi, V.I., Nwodo, S.C., Ebuehi, T. And Onygame, B.M.O. (2010). Pectinolytic Activity of Wild-Type Filamentous Fungi Fermented on agro-wastes. African Journal of Microbiology Research, 4(24): 2729-2734 17. Okpokwasili, G.C. and Nweke, C.O. (2005). Microbial Growth and Substrate Utilization Kinetics.

African Journal of Biotechnology, 5(4): 305-317. 18. Ramachandran, S., Roopesh, K., Nampoothiri, K. M., Szakacs, G. and. Pandey, A. (2005). Mixed Substrate Fermentation for the Production Of Phytase by *Rhizopus* spp. using Oilcakes as Substrates. Proress Biorllem., 40(5): 1749-1754. 19. Rashmi, R., Murthy, S.K., Sneha, S., Syama, A. And Radhika, V.S. (2008).

Partial Purification and Biochemical Characterization of Extracellular Pectinase from *Aspergillus niger* Isolated from Groundnut Seeds. Journal of Applied Biosciences, 9(1): 378 – 384. 20. Rigon, M.S., Alberto, L. J. L., Lorenci, A.W. and Ricardo, C.S. (2009). A

Simplified Model For A. Niger F_{s3} Growth During Phytase Formation In Solid State Fermentation. Brazilian Archives Of Biology And Echnology, 52: 151-158 21. Schuster, E., Dunn-Coleman, N.,

Frisvad, J.C., and van Dijck, P.W.M. (2002). On The Safety of Aspergillus niger. Appl Microbiol Biotechnol, 59:426-435 22. Sharma, R. (2012). Pathogenecity of Aspergillus niger in Plants. Cibtech Journal of Microbiology, 1 (1): 47-51. 23. Shuler M. L., Kargi F. (2002). .Bioprocess Engineering Basic Concepts. 2nd ed. Prentice-Hall, Inc., USA 24. Sukumaran, R.,K., Singhanian, R.,K. and Pandey, A. (2005). Microbial Cellulases – production, application, and challenges.

Journal of Scientific and Industrial Research, 64: 832-844. 25. Thangaratham, T. and Manimegalai, G. (2014). Optimization and Production of Pectinase Using Agro Waste by Solid State and Submerged Fermentation. Int.J.Curr.Microbiol.App.Sci, 3(9): 357-365. 26. Verma, N., Bansal, M. C. and Kumar, V. (2011). Scanning Electron Microscopic Analysis of Aspergillus niger Pellets and Biofilms under Various Process Conditions.

International Journal of Microbiological Research 2(1): 08-11, 2011.

INTERNET SOURCES:

<1% -

<http://eprints.unm.ac.id/3753/7/Proceeding%20Internasional%20-%20Oldeman%20Climate%20Zoning%20for%20the%20Agricultural%20Area.pdf>

5% - <http://ojs.unm.ac.id/index.php/icmstea/article/download/2702/1441>

1% - <http://ojs.unm.ac.id/index.php/icmstea/article/view/2702>

<1% -

https://www.researchgate.net/publication/259700658_Production_of_pectinases_by_Asp_ergillus_niger_in_solid_state_fermentation_at_high_initial_glucose_concentrations

<1% - <https://www.sciencedirect.com/science/article/pii/S1369703X12000769>

<1% -

https://www.researchgate.net/publication/223858246_Utilization_of_whole_wheat_flour_for_the_production_of_extracellular_pectinases_by_some_fugal_strains

<1% - <https://www.sciencedirect.com/science/article/pii/B978012803725600011X>

<1% - <https://www.sciencedirect.com/science/article/pii/B9780444521149500104>

<1% - <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4673338/>

<1% -

https://www.academia.edu/22381501/L-asparaginase_as_a_critical_component_to_combat_Acute_Lymphoblastic_Leukaemia_ALL_a_novel_approach_to_target_ALL

<1% - <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4444909/>
<1% - <https://www.sciencedirect.com/science/article/pii/S0045653519310549>
<1% -
https://mafiadoc.com/yeasts-in-food-and-beverages_5cb27f8d097c476a2a8b4590.html
<1% -
https://mafiadoc.com/sanjay-kumar-gupta-a-anushree-malik-faizal-bux-_5a1df8561723ddb99b1d53dc.html
<1% - <https://www.sciencedirect.com/science/article/pii/S0734975011001650>
<1% - <https://patents.justia.com/patent/20050233421>
<1% -
<https://www.thefreelibrary.com/Reaction+engineering+studies+on+the+biodegradation+of+anthracene+on...-a0159508059>
<1% -
https://www.researchgate.net/post/How_do_you_calculate_the_maximum_growth_rate_of_an_algae_culture
<1% -
https://www.academia.edu/6571529/Exponentially_fed_batch_cultures_as_an_alternative_to_chemostats_The_case_of_penicillin_acylase_production_by_recombinant_E_coli
<1% - <https://www.ijraset.com/files/serve.php?FID=5691>
<1% - <http://eujournal.org/index.php/esj/article/download/3585/3392>
<1% - <https://www.sciencedirect.com/science/article/pii/S0959652619305256>
1% -
http://eprints.utm.my/id/eprint/52540/1/MadiyahMd.Salleh2014_Effectofphysicalpretreatment.pdf
1% -
<https://jurnalteknologi.utm.my/index.php/jurnalteknologi/article/download/3212/2371>
<1% - <https://www.sciencedirect.com/science/article/pii/S0957582017301453>
<1% -
[http://www.ifrj.upm.edu.my/19%20\(04\)%202012/56%20IFRJ%2019%20\(04\)%202012%20Nora%20\(011\).pdf](http://www.ifrj.upm.edu.my/19%20(04)%202012/56%20IFRJ%2019%20(04)%202012%20Nora%20(011).pdf)
<1% -
https://www.canadianjbiotech.com/CAN_J_BIOTECH/Archives/v1/i1/CJB-2016-102.pdf
<1% -
http://www.scielo.br/scielo.php?script=sci_arttext&pid=S1519-69842015000100010
<1% -
https://www.researchgate.net/publication/215714717_Determination_of_microbial_biomass_and_fungal_and_bacterial_distribution_in_cattle_faeces
<1% - <https://patents.google.com/patent/EP1753869B1/en>
<1% - https://explorers.larc.nasa.gov/HPMIDEX/pdf_files/06_NASA-HDBK-6022b.pdf
<1% - <http://biodiversitas.mipa.uns.ac.id/D/D1702/D170253.pdf>

<1% -

<http://ejobios.org/download/biochemical-study-of-the-effect-of-nephrolepis-biserrata-pretreatment-on-exo-polygalacturonase-6193.pdf>

<1% -

https://www.researchgate.net/publication/304241442_The_Effect_of_Some_Sugars_on_the_Growth_of_Aspergillus_niger

<1% - http://shodhganga.inflibnet.ac.in/bitstream/10603/5180/7/07_chapter%202.pdf

<1% - https://issuu.com/sep2011--now/docs/5_355a1a14202664

<1% -

<https://biotechnologyforbiofuels.biomedcentral.com/articles/10.1186/s13068-016-0575-1>

<1% - http://www.engineering.iastate.edu/brl/files/2011/10/brl_ethanolfermentation.pdf

<1% - <https://www.sciencedirect.com/science/article/pii/S1359511315301288>

<1% - <http://slidehtml5.com/zykj/sjgj/basic>

<1% - <http://ojs.unm.ac.id/icmstea/article/download/2654/1392>

<1% - <http://ojs.unm.ac.id/index.php/icmstea/article/download/2713/1452>

<1% - <https://www.mdpi.com/1420-3049/24/12/2298/htm>

2% - <https://jurnalteknologi.utm.my/index.php/jurnalteknologi/article/view/12222>

1% -

https://www.researchgate.net/publication/11850805_Production_of_tannase_by_Aspergillus_niger_Aa-20_in_submerged_and_solid-state_fermentation_Influence_of_glucose_and_tannic_acid

<1% -

https://www.academia.edu/19985568/Solid_state_and_submerged_fermentation_for_the_production_of_bioactive_substances_a_comparative_study

<1% -

https://www.researchgate.net/publication/223802152_Substrate_and_Product_Inhibition_Kinetics_in_Succinic_Acid_Production_by_Actinobacillus_succinogenes

1% - <https://link.springer.com/article/10.1007/s10295-003-0071-z>

<1% -

https://www.researchgate.net/publication/275340162_RESEARCH_ARTICLE_OPTIMIZATION_AND_PRODUCTION_OF_PECTINASE_FROM_BACILLUS_SUBTILIS_MTCC_441_BY_USING_ORANGE_PEEL_AS_A_SUBSTRATE

<1% - <http://www.juit.ac.in/profiles/115.docx>

<1% - <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3544265/>

1% - <http://scholar.google.co.in/citations?user=banSojwAAAAJ>