

PAPER NAME

Characterization And In Vitro Antifungal Assay Against Fusarium oxysporum F.SP . Passiflora Of Endoph

AUTHOR

Hilda Karim

WORD COUNT

4854 Words

CHARACTER COUNT

28129 Characters

PAGE COUNT

9 Pages

FILE SIZE

512.3KB

SUBMISSION DATE

Jul 29, 2022 5:20 PM GMT+8

REPORT DATE

Jul 29, 2022 5:23 PM GMT+8

● **6% Overall Similarity**

The combined total of all matches, including overlapping sources, for each database.

- 6% Internet database
- 0% Publications database

● **Excluded from Similarity Report**

- Crossref database
- Submitted Works database
- Quoted material
- Small Matches (Less than 10 words)
- Crossref Posted Content database
- Bibliographic material
- Cited material
- Manually excluded sources

CHARACTERIZATION AND *IN VITRO* ANTIFUNGAL ASSAY AGAINST *FUSARIUM OXYSPORUM* F.SP.*PASSIFLORA* OF ENDOPHYTIC ACTINOMYCETES FROM PURPLE PASSION FRUIT PLANTS OF SOUTH SULAWESI, INDONESIA

ALIMUDDIN ALI^{1*}, MUHAMMAD JUNDA¹, HILDA KARIM¹ AND SRI NURYANI²

¹Laboratory of Microbiology, Department of Biology, Faculty of Mathematic and Natural Sciences,
Universitas Negeri Makassar. South Sulawesi, 90233, Indonesia

²Graduate School of Biology Department, FMIPA, Universitas Negeri Makassar, Indonesia

(Received 25 October, 2016; accepted 5 December, 2016)

Key words: *Endophytic actinomycetes, Purple passion fruits, Antifungal, Indole-3-acetic acid*

Abstract - Endophytic actinomycetes from purple passion fruit were screened and evaluated for their antifungal activity against of pathogenic fungi. A total of 32 isolates were obtained from people passion garden in Malino distric, South Sulawesi province, Indonesia. Morphological and chemotaxonomical analysis indicated that twenty-two isolates belonged to the streptomycetes like-strain. This genus was the most dominant among the isolates (68.75%) of plant organs (roots, leaves and stem barks) of these showed inhibitory activity against *Fusarium oxysporum* f.sp. *passiflora*. Actinomycetes were most commonly recovered from leaves (50% of all isolates), followed by roots (43.75%) and stems (6.25%). Based on the phenotypic properties and phylogenetic and analysis of 16S rRNA gene, the selected strain EML-D1A5 was assigned to *Streptomyces durhamensis* strain CSSP538 (97% similarity).

INTRODUCTION

Association of endophytic microbe with their hosts are unique interaction because the microbe inhibits the inner organs and tissues of plants such as roots, stems and leaves without causing diseases (Coelho *et al.* 2011). Endophytes with this capacity might profit from association with the plant, because colonization is enhanced. In turn, host plants benefit by biocontrol potential agent against fungal root and seed rots (Yuan *et al.* 1995). Although the interaction between two organisms is not, yet, fully understood, over recent years they have been progressively extensively employed, either in agriculture (Ryan *et al.* 2008) or antimicrobe, anticancer (Li *et al.* 2008) and production of valuable pharmaceutical compounds (Stoble and Long, 1998). The presence of endophytic microbe in healthy plant crops has been demonstrated in the roots of plants such as maize (Araujo *et al.* 2000), banana (Cao *et al.* 2005) and some medicinal plants (Taechowisan and Lumyong 2003; Passari *et al.*

2015). Endophytic Actinomycetes have positive effect on host plant by producing phytohormone. Plant-associated Actinomycetes are rich sources of bioactive compounds including indole-derived molecules such as phytohormone indole-3-acetic acid (IAA) (Ting *et al.* 2008; Jasim *et al.* 2014), production auxin and gibberellins (Brown, 1972; Merckx *et al.* 1987). The metabolites of endopytic Actinomycetes inhibit a number of microbes (Gurney and Mantle, 1993; Yu *et al.* 2010), antibacterial activities (Castillo *et al.* 2006) and antifungal activities (Gupta *et al.* 1995; Ezra *et al.* 2004). Some endophytic bacteria exert several beneficial effect on host plants such as induction of resistances to plant pathogen (Chen *et al.* 1995; Sturz and Matheso, 1996), and nitrogen fixation (Kirchhorf *et al.* 1997). Numerous recent studies showed promising in endophytic Actinomycetes research. Application of microbial consortia that interact synergistically each other, they enhance the plant growth and protect from phytopathogens (Kurth *et al.* 2014). Therefore, studies on

*Corresponding Author's email- muddin_69@unm.ac.id

biological control of plant diseases used endophytic Actinomycetes origin to be an alternative to protect locally agricultural crop plants.

Purple passion fruit is one of the economically important fruits particularly in South Sulawesi, Indonesia for decades. Recently, these plants began to decrease due to a fungal diseases pathogen. *F. oxysporum* f. sp. *passiflora* is a strain which causing a vascular wilt plant of purple passion fruit. The fungal are the most destructive plant pathogens causing wilt fungal diseases in many crop plants. Fungal strain devastated a great number of passion fruit plantation. Various methods were applied to controlling the diseases such as application of fungicides and planting of fungal resistance varieties, but these method were uneffective. A cost-effective measure of control for the disease is still not available. At this time, many methods practically use to control of diseases along with the need to develop sustainable methods of diseases management has started the hunt for a suitable alternative control. Biological control agent may be used as an alternative management strategy because it is not only suppressing the diseases and increasing crop yield but also reducing the enviromental pollution due to chemical pesticides (Achari and Ramesh. 2014). Several studies have shown that endophytic bacteria isolated from crops such as tamato and chili (Amaresan *et al.* 2012), chickpea (Misk and Franco. 2011) and citrus (Araujo *et al.* 2002). However, the screening of antifungal endophytic actinomycetes of purple passion fruits has remained unexplored.

The present study was carried out to isolate Actinomycetes from roots, stem bark and leaves of passion fruits, widely cultivated in South Sulawesi, Indonesia. The microbes were screened for antifungal activities for the purpose of this study, endophytic Actinomycetes were isolated and screened *in vitro* for their abilities to inhibit fungal pathogens.

MATERIALS AND METHODS

Plant Material. Samples (roots, leaves, stem bark) of purple passion fruits were collected from passion fruits of farm garden at Malino district, South Sulawesi province, Indonesia. Plant samples were kept in sterilized plastic bags and stored at 4°C until isolation. Isolation of strain was carried out immediately after samples were sent back to the laboratory.

Isolation and selection of Actinobacteria. All the collected samples of the purple passion fruits were washed with running tap water to remove soils and debris. Isolation of endophytic actinomycetes was done by cutted the tissues into small pieces (0.2 x 4 cm²) and subjected to surface sterilization. The excised roots, leaf and bark stems were surface-sterilized using by serial treatment of etanol 70% (v/v) for 10 min, 1% sodium hypochlorite for 2 min. Finally, root tissue, leaf and stem bark were washed in sterilized distilled water for a 5 minutes, then dried using sterile filter paper. Approximately 0.1 x 1 cm² of plant tissue were cutted and transferred onto starch casein (SC) agar supplemented with nystatin 100µg/mL. The surface sterilization process was confirmed by spreading aliquots of the sterile distilled water from the final rinse on SC agar medium, followed by incubation at 35°C, and observation of microbial growth. If there was no visible growth of microbial colony on the surface of agar plates, the sterilization was assumed completed. Colonies of endophytic actinomycetes appeared surrounding of plant sample tissue after incubation at 35°C for 2-3 weeks. Purified of endophytic actinomycetes cells was done by transferred of colonies onto freshly mannitol-soy agar medium plate until a single colony showed purity. The pure culture was maintained at a 15% sterile glycerol suspension at -80°C for long-term preservation.

Screening for antifungal antagonism. *In vitro* antifungal activities of endophytic actinomycetes isolates were assessed by using dual assay antagonistic method against the *Fusarium oxysporum* f. sp. *passiflora* (Barakate *et al.* 2002). Each of actinomycetes isolated obtained from isolation proces was spreaded onto SC agar medium and incubated 35°C for 7 days. A plug agar of isolates (6 mm in diameter) was tranferred onto the edgen of sabaroud agar plate, while the tested fungi inoculated at the edge of the 9 cm plate on the same media. The plates were incubated 30°C for 7 days. Tested fungal plugs were also placed on uninoculated actinomycetes used as control. The endophytic actinomycetes was showed inhibition growth against to inhibited growth fungi considered as endophytic actinomycetes producing antifungal. The fungal inhibition was calculated from the equation:

$$R = \frac{(R1 - R2)}{R1} \times 100$$

Where: R = fungal inhibition (%), R1 = the fungal growth radius of control culture, R2 = the distance of fungal growth in the direction of actinomycete colony.

Morphological and physiological characterization

To study cultural characteristic of selected isolate, the strain EML-D1A5 was grown on several media, namely yeast extract-malt extract agar (ISP2), oatmeal agar (ISP3), inorganic salts-starch agar (ISP4), (ISP7). Aerial and substrate mycelia were determined by comparison with colour chips from the Color Harmony Manual (Jacobson *et al.* 1958). In order to examine morphological characteristics, the selected strains were grown on SC agar by using slide culture method and spore morphology were assessed by light microscopy.

Physiological characteristics of strain EML-D1A5 were determined as follow. Hydrolysis of starch was determined as described by Gordon *et al.* (1974). Utilization of carbohydrates and nitrogen as sole carbon and nitrogen sources was tested by using medium (ISP9), respectively as described by Shirling and Gottlieb. (1966). The temperature range for growth was determined on ISP2 in a temperature gradient incubator.

Molecular characteristics and Phylogenetic analysis.

The 16S rRNA gene from selected strain was amplified and sequenced. The DNA genome of strain EML-D1A5 extraction and purification was carried out using the Pure link genomic DNA kit (In-vitrogen, Carlsbad, CA, USA) according to the manufacturer instructions. The 16S rRNA gene was amplified using the following primers following primers: 27f (5'-AGAGTTTGATCCTGGCTCAG-3') and 1492r (5'-GGTACCTTGTTACGACTT-3') to amplify 1500bp fragment. The PCR cycling was conducted as follow: pre-denaturation of the target DNA at 96°C for 3 min followed by 30 cycles at 95°C for 1 min, primer annealing at 56°C for 1 min, and primer extension at 72°C for 5 min, the reaction mixture was held at 72°C for 5 min. The resulting PCR products were visualized by electrophoresis in 1.5% (w/v) agarose gels stained with ethidium bromide (Petrosyan *et al.* 2003). The amplicon of DNA fragments was sequenced using sequencer model ABI 3100 sequencer according to manufacturers' instructions (ABI PRISMA 3100 Genetic Analyzer User's Manual).

The nucleotides obtained were subjected to BLAST analysis using by the NCBI database deposited in NCBI GenBank. The 16S rRNA gene sequence of strain EML-D1A5 was aligned with

those of representatif strain members of other 39 Streptomycete genera using CLUSTAL-X. Phylogenetic trees were deduced by using the neighbor-joining method using the Phylip version 3.5 (Saitou and Nei. 1987) with bootstrap values based on 1000 replication (Felsenstein. 1985).

RESULTS

Evaluation of the surface sterilization protocol

A total of 32 of Actinobacteria endophytic strain were collected from purple passion plant health at five sampling locations of passion fruit plants of farming garden in Malino distric, South Sulawesi, Indonesia. Evaluation of samples from all actinomycetes cultured with the surface sterilization protocol showed no Actinomycetes growth on SC agar. It showed that epiphytic Actinomycetes were eliminated, so the strain were obtained indeed endophytic. The strains were classified into streptomycete-like strain (characterized by abundant aerial mycelium with powdery spores) and non-streptomycetes strain (slimy colonies coloured red, oranges and brown to black). The strains were grouped into different plant organs such as root, leaves and stem bark on agar starch casein agar media. In vitro antagonistic assay was used to select Actinomycetes which produce antifungal against phytopathogenic fungi. There are five isolates obtained from roots and leaves inhibited the growth of fungal tested. However, no isolate from stems bark inhibites fungi (Table 1).

Screening for antifungal antagonistic

Five of 23 isolates obtained showed strong antagonisms against to *Fusarium oxysporum* f.sp.

Table 1. Endophytic actinomycetes was obtained from plant organs of purple passion fruit plantation in South Sulawesi, Indonesia.

Group	No. of isolates		
	Roots	Leaves	Stem bark
Streptomycete-like strain	8 (3)	13 (2)	1
Non-streptomycetes strain	6 (1)	3	1
Total	14	16	2
Percentage	43.75	50	6.25

Number in brackets indicate that the number of isolates that inhibit of *Fusarium oxysporum* f.sp. *passiflora* by testing *in vitro*

passiflora (Fig. 1). Moreover, the selected strain has been demonstrated for a variety of plant pathogen and showing inhibitory activities against many species of tested fungi such as *Phytophthora* sp, *Aspergillus* sp, and *Trichoderma* sp (data not shown). The percentage of inhibitory of selected actinomycetes strain showed that the five selected actinomycetes strains have inhibitory potency pathogenic fungi with inhibitory level more than 60 % at the end of incubation (Fig. 2). Strain EML-D1A5 showed the highest inhibitory among selected strains (83.6%), whereas the the lowest is EML-A2P1(69 %).

Preliminary identification of selectes endophytic strain

Morphological and physiological properties of strain EML-D1A5 shown in Figure 3. According to microscopic morphology observation (100x magnification) showed that selected strain has a typical character as the *Streptomyces* genera. The strain showed the structure of spore chains opened spiral.

DISCUSSIONS

In the present study, there were 23 Actinomycetes strains obtained from the purple passion fruit plants. The result showed that endophytic Actinomycetes can be isolated amongs inside the roots, leaves and stem bark of purple passion fruits. Majority of the strain from roots and leaves were *Streptomyces* like strain indicating that mainly *Streptomyces* spp genera are existing as endophytic in purple passion fruits. Endophytic streptomycete actinomycetes (SA) and nonstreptomycete actinomycetes (NSA) have been isolated from within live tissues of various plant species (Coombs and Franco 2003; Rosenblueth and Martinez-Romero 2006). *Streptomyces* is also the most dominant endophytic genus in the roots of plants such as tomato (Coa *et al.* 2004; Sardi *et al.* 1992), wheat (Justin and Christopher. 2003) and some medicinal plants (Qin *et al.* 2009).

Predominance of streptomycetes in any habitat is well known both in rhizosphere area and or in plant tissue. The presence of endophytic Actinobacteria associated with plant tissues may play important roles in plant both health and development. Many studies reported that bacteria (such as Actinobacteria) protect plant against pathogens like fungal diseases (Zhao *et al.* 2012).

Moreover, the endophytic bacteria may influences of their metabolic product for plant growth and physiologic (Gupta *et al.* 1995). Endophytes with this capacity might profit from association with the plant, because colonization is enhanced.

Antagonistic activities of microbe have been reported against several fungal plant pathogens fungal such as bacteria, actinomycetes and fungi. Here we report the antifungal properties of endophytic actinomycetes isolates of passion fruit against *Fusarium oxysporum* f.sp. *passiflorae*. Inhibiting capacity of pathogenic fungi beside passion fruits-*Fusarium* wilt fungi showed that strain EML-D1A5 has high potency as fungal biocontrol agent. The capabilities of the strain to inhibit of passion fruit-*Fusarium* wilt pathogen and other genera of fungal show that this strain most prospect applied for pathogen fungal controlling particulary on agricultural crop plant. Some endophytic bacteria direct the potential benefit on the host plant, such as antagonist against fungal pathogen including including *Fusarium oxysporum* (Coa *et al.* 2005), *Rhizoctonia solani* (Sadeghi *et al.* 2006) and *Verticillium dahliae* (Meschke and Schrempf. 2010), *Gaeumannomyces graminis* var. *tritici* and *R. solani* (Coombs *et al.* 2004). In addition, endophytic Actinomycetes has been reported the stimulation of plant growth through the formation of growth hormones (Sturz *et al.* 1997; Stoltzfus *et al.* 1997; Reinhold-Hurek and Hurek 1998) and induction of plant resistance to pathogens (Liu *et al.* 1995). Mechanisms of action of endophytic actinomycetes in suppressing pathogen are production of substances particulary secondary metabolite (antibiotics), cell wall degrading enzymes (chitinase) and competition for nutrient (sideropore)(El-Tarabily and Sivasithamparam 2006).

Strain EML-D1A5 was observed to grow well on some of the media tested, including ISP1, ISP 2, ISP 3 (oatmeal agar), ISP 5 and Bennet but it has no growth on ISP4 (inorganic salts-starch). The aerial and substrate mycelia were white on all tested agar media. Utilization on of carbon sources was determined on ISP 9 medium supplemented with sterile carbon sources. The strain utilized glucose, galactose, manitol, xylose, raffinose, sucrose and dextrose but not lactose, rhamnase, fructose, sorbitol. Besides the different of genotypic character, there are many phenotypic differences between strain EML-D1A5 and the most closely related species of the genus *Streptomyces* (Table 5). The phylogenetic tree (Fig.3) reconstructed on




Fig 1. *In vitro* inhibition activity of selected endophytic actinomycetes strain against test fungi A). *Fusarium oxysporum* f. sp. *passiflorae*, B); Strain EML-A₂P₁; C). Strain EML-A₁L₁; D). Strain EML-A₁N₃; E). Strain EML-D₁A₃ and F). Strain EML-D₁A₅

Table 2. Cultural characteristics of strain EML-D1A5 on different culture medium

Agar Medium	Aerial Mycelium		Substrate mycelium		Growth	Soluble pigment
	HCM	Colour	HCM	Colour		
ISP1	RAL 1013	Oyster white	RAL 1014	Ivory	Good growth	-
ISP2	RAL 1013	Oyster white	RAL 1014	Ivory	Good growth	-
ISP3	RAL 1015	Light ivory	RAL 1015	Light ivory	Moderate growth	-
ISP4	-	-	-	-	No growth	-
ISP5	RAL 1015	Light ivory	RAL 1014	Ivory	Good growth	-
BENNET	RAL 1013	Oyster white	RAL 1013	Oyster white	Moderate growth	-

HCM, Harmony Colour Manual

Table 3. Physiological properties of strain EML-D1A5 on physico-chemical factor

	pH	NaCl (%)	Temperature (°C)
3	-	0	4
4	-	3	25
5	+	4	30
6	+	5	35
7	+	6	37
8	++	7	40
9	+	8	45
10	+/-	9	
11	+/-	10	
12	-	11	

++= good growth (), += moderate growth (), +/- = weak growth, - = no growth

Table 4. Utilization of nitrogen and carbon sources of strain EML-D1A5

No	Nitrogen sources utilization	Carbon sources utilization
1	(NH ₄) ₂ SO ₄	D-glucose
2	AgNO ₃	D-lactose
3	(NH ₄) ₂ Fe(SO ₄) ₂ 6H ₂ O	D-galactose
4	C ₂ H ₈ N ₂ O ₄	D- mannitol
5	(NH ₄)(NO ₃)	D-xylose
6	NaNO ₃	Rhamnose
7	KNO ₃	Raffinose
8	Fe(NO ₃) ₃	Sucrose
9	Ba(NO ₃) ₂	L-fructose
10	L-valine	Sorbitol
11	L-glutamic acid	Cellobiose
12	L-asparagine	Maltose
13	L-arginine	Dextrose
14	Iso leucine	Myo-Inositol
15	DL α-amino n- butyric acid	Arabinose

++ : growth, well utilized , +: growth, moderate utilized, +/- : growth, weakly utilized, -: not utilized

Table 5. Phenotype and physiological properties of strain EML-D1A5 and related species

Characteristic	Strain EML-D1A5	<i>Streptomyces durhamensis</i> AS 4.1699	<i>Streptomyces puniscabiei</i> KACC 20253	<i>Streptomyces filipinensis</i> AS 4.1452
Morphology and pigmentation				
Aerial mass colour	W	GR	WO	G
Substrate mycelium	W	WB	WB	Y
Spore chain morphology	Spiral	Spiral	Rectiflexuous	Spiral
Soluble pigment on ISP3	-	-	-	-
Carbon utilization				
Raffinose	+	+	+	+
D-xylose	+	W	+	+
Rhamnose	-	-	+	-
D-Manitol	+	+	+	+
Lactose	-	-	ND	ND
L-Arginine	+	+	+	-
pH4	-	-	+	+
7% NaCl	-	-	+	-

+, Well utilized/ present; W, weakly utilized; 2, not utilized/absent; ND, not determined. W, white; GR, grey-red; WO, white-orange; G, grey; WP, white-purple; WB, white-brown; Y, yellow; BR, brown-red.




Fig 2. Inhibition of tested fungal of *Fusarium oxysporum* f.sp. *passiflorae* by selected endophytic actinomycetes strain on Starch Casein agar for different incubation periods

the basis of 16S rRNA gene sequences indicated that strain EML-D1A5 was phylogenetically most closely affiliated to the genus *Streptomyces*. Strain EML-D1A5 shared similarity to *Streptomyces durhamensis* ATCC 23194 (97 % similarity). Based

on the phenotypic properties and phylogenetic analysis of 16S rRNA gene, the strain EML-D1A5 was assigned to *Streptomyces durhamensis* strain CSSP538 clade (NCBI Genbank).




Fig 3. Morphology of colony and mycelium structure of endophytic selected strain (magnification 400X). A Strain EML-A₂P₁; B). Strain EML-A₁L₄; C). Strain EML-A₁N₃; D). Strain EML-D₁A₃ and E). Strain EML-D₁A₅

ACKNOWLEDGMENT

We would like to acknowledge the Ministry of Research and Technology, Higher Education of

Indonesia for the financial support under grant HIKOM 2016.

REFERENCES

- Achari, G.A and Ramesh, R. (2014). Diversity, biocontrol, and plant growth promoting abilities of xylem residing bacteria from Solanaceous crops. *Int J Microbiol.* 58: 135-147.
- Amaresan, N., Jayakumar, V., Kumar, K and Thajuddin, N. (2012). Endophytic bacteria from tomato and chili, their diversity and antagonistic potential against *Ralstonia solanacearum*. *Arch Phyto Plant Protec.* 45 (3): 344-355
- Araujo, J.M., Silva, A.C and Azevedo, J.L. (2000). Isolation of endophytic actinomycetes from roots and leaves of maize (*Zea mays* L.). *Braz Arch Biol Technol.* 43: 447-451.
- Araujo, W.L., Marcon, J.W., Jr Macheroni, V.E and Azevedo, J.L. (2002). Diversity of endophytic bacterial populations and their interaction with *Xylella fastidiosa* in citrus plants. *Appl Environ Microbiol.* 68: 4906-4914
- Bano, N and Musarrat, J. (2003). Characterization of a new *Pseudomonas aeruginosa* strain NJ-15 as a potential biocontrol agent. *Curr Microbiol.* 46: 324-328
- Barakate, M., Ouhdouch, Y., Oufdou, K.H and Beaulieu, C. (2002). Characterization of rhizospheric soil Streptomyces from Moroccan habitats and their antimicrobial activities. *World J Microb Biot.* 18: 49-54.
- Brown, M.E. (1972). Plant growth substances produced by microorganisms of soil and rhizosphere. *J Appl Bacteriol.* 35: 443-451
- Cao, L., Qiu, Z., You, J., Tan, H and Zhou, S. (2005). Isolation and characterization of endophytic streptomycete antagonists of Fusarium wilt pathogen from surface-sterilized banana roots. *FEMS Microbiol Lett.* 247:147-152. doi: 10.1016/j.femsle.2005.05.006
- Cao, L. X., Qiu, Z. Q., You, J. L., Tan, H. M and Zhou, S. (2004). Isolation and characterization of endophytic Streptomyces antagonists of Fusarium wilt pathogen from surface sterilized banana roots. *FEMS Microbiol Lett.* 247: 147-152. doi: 10.1016/j.femsle.2005.05.006
- Castillo, U., Strobel, G.A and Mullenberg, K. (2006). Munumbicins E-4 and E-5: novel broad spectrum antibiotics from Streptomyces NRRL 3052. *FEMS Microbiol Lett.* 255: 296-300
- Chen, C., Bauske, E.M., Mussan, G., Rodriguez-Kabana, R and Kloepper, J.W. (1995). Biological control of Fusarium wilt on cotton by use of endophytic bacteria. *Biol Control.* 5: 83-91
- Coelho, M.M., Ferreira-Nozawa, M.S., Nozawa, S.R and Santos, L.W. (2011). Isolation of endophytic from arboreal species of the Amazon and identification by sequencing of 16S rRNA encoding gene. *Gen Mol Biol.* 34(4): 676-680
- Coombs, J. T and Franco, C. M. M. (2003). Isolation and identification of actinobacteria from surface sterilized wheat roots. *Appl Environ Microbiol.* 69: 5603-5608. doi: 10.1128/AEM.69.9.5603-5608.2003




Fig 3. Neighbor-joining phylogenetic tree inferred from 16S rRNA gene sequences. The phylogenetic tree shows the phylogenetic relationship of endophytic bacteria strain EML-D1A5 with related genera. Bootstrap values are expressed as percentages of 1000 replications. Bootstrap values, $\geq 50\%$ are shown at branch points. Score bar represents 1 nucleotide substitution per 100 nucleotides.

Coombs, J.T., Michelsen, P.P and Franco, C.M.M. (2004). Evaluation of endophytic actinobacteria as antagonists of *Gaeumannomyces graminis* var. *tritici* in wheat. *Biol Control*. 29:359-366

Felsenstein, J. (1985). Confidence limits of phylogenies: an approach using the bootstrap. *Evol.* 39: 783-791. doi:10.2307/2408678

El-Tarabily, K.A and Sivasithamparam, K. (2006). Nonstreptomycete actinomycetes as biocontrol agents of soil-borne fungal plant pathogens and as plant growth promoters. *Soil Biol Biochem*. 38: 1505-1520

Ezra, D., Castillo, U and Strobel, G.A. (2004). Coronamycins, peptide antibiotics produced by a verticillated *Streptomyces* sp. (MSU-2110) endophytic on *Monstera* sp. *Microbiol* 150: 785-793.

Gordon, R. E., Barnett, D. A., Handerman, J. E and Pang, C. H.-N. (1974). *Nocardia coeliaca*, *Nocardia autotrophica*, and the nocardin strain. *Int J Syst Bacteriol*. 24:54-63.

Gupta, R., Saxena, R.K., Chaturverde, P., Virdi and J. S. (1995). Chitinase production by *Streptomyces viridificans*: its potential in fungi cell wall lysis. *J Appl Bacteriol*. 78: 378-383

Gurney, K. A and Mantle, P.G. (1993). Biosynthesis of 1-N-methylalbonoursin by an endophytic *Streptomyces* sp. isolated from perennial ryegrass. *J Nat Prod* (Lloydia). 56 :1194-1198

Jacobson, E., Grauville, W. C and Fogs, C. E. (1958). *Color Harmony Manual*, 4th ed. Chicago: Container Corporation of America.

Jasim, B., Joseph, A. A., John, C.J., Mathew J and Radhakrishnan, E. K. (2014). Isolation and characterization of plant growth promoting endophytic bacteria from the rhizome of *Zingiber officinale*. *Biotech*. 4:197-204

- Kirchhorf, G., Reis, V.G., Baldani, J.L., Eckert, B., Dobereiner, J and Hartmann, A. (1997). Occurrences, physiological and molecular analysis of endophytic diazotrophic bacteria in gramineous energy plants. *Plant Soil*. 194: 45-55
- Li, J., Zhao, G.Z., Chen, H.H., Wang, H.B., Qin, S and Zhu, W.Y. (2008). Antitumour and antimicrobial activities of endophytic streptomycetes from pharmaceutical plants in rain forest. *Lett Appl Microbiol.* 47:574-580. doi: 10.1111/j.1472-765X.2008.02470.x
- Liu, L., Kloepper J.W and Tuzun, S. (1995). Induction of systemic resistance in cucumber against *Fusarium wilt* by plant growth-promoting rhizobacteria. *Phyto Path.* 5:695-698
- Misk, A and Franco, C. (2011). Biocontrol of chickpea root rot using endophytic actinobacteria. *BioControl* 56: 811-822. doi:10.1007/s10526-011-9352-z
- Merckx, R., Dijkstra, A., Hartog, A. D and Veen, J. A. V. (1987). Production of root-derived material and associated microbial growth in soil at different nutrient levels. *Biol. Fert. Soils*, 5: 126-132.
- Meschke, H and Schrenpf, H. (2010). *Streptomyces lividans* inhibits the proliferation of the fungus *Verticillium dahliae* on seeds and roots of *Arabidopsis thaliana*. *Microb Biotechnol*, 3:428-443
- Qin, S., Li, J., Chen, H.H., Zhao, G.Z., and Zhu, W.Y. (2009). Isolation, diversity and antimicrobial activity of rare actinobacteria from medicinal plants of tropical rain forests in Xishuangbanna, China. *Appl Environ Microbiol.* 75:-6186.
- Passari, A. K., Mishra, V. K., Saikia, R., Gupta, V. K and Singh, B. P. (2015). Isolation, abundance and phylogenetic affiliation of endophytic actinomycetes associated with medicinal plants and screening for their in vitro antimicrobial biosynthetic potential. *Front Microbiol.* 6: 273. Doi:10.3389/fmicb.2015.00273
- Petrosyan, P., Garcia-Verela, M., Luz-Madrigal, A., Eluitron, C and Flores, M.E. (2003). *Streptomyces mexicanus* sp. nov., a xylanolytic microorganism isolated from soil. *Int J Sys Evol Microbiol.* 53: 269-273.
- Reinhold-Hurek, B and Hurek, T. (1998). Life in grasses: diazotrophic endophytes. *Trend Microbiol.* 6:139-144
- Rosenblueth and Martinez-Romero. (2006). Bacterial endophytes and their interactions with hosts. *Mol Plant Micro Inter.* 19 (8): 827-837.
- Ryan, R.P., Germaine, K and Franks, A. (2008). Bacterial endophytes: recent developments and applications. *FEMS Microbiol Lett.* 278: 1-9.
- Saitou, N and Nei, M. (1987). The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol.* 4:406-425
- Sadeghi, A., Hessian, A. R., Askari, H., Aghighi, S and Shahidi-Bonjar, G.H. 2006. Biological control potential of two *Streptomyces* isolates on *Rhizoctonia solani*, the causal agent of damping-off of sugar beet. *Pak J Biol Sci.* 9:904-910
- Sardi, P., Saracchi, M., Quaroni, S., Petrolini, B., Borgonovi, G. E and Merli, S. (1992). Isolation of endophytic *Streptomyces* strains from surface-sterilized roots. *Appl Environ Microbiol.* 58: 2691-2693.
- Shirling, EB and Gottlieb, D. (1966). Methods for characterization of *Streptomyces* species. *Int J Syst Bacteriol.* 16:313-340
- Stoble, G.A and Long, D.M. (1998). Endophytic microbes embody pharmaceutical potential. *ASM News.* 64: 263-268
- Sturz, A.V and Matheson, B.G. (1996). Populations of endophytic bacteria which influence host-resistance to *Erwinia*-induces bacterial soft rot in potato tubers. *Plant Soil.* 184:265-271
- Sturz, V.V., Christie, B.R., Matheson, B.G and Nowak, J. (1997). Biodiversity of endophytic bacteria which colonize red clover nodules, roots, stems and foliage and their influence on host growth. *Biol Fertil Soils.* 25:13-19
- Stoltzfus, J.R., So, R., Malarvithi, P.P., Ladhaet, J.K and de Bruijn, F.J. (1997). Isolation of endophytic bacteria from rice and assesment of their potential for supplying rice with biologically fixed nitrogen. *Plant Soil.* 194:25-36
- Taechowisan, T and Lumyong, S. (2003). Activity of endophytic actinomycetes from roots of *Zingiber officinale* and *Alpinia galena* against phytopathogenic fungi. *Ann Microbiol.* 53, 291-298.
- Taechowisan, T., Peberdy, J. F and Lumyong, S. (2003). Isolation of endophytic actinomycetes from selected plants and their antifungal activity. *World J. Microbiol Biotechnol.* 19: 381-385.
- Kurth, F., Mailander, S., Bonn, M., Feldhahn, S., Herrmann, S., Buscot, F., Schrey, S. D and Takka, M. T. (2014). *Streptomyces*-induced resistance against oak powdery mildew involves host plant responses in defense, photosynthesis, and secondary metabolism pathways. *Mol Plant Micro Inter.* 27: 891-900
- Yu, H., Zhang, L., Li, L., Zheng, C and Guo, L. (2010). Recent developments and future prospects of antimicrobial metabolites produced by endophytes. *Microbiol Res.* 165: 437-449. doi: 10.1016/j.micres.2009.11.009
- Yuan, W. M. and Crawford, D. L. (1995). Characterization of *Streptomyces lydicus* WYEC108 as a potential biocontrol agent against fungal root and seed rots. *Appl. Environ. Microbiol.* 61: 3119-3128
- Zhao, S., Du, C. M and Tian, C.Y. (2012). Suppression of *Fusarium oxysporum* and induced resistance of plants involved in the biocontrol of Cucumber *Fusarium Wilt* by *Streptomyces bikiniensis* HD-087. *World J Microbiol Biotechnol.* 28: 2919-2927. doi:10.1007/s11274-012-1102-6

● **6% Overall Similarity**

Top sources found in the following databases:

- 6% Internet database
- 0% Publications database

TOP SOURCES

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

1	core.ac.uk Internet	1%
2	link.springer.com Internet	1%
3	pubmed.ncbi.nlm.nih.gov Internet	<1%
4	iopscience.iop.org Internet	<1%
5	tjnpr.org Internet	<1%
6	vestre.com Internet	<1%
7	frontiersin.org Internet	<1%
8	microbiologyresearch.org Internet	<1%
9	tandfonline.com Internet	<1%

● Excluded from Similarity Report

- Crossref database
- Submitted Works database
- Quoted material
- Small Matches (Less than 10 words)
- Crossref Posted Content database
- Bibliographic material
- Cited material
- Manually excluded sources

EXCLUDED SOURCES

eprints.unm.ac.id	46%
Internet	
envirobiotechjournals.com	6%
Internet	
envirobiotechjournals.com	6%
Internet	
pertanika.upm.edu.my	2%
Internet	
patents.google.com	2%
Internet	