

Usnic Acid Derivate from Usnea sp. and Bioactivity against Arthemisia salina Leach

by Muharram M

Submission date: 25-Dec-2020 09:37AM (UTC-0500)

Submission ID: 1481227760

File name: id_Derivate_from_Usnea_sp._and_Bioactivity_against_Arthemisia.docx (1.01M)

Word count: 2200

Character count: 11907

Usnic Acid Derivate from *Usnea sp.* and Bioactivity against *Arthemiasalina* Leach

MARYONO^{1,a}, MUHARRAM¹, A. IRMA Suryani², IWAN Dini¹

¹Jurusan Kimia, Fakultas Matematika dan Ilmu Pengetahuan Alam, Universitas Negeri Makassar
Jl. Daeng Tata Raya, Kampus UNM Parangtambung, Makassar 90244, Indonesia

²Jurusan Biologi, Fakultas Matematika dan Ilmu Pengetahuan Alam, Universitas Negeri Makassar
Jl. Daeng Tata Raya, Kampus UNM Parangtambung, Makassar 90244, Indonesia

^amaryounm76@gmail.com

Keywords: *Usnea sp.*, Usnic acid, *A. Salina* Leach.

Abstract. Usnic acid is chemical compound commonly found in lichen. These derivatives compounds both from the synthesis and from the isolation of various lichens have been studied as antibacterial, antifungal, antiparasitic, antiviral, and anticancer. Usnic acid derivative with not hydroxyl group on carbon C-3 was isolated from lichen *usnea sp.* taken from the Sinjaiselatan described in this paper. Its bioactivity against *Arthemiasalina* Leach. LC₅₀ value of 19.49 µg/mL.

Introduction

Usnic acid is a lichen metabolite was first isolated in 1844 and it's still studied because it has a high commercial value. Pharmacological research at the beginning of the 1950s antibiotic era proposed usnic acid as an active compound [1]. Further investigation shows usnic acid as a lichen secondary metabolite which has many interesting bioactivity properties for industrial and health purposes, especially those related to human diseases caused by microbes. Unique biological and physiological properties make these compounds and their derivatives have potential applications in clinical pharmacology [2]. Usnic acid is found in lichen, especially in the *Usnea*, and also in Lichen *Alectoria*, *Cladonia*, *Lecanora*, *Ramalina* and *Evernia* [3]. Compounds that are closely related to usnic acid are also found in fungi such as *phytoxin mycosuine* from the *Mycosphaerella nawae* fungus [4] and *cercosporamide* and *usnic acid amide* from the *Cercosporidium henningii* [5].

Usnic acid (2,6-diacetyl-7,9-dihydroxy-8,9b-dimethyl-1,3 (2H, 9bH) -dibenzo-furandione) is a general chemical component found in lichen cells and is a chemical component of secondary metabolites main finding in all species of *Usnea sp.* From the basic structure of the molecule, this compound is found in the form of a pair of enantiomers as optically active namely; (-) - Usnic acid and (+) - Usnic acid. Physically the two enantiomers have the same physical properties and basic structure as the molecular formula C₁₈H₁₆O₇, it is very easy to form yellow crystals with needle or prism forms, having a melting point of around 203-204 °C. Both enantiomers are distinguished from optical rotation which rotates the plane of right and left polarization depending on the polarization of the rotating plane of the methyl group on carbon 9b with each specific optical rotation $[\alpha]_D^{20} = 495^\circ$ and -495° [6]. In the practice of testing the results of isolation are not known for the mixture composition of the two. Different compositions or at a pure level will provide different biological activities [7].

Metabolism of usnic acid is known to be included in the *dibenzofurano* group biosynthesis pathway from polyketide through acetyl coenzyme A then cyclization to form *methylphloracetophenone* then dimerization with the C1 bonding then aromatization involving *stereospecific oxidative phenolic coupling* of two units *methylphloracetophenone* hydrated and then dehydrated to ether formation produce usnic acid. The relevance of the molecular structure changes in functional groups and derivatives of usnic acid has been validated and further explored in the investigation of the many biological properties for medicinal purposes demonstrating the broad biological and physiological activity relevant in the field of clinical pharmacology with antimicrobial activity pathogenic to humans and plants. It's showed antiviral, antiprotozoal, antiproliferative, anti-inflammatory and

analgesic activity also showed ecological effects such as grouter, antiherbivorous and insecticidal properties. Differences in biological activity in some cases have been observed between two forms

of enantiomer usnic acid. Recently health food supplements containing usnic acid have been promoted for use in weight reduction, with little scientific support. The current review emphasis is on the chemical and biological activity of usnic acid and its derivatives in addition to rational and ecologically acceptable methods for the supply of these natural compounds on a large scale [8].

Usnic acid as a pure substance has been applied in the form of cream formulations, toothpastes, mouthwashes, sunscreen products, and some cases as preservatives. At present, a review of usnic acid's biological and biological activities is needed and its derivatives are rational and can be accepted ecologically to provide usnic acid on a large scale [8]. Lichen as the main source of the discovery of usnic acid is widely found growing in the Sinjaiborong area of Sinjaiselatan district, South Sulawesi province, especially in high altitude areas with low humidity. In this study, usnea sp. obtained from the area, which is one of the lichen species found in many trees and branches of coffee plants. 1,8-diaminooctane compounds as derivative of usnic acid were evaluated on murine and human cancer cell lines. It showed significant cytotoxicity against L1210 cancer cells [9]. Therefore, usnic acid derivative it found in this research was tested for cytotoxicity against *Artemia salina* as a preliminary test for anticancer correlation.

Experimental

The plant material is usnea sp. taken from Sinjai Borong, South Sulawesi Province, Indonesia. Specimens were identified in Herbarium Bogoriensis LIPI Bogor document number FR-7.5.1PU.01-02.Ed. Organic solvents (E. Merck) n-hexane, chloroform, ethyl acetate, acetone, and methanol. Silica gel TLC plate GF₂₅₄, G 60 silica gel (70-230 mesh), catalog 1.07734, and CeSO₄ 10%, in 2N sulfuric acid. A total of 500 g of usnea sp. dry macerated with 6.5 liter chloroform previously with n-hexane, the obtained macerate was filtered and evaporated obtained 16 g extract. Chloroform extract was fractionated with chromatography column with the stationary phase are silica gel and mobile phase with n-hexane, ethyl acetate and methanol with gradient. Fractionation carryout six fraction (A-F). Continuously fractionating of D (2.54 g) with chromatography column, and with hexane: chloroform: ethyl acetate (2: 7: 1) as mobile phase and then crystallization of D1 obtained compound 2 (34 mg). Compound 2 was measured by melting point with kruss M5000 melting point, elucidation of the structure by Shimadzu FT-IR spectrophotometer prestige-21 KBr plate, 500NMR spectrometry Agilent (¹H at 500 MHz and ¹³C at 125 MHz). *Brine Shrimp Lethality Test* (BST) follows BN Meyer's procedure [10].

Results and Discussion

Structure elucidation: Compound 2 (Fig. 1) obtained as solid yellow crystals, melting point 198-200°C. IR (KBr) ν (cm⁻¹) 3438, 2918, 1708, 1614, and 1136. IR spectrum gives absorption wave numbers (ν , cm⁻¹) ν = 3438 cm⁻¹ as the OH group, ν = 2918 cm⁻¹ as the asymmetric stretching vibration CH, ν = 1707 cm⁻¹ as the vibration of the carbonyl group C=O, ν = 1614 cm⁻¹ is vibration of the C=C bond from aromatics, ν = 1136 cm⁻¹ as vibration of the COC. The spectrum representing the functional group in usnic acid.




Figure 1. Structure of compound 2

The spectrum of $^1\text{H-NMR}$ (Fig. 2) shows proton signals are indicating seven proton namely; four aliphatic protons δ_{H} (1.76 ppm (3H, s); 2.10 ppm (3H, s); 2.66 ppm (3H, s) and 2.68 ppm (3H,s) respectively as methyl proton at position 9b-CH₃, H8-CH₃, H-2-OCCH₃ and 6-OCCH₃. An aromatic proton δ_{H} 5.98 ppm (1H, s) at position 4, as well as two hydroxyl proton attached to the aromatic ring δ_{H} (13.31ppm (1H, s) and 11:02 ppm (1H, s). The $^{13}\text{C-NMR}$ spectrum showed 18 carbon signal; six aromatic carbon δ_{C} (155.34; 101.66; 164.02; 109.46; 157.64; 104.09), four-carbon olefin δ_{C} (105.37; 191.84; 98.46 ; 179.50), three carbonyl carbon δ_{C} (198.18; 201.89; 200.44), four aliphatic carbon methyl δ_{C} (28.03; 31.41; 7.68; 32.26), and one quaternary carbon δ_{C} (59.22).




Figure 2. Spectrum $^1\text{H-NMR}$ and $^{13}\text{C-NMR}$ of Compound 2

Carbon framework and the location of each proton are shown from the HMBC and HMQC correlation (Fig. 3). All correlation of HMBC, is shown below; proton 2-OCCH₃ δ H = 2.66 (s) with carbon C-2 (δ C = 105.37), 2-C=O (δ C = 201.89). Proton 4H δ H = 5.98 (s) with carbon C3 (δ C = 191.84), C4 (δ C = 98.46), and C9b (δ C = 59.22). Proton 6-OCCH₃ δ H = 2.68 (s) with carbon 6-CCO (δ C = 200.44), C-6 (δ C = 101.66). Proton 7-OH δ H = 13.31 (s) with carbon C7 (δ C = 164.02), C6 (δ C = 101.66), C8 (δ C = 109.46). Proton 9b-CH₃ δ H = 1.76 with carbon C1 (δ C = 198.18), C9b (δ C = 59.22), C9a (δ C = 104.09), and C4 (δ C = 98.46). Proton 8-CH₃ δ H = 7.68 with carbon C8 (δ C = 109.46), C9 (δ C = 157.64), C7 (δ C = 164.02). Proton 9-OH δ H = 11.02 with carbon C9 (δ C = 157.64), C9b (δ C = 59.22), C8 (δ C = 109.46). HMQC showed that the proton correlation with carbon are below; proton (A) 9b-CH₃ (δ H = 1.7) with C9b-C carbon δ H = 59.22; proton (B) 8-CH₃ (δ H = 2.10) with carbon C8-CH₃ δ H = 7.68; proton (C) 6-OCCH₃ (δ H = 2.68) with carbon 6-OC δ H = 200.44; proton (D) 2-OCCH₃ (δ H = 2.66) with carbon 2-OC δ H = 201.89 ppm).




Figure 3. HMBC and HMQC Correlation of Compound 2

Based interpretation of $^1\text{H-NMR}$, $^{13}\text{C-NMR}$, NMR-2D, IR, data and by comparing NMR spectrum data of usnic acid [11], the compound (1) was ascertained as usnat acid derivative (molecular structure as shown in Figure 1) which is minus the hydroxyl group on the C-3 atom because there is no aliphatic hydroxyl observed at chemical shift ≥ 17 ppm.

Bioactivity against *Arthemia Salina* Leach

Based on the LC_{50} cytotoxicity BST against *A. Salina*, the compound 2 be considered as moderately toxic considering to the 50% of lethality it caused at the concentration of 19.49 $\mu\text{g/mL}$, the data calculate showed in Fig. 4 and Table 1. This criterion is in agreement with the American National Cancer Institute that fixed the limitation of IC_{50} must be lower than 30 $\mu\text{g/mL}$. The compound 2 as moderately toxic it have correlation as anticancer [11]. Synthesis of several usnic acid derivate has catalytic activity of tyrosyl-DNA phosphodiesterase enzyme with IC_{50} 0.33-2.7 μM and low cytotoxicity to human cells. That derivate can be used for anticancer therapies that are more effective when combined with topoisomerase 1 inhibitors [12].

Table 1. The Data %death of *A. Salina* Leach. versus concentration of compound 2

Con. (ppm)	log con.	Number of <i>A. Salina</i> Leach				% death			probit
		Compound 2		Control		sample	control	% death corrected	
		first	death	first	death				
15.62	1.19	28.00	16.00	20.0	0.0	57.14	0.00	57.14	5.18
31.25	1.49	28.00	18.00	21.0	1.0	64.29	4.76	62.50	5.33
62.50	1.80	30.00	22.00	21.0	2.0	73.33	9.52	70.53	5.55
125.00	2.10	27.00	22.00	23.0	3.0	81.48	13.04	78.70	5.81
250.00	2.40	25.00	22.00	20.0	4.0	88.00	20.00	85.00	6.04
500.00	2.70	36.00	32.00	24.0	5.0	88.89	20.83	85.96	6.08
1000.00	3.00	35.00	35.00	22.0	8.0	100.00	36.36	100.00	8.09




Figure 4. Graph of probity %lethality versus \log_{10} concentration for chronic (24 hours) toxicities of compound 2

Summary

Based on the results of research that concluded that the usnat acid derivat have been isolated from the chloroform extract of *usnea sp.* which has no hydroxyl group on the carbon C3. The compound dissolved in DMSO 5% concentration showed cytotoxic properties against *Arthemiasalina* Leach. LC₅₀ value of 19.49 µg/mL.

Acknowledgment

Thanks are due to the Microbiology Laboratory of the Department of Biology, FMIPA UNM for the antibacterial test and the Ministry of Research, Technology and Higher Education for the research funding grant through the State University of Makassar LPM in 2018.

References

- [1] Michael Lauterwein, Margret Oethinger, Klaus Belsner, Thies Peters, Reinhard Marre. (1995), in vitro activities of the lichen secondary metabolites vulpinic acid, (1)-usnic acid, and (2)-usnic acid against aerobic and anaerobic microorganisms. *Antimicrobial Agents and Chemotherapy*, 39, 2541–2543,
- [2] Zhiheng Zhang, Yu Zheng, Yue Li, Hui Bai, Tianwen Ma, Xiaopeng Song, Jinghua Zhao, Li Gao. (2018), the effects of sodium usnic acid by topical application on skin wound healing in rats. *Biomedicine & Pharmacotherapy*, 97, 587–593,
- [3] Kristin Ingólfssdóttir, Gavin A.C. Chung, Vilhjálmur G. Skulason, Stefan R. Gissurarson, Margret Vilhelmsdóttir. (1998) antimycobacterial activity of lichen metabolites in vitro. *European Journal of Pharmaceutical Sciences*, 6, 141–144,
- [4] Sassa and Masayuki Igarashi. (1990) structures of (-)-mycousnine, (+)-isomycousnine and (+)-oxymycousnine, new usnic acid derivatives from *Phytopathogenic*. *Agric. Biol. Chem.*, 54, 2231–2237,
- [5] Michael A. Conover, Ronald Mierzwa, Arthur King, David Loebenberg, W. Robert bishop, Mohindar Puar, Mahesh Patel, Stephen J. Coval, Joseph Hershenhorn, Gary A. Strobel. 1992 usnic acid amide, a phytotoxin and antifungal agent from *Cercosporidium i-enningii*. *Phytochemistry*, 31, 2999–3001,
- [6] Huneck, S., Yoshimura, I.. (1996) identification of lichen substances, 2nd ed., Springer, Berlin, New York, London, Singapore, Tokyo,
- [7] Jonathan McConathy and Michael J. Owens. (2003), stereochemistry in drug action, *Journal of Clinical Psychiatry*, 5, 70–73
- [8] K. Ingólfssdóttir. (2002), Molecules of Interest Usnic acid, *Phytochemistry*, 61, 729–736,
- [9] Bazin Marc-Antoine, Bazin, Anne-Cecile Le Lamer, Jean-Guy Delcros, Isabelle Rouaud, Philippe Uriac, Joel Boustie, Jean-Charles Corbel, Sophie Tomasi. (2008), Synthesis and cytotoxic activities of usnic acid derivatives. *Bioorganic & Medicinal Chemistry*, 16 (14), 6860–6866,
- [10] B.N. Meyer, N.R. Ferrigni, J.E. Putnam, L.B. Jacobsen, D.E. Nicholas, J.L. McLaughlin. (1992), bhrine shrimp: a convenient general bioassay for active plant constituent, *Planta Medica*, 45, 31–34.
- [11] M.A. Rashida, U. M.A. Majida, M.A. Quader. (1999), Complete NMR assignments of (+)-usnic acid, *Fitoterapia*, 70, 113–115,
- [12] Dyrkheeva, N., Luzina, O., Filimonov, A., Zakharova, O., Ilina, E., Zakharenko, A., Lavrik, O. 2018. Inhibitory Effect of New Semisynthetic Usnic Acid Derivatives on Human Tyrosyl-DNA Phosphodiesterase 1. *Planta Med.* doi:10.1055/a-0681-7069.

Usnic Acid Derivate from Usnea sp. and Bioactivity against *Arthemia salina* Leach

ORIGINALITY REPORT

10%

SIMILARITY INDEX

7%

INTERNET SOURCES

7%

PUBLICATIONS

%

STUDENT PAPERS

PRIMARY SOURCES

- 1 John A. G. Roach, Steven M. Musser, Kim Morehouse, Jason Y. J. Woo. "Determination of Usnic Acid in Lichen Toxic to Elk by Liquid Chromatography with Ultraviolet and Tandem Mass Spectrometry Detection", *Journal of Agricultural and Food Chemistry*, 2006
Publication 1%
- 2 eprints.unm.ac.id
Internet Source 1%
- 3 Francisco A. Durazo. "Fulminant Liver Failure Due to Usnic Acid for Weight Loss", *The American Journal of Gastroenterology*, 5/2004
Publication 1%
- 4 doi.org
Internet Source 1%
- 5 Baldenebro-López, Jesús, José Castorena-González, Norma Flores-Holguín, Jorge Almaral-Sánchez, and Daniel Glossman-Mitnik. "Computational Molecular Nanoscience Study of

the Properties of Copper Complexes for Dye-Sensitized Solar Cells", International Journal of Molecular Sciences, 2012.

Publication

6	hal.archives-ouvertes.fr Internet Source	1%
7	Zhiheng Zhang, Yu Zheng, Yue Li, Hui Bai, Tianwen Ma, Xiaopeng Song, Jinghua Zhao, Li Gao. "The effects of sodium usnic acid by topical application on skin wound healing in rats", Biomedicine & Pharmacotherapy, 2018 Publication	1%
8	Franck-Neumann, M.. "(@h^3: @h^1 allylcarbonyl)iron tricarbonyl complexes. I. Transformation of complexes from electrophilic gem-dimethylcyclopropenes into functionalized isoprenic diene-iron tricarbonyl complexes by thermolysis or photolysis", Tetrahedron Letters, 1981 Publication	1%
9	ijpbs.net Internet Source	1%
10	eprints.usm.my Internet Source	<1%
11	ntp.niehs.nih.gov Internet Source	<1%

12

repositorio.ufpe.br

Internet Source

<1%

13

femsle.oxfordjournals.org

Internet Source

<1%

14

portalgaruda.org

Internet Source

<1%

15

www.jurnal.unsyiah.ac.id

Internet Source

<1%

16

Jeng-Sheng Huang. "Plant Pathogenesis and Resistance", Springer Science and Business Media LLC, 2001

Publication

<1%

17

James P. Kutney, Ignacio H. Sanchez. "Studies in the usnic acid series. I. The condensation of (+)-usnic acid with aliphatic and aromatic amines", Canadian Journal of Chemistry, 1976

Publication

<1%

Exclude quotes

Off

Exclude matches

Off

Exclude bibliography

On

Usnic Acid Derivate from Usnea sp. and Bioactivity against Arthemia salina Leach

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6
