

Volume 35 Number 9 May 2021

ISSN 1478-6419


Natural Product Research

Part B – Bioactive Natural Products

Editor-in-Chief
Armandodoriano Bianco



Taylor & Francis
Taylor & Francis Group





Natural Product Research

Publish
with us

[Submit an
article](#) ▼

[About this
journal](#) ▼

Explore

[Browse all
articles & issues](#)
▼



Subscribe

[Alerts &
RSS feed](#) ▼

[+ Purchase a
subscription](#)

Ready to submit?

Start a new manuscript submission or continue a submission in progress

[Go to submission site](#) ↗

Submission information

[▶ Instructions for authors](#)

[▶ Editorial policies](#) ↗

Editing services

[▶ Editing services site](#) ↗

About this journal

[▶ Journal metrics](#)

[▶ Aims & scope](#)

[▶ Journal information](#)

[▶ Editorial board](#)

[▶ News & call for papers](#)

[▶ Advertising information](#)

Editorial board

Editor-in-Chief:

Armandodoriano Bianco - Dipartimento di Chimica, Università degli Studi di Roma "La Sapienza", Piazzale Aldo Moro 5, 00185 Roma, Italy

Editors:

Søren Rosendal Jensen - DTU Chemistry, Department of Chemistry, Emeriti, Organic Chemistry, DK-2820 Lyngby, Denmark

Marcello Nicoletti - Facoltà di Farmacia e Medicina, Università degli Studi di Roma "La Sapienza", Piazzale Aldo Moro 5, 00185 Roma, Italy

Editorial Office:

Dr Ilaria Serafini - Dipartimento di Chimica, Università degli Studi di Roma "La Sapienza", Piazzale Aldo Moro 5, 00185 Roma, Italy

Alessandro Venditti - Dipartimento di Chimica, Università degli Studi di Roma "La Sapienza", Piazzale Aldo Moro 5, 00185 Roma, Italy

Dr Claudio Frezza - Dipartimento di Biologia Vegetale, Università degli Studi di Roma "La Sapienza", Piazzale Aldo Moro 5, 00185 Roma, Italy

Associate Editors:

D. J. Aitken - Institute of Molecular Sciences of Orsay, France

B Charlwood - Universidade Federal de Alagoas, Campus Universitario, Brazil

P. Ciminiello - Università di Napoli "Federico II", Italy

L. N. Mander - The Australian National University, Australia

K. Mori - University of Tokyo, Japan

I. Ninomiya - Kobe Pharmaceutical University, Japan

R. J. Parry - Rice University, Texas

L. Pistelli - Università di Pisa, Italy

O. Sterner - Lund University, Sweden

J Stöckigt - Johannes Gutenberg University, Germany

Editorial Board:

G. Appendino - Italy

Y. Asakawa - Japan
R. Ascrizzi - Italy
M. Ballero - Italy
R. Bernini - Italy
E. Block - USA
E. Brasili - Italy
M. Bruno - Italy
H. Budzikiewicz - Germany
S. D. Burke - USA
Z. Candan - Turkey
L. Castedo - Spain
N. Cicero - Italy
G. Cimino - Italy
G. A. Cordell - USA
P. Cortis - Italy
B. Danieli - Italy
S. Danishefsky - USA
V. De Feo - Italy
D. De Vita - Italy
I. Fleming - UK
S. Foddai - Italy
B. Fraser-Reid - USA
C. Frezza - Italy
F. R. Grippaudo - Italy
F. Guarino - Italy
L. Gunatilaka - USA
G. M. Halpern - China
S. Hanessian - Canada
J. R. Hanson - UK
K. Hostettmann - Switzerland
M. Isobe - Japan
D. Kinghorn - USA
J. Lehn - France
P. W. Le Quesne - USA
S. V. Ley - UK
Y. Liu - China
P. D. Magnus - USA
K. Nakanishi - USA

R. Noyori - Japan
S. Persechino - Italy
J. M. Pezzuto - USA
G. A. M. Pintore - Italy
F. Poli - Italy
M. Rizk - Egypt
E. Rodriguez - USA
D. S. Rycroft - UK
A. Salvo - Italy
C. Sanna - Italy
F. Sciubba - Italy
B. Sener - Turkey
M. Serafini - Italy
L. Y. Sheen - Taiwan
G. Solladie - France
D. Spitzner - Germany
C. Toniolo - Italy
G. Topcu - Turkey
E. Tramontano - Italy
B. M. Trost - USA
R. Verpoorte - Netherlands
H. Wagner - Germany
P. G. Waterman - Australia
J. D. White - USA
K. R. Zeller - Germany





Virulence
Find the latest research

 [Learn More](#) 

This banner features a background of microscopic images in shades of purple and red. The text is white on a dark blue background. A small circular logo with a white icon is in the bottom left corner.



RNA Biology
Find the latest RNA research

 [Learn More](#) 

This banner features a background image of a microscope. The text is white on a dark blue background. A small circular logo with a white icon is in the bottom left corner.



Pharmaceutical Biology
A Taylor & Francis journal

Natural medicines, bioactive natural products,
and complex traditional medicine formulas

[Start your submission today](#) 

This banner features a background image of laboratory equipment. The text is white on a dark blue background. A small circular logo with a white icon is in the bottom left corner.

Information for

[Authors](#)[R&D professionals](#)[Editors](#)[Librarians](#)[Societies](#)

Opportunities

[Reprints and e-prints](#)[Advertising solutions](#)[Accelerated publication](#)[Corporate access solutions](#)

Open access

[Overview](#)[Open journals](#)[Open Select](#)[Dove Medical Press](#)[F1000Research](#)

Help and information

[Help and contact](#)[Newsroom](#)[All journals](#)[Books](#)

Keep up to date

Register to receive personalised research and resources by email



Sign me up



Copyright © 2022 Informa UK Limited

[Privacy policy](#)

[Cookies](#)

[Terms &](#)

[conditions](#) [Accessibility](#)



Taylor & Francis Group
an informa business

Registered in England & Wales No. 3099067

5 Howick Place | London | SW1P 1WG



[Home](#) ▶ [All Journals](#) ▶ [Natural Product Research](#) ▶ [List of Issues](#) ▶ [Volume 35, Issue 9](#)

Natural Product Research

Publish
with us

[Submit an article](#) ▼

[About this journal](#) ▼

Explore


[Browse all articles & issues](#) ▼

 [Latest issue](#)

Subscribe

[Alerts & RSS feed](#) ▼

[+ Purchase a subscription](#)



Browse this journal

[▶ Latest articles](#)

[▶ Current issue](#)

[▶ List of issues](#)

[▶ Special issues](#)

[▶ Open access articles](#)

[▶ Most read articles](#)

[▶ Most cited articles](#)

Sample our
Bioscience
Journals




>> [Sign in here](#) to start your access
to the latest two volumes for 14 days

Natural Product Research, Volume 35, Issue 9 (2021)

[See all volumes and issues](#)

[<](#)
[Volume 35, 2021](#)
[Vol 34, 2020](#)
[Vol 33, 2019](#)
[>](#)

[<](#)
[Volume 35, Issue 4](#)
[Volume 35, Issue 3](#)
[Volume 35, Issue 2](#)
[Volume 35, Issue 1](#)
[>](#)


[Browse by section \(All\)](#)


[Display order \(Default\)](#)

[Download citations](#)
 [Download PDFs](#)


Research Articles

Article
[Sesquiterpenoids from *Saussurea costus*](#) >

Tran Thi Hong Hanh, Pham Thi Cham, Nguyen Thi Thuy My, Nguyen The Cuong, Nguyen Hai Dang, Tran Hong Quang, Tran Thu Huong, Nguyen Xuan Cuong, Nguyen Hoai Nam & Chau Van Minh

Pages: 1399-1405

Published online: 12 Aug 2019



326	6	0
Views	CrossRef citations	Altmetric


Article

New naphthoquinone derivatives from *Fusarium napiforme* of a mangrove plant >

Unang Supratman, Nami Hirai, Shiho Sato, Keiko Watanabe, Abdul Malik, Suwardi Annas, Desi Harneti, Rani Maharani, Takuya Koseki & Yoshihito Shiono

Pages: 1406-1412

Published online: 12 Aug 2019



Phytotoxic action in lettuce seeding

235	9	1
Views	CrossRef citations	Altmetric


Article


An unambiguous and practical synthesis of Oroxylin A: a commonly misidentified flavone >

Hosahalli P. Hemantha, Rajendran Ramanujam, Muhammed Majeed & Kalyanam Nagabhushanam

Pages: 1413-1420

Published online: 12 Aug 2019






182	3	0
Views	CrossRef citations	Altmetric

Article

Three new bioactive diterpenoids from the roots of *Croton crassifolius* >

Guo-Hua Ye, Jun-Juan Xue, Wei-Ling Liang & Shou-Juan Yang

Pages: 1421-1427

Published online: 14 Aug 2019

238	0	0
Views	CrossRef citations	Altmetric


Article

α -Coticic acid, a plant sesquiterpene with acaricidal activity against *Varroa destructor* parasitizing the honey bee >

Alessio Cimmino, Fabrizio Freda, Ernesto Santoro, Stefano Superchi, Antonio Evidente, Massimo Cristofaro & Marco Masi

Pages: 1428-1435

Published online: 16 Aug 2019



320	9	1
Views	CrossRef citations	Altmetric


Article

Pentacyclic triterpenes from *Lavandula coronopifolia*: structure related inhibitory activity on α -glucosidase >

Marwa Elsbaey, Rogers Mwakalukwa, Kuniyoshi Shimizu & Tomofumi Miyamoto

Pages: 1436-1444

Published online: 21 Aug 2019



157	3	0
Views	CrossRef citations	Altmetric


Article

Two new dihydroisocoumarins with antimicrobial activities from the fungus *Penicillium* sp. XR046 collected from Xinren coal area >

Guo-Bo Xu, Fei-Yu Yang, Xin-Yu Wu, Rui Li, Meng Zhou, Bing Wang, Xiao-Sheng Yang, Ting-Ting Zhang & Shang-Gao Liao

Pages: 1445-1451

Published online: 28 Aug 2019



189	2	0
Views	CrossRef citations	Altmetric


Article

Anti-inflammatory glycosides from the roots of *Paeonia intermedia* C. A. Meyer >

Liang Yu, Ling-Juan Zhu, An-Hua Wang, Yu Qin, Xue Zhang, Jing-Ming Jia & Xin-Sheng Yao

Pages: 1452-1458

Published online: 27 Aug 2019



171	0	0
Views	CrossRef citations	Altmetric


Article

A new isoflavone glycoside from flowers of *Pueraria Montana* var. *lobata* (Willd.) Sanjappa & Pradeep >

Yilong Ma, Yafang Shang, Zhifeng Zhong, Yingshuo Zhang, Yang Yang, Jun Feng & Zhaojun Wei

Pages: 1459-1464

Published online: 21 Aug 2019





179	3	0
Views	CrossRef citations	Altmetric


Article

Two novel diterpenes from the stems and leaves of tropical seagrass *Enhalus acoroides* in the South China sea >


Xiao-Bing Wang, Zhong-Hao Sun, Li-Xia Fan, Yang-Yang Liu, Jian Feng, Guo-Xu Ma & De-Li Chen

Pages: 1465-1473

Published online: 20 Aug 2019



Enhoidin A (1)



Enhoidin B (2)

161	1	0
Views	CrossRef citations	Altmetric

Article

Isolation and structural determination of flavan-3-ol derivatives from the *Polypodium vulgare* L. rhizomes water extract >

Michał Gleńsk, Marta K. Dudek, Maciej Ciach & Maciej Włodarczyk

Pages: 1474-1483

Published online: 21 Aug 2019



336	8	0
Views	CrossRef citations	Altmetric

Article




Design and synthesis of novel hydrocarbylidene nitrohydrazinecarboximidamides against aphids based on natural product galegine >

Qinglai Wu, Fenghai Zhao, Xiang Zhu, Ziyuan Zhou, Dongyan Yang, Zhihong Xu, Junkai Li & Zhaohai Qin

Pages: 1484-1490

Published online: 20 Aug 2019

Galega officinalis L.



Galegine

IIa-III

118	2	0
Views	CrossRef citations	Altmetric


Article

[A new chlorine-containing iridoid glycoside from *Plantago maxima* >](#)

Liang-Liang Ma, Lu-Lu Wang, Yue-Feng Zhang, Xiang-Fu Jiang, Xin-Liu Zhu, Ke Pan, Chuan-Xing Wan & Zhong-Bo Zhou

Pages: 1491-1496

Published online: 20 Aug 2019



140	1	0
Views	CrossRef citations	Altmetric


Article

[A pair of novel bisindole alkaloid enantiomers from marine fungus *Fusarium* sp. XBB-9 >](#)

Sharpkate Shaker, Run-Zhu Fan, Hou-Jin Li & Wen-Jian Lan

Pages: 1497-1503

Published online: 22 Aug 2019



489	7	0
Views	CrossRef citations	Altmetric

Article


Secondary metabolites from the marine-derived fungus *Phaeosphaeria spartinae* >

Mahmoud Fahmi Elsebai

Pages: 1504-1509

Published online: 23 Aug 2019





215	0	0
Views	CrossRef citations	Altmetric



Article

Isolation and characterization of *ent*-pimarane diterpenoids from *Sigesbeckia pubescens* >

Jianbin Wang, Kehui Xie, Qing Wang, Wenchao Li & Hongzheng Fu

Pages: 1510-1517

Published online: 10 Sep 2019



269	1	0
Views	CrossRef citations	Altmetric


Article

Vitexnegheteroin M, a new phenolic glycoside from *Vitex negundo* var. *heterophylla* >

Ye Huang, Guo-Yu Ding & Ping Hu

Pages: 1518-1524

Published online: 26 Aug 2019



vitexnegheteroin M (1)

Vitex negundo var. *heterophylla*

187	4	0
Views	CrossRef citations	Altmetric


Article

New metabolites produced by endophyte *Clonostachys rosea* B5-2 >

Unang Supratman, Takuma Suzuki, Tomoki Nakamura, Yasuhiro Yokoyama, Desi Harneti, Rani Maharani, Supriatno Salam, Fajar Fauzi Abdullah, Takuya Koseki & Yoshihito Shiono

Pages: 1525-1531

Published online: 27 Aug 2019



450	16	2
Views	CrossRef citations	Altmetric


Article

Rice germ macro- and micronutrients: a new opportunity for the nutraceutics >

Mariangela Rondanelli, Alessandra Miccono, Gabriella Peroni, Mara Nichetti, Vittoria Infantino, Daniele Spadaccini, Tariq A. Alalwan, Milena Anna Faliva & Simone Perna

Pages: 1532-1536

Published online: 03 Sep 2019





- Good lipid intake
- Good fiber intake
- High content of thiamine (B1) and pyridoxine (B6)
- Iron (77% of RDA) and magnesium (108% of RDA)

233	5	0
Views	CrossRef citations	Altmetric



Article

Sesquiterpenoids from *Pilea cavalieriei* >

Yong Zhou, Ling-Yu Li, Xu Yin, Qing-Ying Zhang, Hong Liang & Peng-Fei Tu

Pages: 1537-1543

Published online: 05 Sep 2019



200	0	0
Views	CrossRef citations	Altmetric


Article



Iridoid compounds from the aerial parts of *Swertia mussotii* Franch. with cytotoxic activity >

Haixia Wang, Huiming Huang, Jingwei Lv, Ning Jiang, Yujiao Li, Xinmin Liu & Hongping Zhao

Pages: 1544-1549

Published online: 06 Sep 2019





184

Views

2

CrossRef citations

0

Altmetric

Short Communications


- Rapid Communication
Chemical composition and anti-inflammatory activity of the essential oil from the leaves of *Limnocitrus littoralis* (Miq.) Swingle from Vietnam >

Tuan Quoc Doan, Duc Viet Ho, Nhan Trong Le, Anh Tuan Le, Kiem Van Phan, Hoai Thi Nguyen & Ain Raal

Pages: 1550-1554

Published online: 20 Aug 2019

Limnocitrus littoralis leaves from Vietnam



200

Views

4

CrossRef citations

3

Altmetric

- Rapid Communication
Ustilaginoidin M₁, a new bis-naphtho- γ -pyrone from the fungus *Villosiclava virens* >


Jiajia Meng, Siji Zhao, Pengqin Dang, Zhiyao Zhou, Daowan Lai & Ligang Zhou

Pages: 1555-1560

Published online: 12 Aug 2019



***Villosiclava virens* UV-8b**



230	7	0
Views	CrossRef citations	Altmetric


Rapid Communication

Chemical Investigation Of Secondary Metabolites Produced By Mangrove Endophytic Fungus *Phyllosticta Capitalensis* >

Zhiyong Xu, Bingxue Xiong & Jing Xu

Pages: 1561-1565

Published online: 23 Aug 2019



436	11	1
Views	CrossRef citations	Altmetric


Rapid Communication

Coumarin (2H-1-benzopyran-2-one): a novel and eco-friendly aphicide >

Roman Pavela, Filippo Maggi & Giovanni Benelli

Pages: 1566-1571

Published online: 11 Sep 2019



308

Views

1

CrossRef citations

1

Altmetric

Explore articles

Latest

Open access

Most read

Most cited

Trending

Synthesis of ring-A modified lupeol derivatives and their anti-proliferative activity: identification of potent lead active against MCF-7 breast cancer cells >

Kummari Narendar et al.

Article | Published online: 9 Sep 2022

Structural characterization of a pectic polysaccharide from laoshan green

[tea and its inhibitory effects on the production of NO, TNF- \$\alpha\$ and IL-6 >](#)

Fan Liu et al.

Article | Published online: 9 Sep 2022

[Flavonoids composition, antioxidant, anticholinesterase, and cytotoxic activities of flowers extract of *Senna cearensis* Afr. Fern >](#)

Maria G. A. Franca et al.

Brief Report | Published online: 9 Sep 2022

[Bioactive compounds of the *Opuntia monacantha* fruit >](#)

Vinicius M. da Silva et al.

Brief Report | Published online: 8 Sep 2022

[View all latest articles >](#)

Information for

- Authors
- R&D professionals
- Editors
- Librarians
- Societies

Opportunities

- Reprints and e-prints
- Advertising solutions
- Accelerated publication
- Corporate access solutions

Open access

- Overview
- Open journals
- Open Select
- Dove Medical Press
- F1000Research

Help and information

- Help and contact
- Newsroom
- All journals
- Books

Keep up to date

Register to receive personalised research and resources by email


 Sign me up



Copyright © 2022 Informa UK Limited Privacy policy Cookies Terms & conditions Accessibility



Registered in England & Wales No. 3099067
5 Howick Place | London | SW1P 1WG



Natural Product Research

Formerly Natural Product Letters



ISSN: 1478-6419 (Print) 1478-6427 (Online) Journal homepage: <https://www.tandfonline.com/loi/gnpl20>


New naphthoquinone derivatives from *Fusarium napiforme* of a mangrove plant

Unang Supratman, Nami Hirai, Shiho Sato, Keiko Watanabe, Abdul Malik, Suwardi Annas, Desi Harneti, Rani Maharani, Takuya Koseki & Yoshihito Shiono

To cite this article: Unang Supratman, Nami Hirai, Shiho Sato, Keiko Watanabe, Abdul Malik, Suwardi Annas, Desi Harneti, Rani Maharani, Takuya Koseki & Yoshihito Shiono (2019): New naphthoquinone derivatives from *Fusarium napiforme* of a mangrove plant, Natural Product Research, DOI: [10.1080/14786419.2019.1650358](https://doi.org/10.1080/14786419.2019.1650358)

To link to this article: <https://doi.org/10.1080/14786419.2019.1650358>

 View supplementary material 

 Published online: 12 Aug 2019.

 Submit your article to this journal 

 View Crossmark data 



New naphthoquinone derivatives from *Fusarium napiforme* of a mangrove plant

Unang Supratman^a, Nami Hirai^b, Shiho Sato^b, Keiko Watanabe^b, **Abdul Malik^c**,
Suardi Annas^d, Desi Harneti^a, Rani Maharani^a, Takuya Koseki^b and
Yoshihito Shiono^b

^aDepartment of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Padjadjaran, Sumedang, Indonesia; ^bDepartment of Food, Life, and Environmental Science, Faculty of Agriculture, Yamagata University, Tsuruoka, Yamagata, Japan; ^cDepartment of Geography, Faculty of Mathematics and Natural Sciences, Universitas Negeri Makassar, Makassar, Indonesia; ^dDepartment of Statistics, Faculty of Mathematics and Natural Sciences, Universitas Negeri Makassar, Makassar, Indonesia

ABSTRACT


Two new naphthoquinone derivatives, 6-hydroxy-astropaquinone B (**1**) and astropaquinone D (**2**) as well as the known compound 3-*O*-methyl-9-*O*-methylfusarubin (**3**) were isolated from *Fusarium napiforme*, an endophytic fungus isolated from the mangrove plant, *Rhizophora mucronata*. The structures of **1** and **2** were determined by 1D and 2D NMR spectroscopic analyses. Compounds **1**, **2** and **3** exhibited moderate antibacterial activity against *Staphylococcus aureus* and *Pseudomonas aeruginosa*. Furthermore, **1**, **2** and **3** were phytotoxic action in lettuce seeding at a concentration of 30 $\mu\text{g} \cdot \text{mL}^{-1}$.

ARTICLE HISTORY

Received 26 May 2019
Accepted 21 July 2019

KEYWORDS

Endophyte; naphthoquinone; 6-hydroxy-astropaquinone B; astropaquinone D; antimicrobial activity; phytotoxicity



Phytotoxic action in lettuce seeding



1. Introduction

Endophytic fungi, which asymptotically inhabit plant tissues, may produce specific biologically active secondary metabolites in some cases. Actually, endophytes have also been a prominent source of new secondary metabolites with unique structures over past two decades. In addition, endophyte can evolve to produce the same or similar bioactive compounds as their host plants (Venugopalan and Srivastava 2015). Further, to date, several novel endophytic fungal-derived natural products have been isolated and described (Li et al. 2018). A previous investigation by our team has demonstrated that several secondary metabolites secreted by endophytic fungi isolated from different habitats were biologically active and had interesting structures (Shiono et al. 2016).

In our continuing search for fungal metabolites, a batch of approximately 50 fungal strains isolated from mangrove plants collected from Makassar, South Sulawesi, Indonesia, were fermented on a small scale. Subsequently, the methanolic fermentation extracts were subjected to TLC chemical screening. The fungal strain *F. napiforme* IP-28 isolated from the stem of *Rhizophora mucronata* was selected for further chemical investigation. The genus *Fusarium* is known as one of the mycotoxin producing fungi with diverse structures (Zakaria 2017). *F. napiforme* has been firstly isolated from millet and sorghum from Southern Africa, and soil debris from one grassland in Australia (Marasas et al. 1987) and ability to produce fumonisin B1 has been reported (Zakaria 2017). Fungal culture of IP-28 strain was extracted with solvents and fractionated using different chromatographic techniques, which yielded two new compounds and one previously identified compound. Based on different spectroscopic data including HRESITOFMS, as well as ^1H , ^{13}C , and 2D NMR, the structures of the new compounds were found to be naphthoquinone derivatives, which were named 6-hydroxy-astropaquinone B (**1**) and astropaquinone D (**2**). We found that **1** and **2** exhibited antimicrobial activities against *S. aureus* and *P. aeruginosa*. Additionally, all three compounds regulated the growth of lettuce seedlings.

2. Result and discussion

Compound **1** was obtained as a red amorphous powder. The molecular formula of **1** was determined as $\text{C}_{17}\text{H}_{18}\text{O}_7$ by HRESITOFMS, in combination with ^{13}C NMR spectroscopy. The IR spectrum showed major absorption bands at 3397 and 1650 cm^{-1} corresponding to a hydroxyl and a carbonyl group, respectively. The UV absorption bands at 226, 283 and 491 nm together with IR absorption bands indicated the presence of a 1,4-naphthoquinone moiety in **1**. The ^{13}C NMR spectrum of **1**, when analysed in combination with the HSQC data, revealed the presence of two quinone carbonyls (δ_{C} : 190.3 and 179.5), eight aromatic carbons, of which three were oxygenated (δ_{C} : 148.4, 156.0 and 155.5) and one was protonated (δ_{C} : 102.8), two oxygenated methine carbons, of which one was dioxygenated (δ_{C} : 94.1), three OCH_3 carbons (δ_{C} : 56.4, 56.5 and 56.8), one methylene carbon (δ_{C} : 28.6), and a methyl carbon (δ_{C} : 21.1). The ^1H NMR spectrum of **1** showed signals for one aromatic proton [δ_{H} : 6.74 (1 H, s, H-8)], two methine protons [δ_{H} : 4.21 (1 H, m, H-3), 5.52 (1 H, s, H-1)], three methoxy groups [δ_{C} : 56.4 (q, 1-OMe), 56.5 (q, 7-OMe), 56.8 (q, 7-OMe)], one doublet methyl group [δ_{H} : 1.38

(3 H, d, $J=6.7$ Hz, H-11)], and one methylene group [δ_{H} : 2.22 (1 H, dd, $J=18.0, 10.0$ Hz, H-4), 2.66 (1 H, dd, $J=18.0, 3.5$ Hz, H-4)]. The cross peaks between H-4/H-3/H-11 were found in the ^1H - ^1H COSY spectrum of **1** (Supplementary information Figure S1). The HMBC correlations from H-8 to C-6, C-9a, and C-10, from H-1 to C-10, from H-4 to C-5 and from H-11 to C-3 and C-4 established a methyl-pyranonaphthoquinone nucleus (Supplementary information Figure S1). In addition, NOE correlations between H-8 and two methoxys at C-7 and C-9 were found, which established the positions of the methoxyl groups at C-7 and C-9 in the aromatic ring (Supplementary information Figure S1). The location of a methoxyl group at C-1 of the pyran ring was indicated by HMBC correlations between these protons and the corresponding carbon (Supplementary information Figure S1). Comparison of the C-6 ^{13}C NMR chemical shifts (δ_{C} : 148.4) of **1** with that for the corresponding C-atom (δ_{C} : 148.4) of **3** (Kornsakulkarn et al., 2011) revealed the position of the hydroxyl group at C-6. Thus, the planar structure of **1** was assigned as depicted in Figure 1. Compound **1** is structurally related to astropaquinones B (**4**) and C (**5**) previously isolated from *Astrosphaeriella papuana* (Geng et al. 2012; Wang et al. 2009). The relative configuration of **1** was deduced on the basis of NOE experiments (Supplementary information Figure S2). The NOE correlation of OMe-1/H-3 suggested that its relative configuration was identical to that reported for **4**. In addition, the specific optical rotation of **1** exhibited a positive specific rotation value [$[\alpha]_{\text{D}} +45.1^\circ$ (c 0.12, CHCl_3)] which was similar to the previously described **4**. Thus, the structure of **1** was identified as (1*R*, 3*S*)-6-hydroxy-astropaquinone B.

Compound **2** was obtained as an orange amorphous powder. The molecular formula of **2** was determined as $\text{C}_{16}\text{H}_{16}\text{O}_6$ by a combination of its HRESITOFMS, ^{13}C NMR, and HSQC data, which also indicated that the molecule has nine degrees of unsaturation. Its UV spectrum exhibited absorption maxima at 280 and 490 nm, which is characteristic of a pyranonaphthoquinone (Masi et al. 2018). Furthermore, its IR spectrum showed absorption bands at 3502 and 1631 cm^{-1} , indicating the presence of hydroxy and quinone ketone functions. The ^1H NMR data of **2** indicated that it was structurally related to **1**. The essential difference between the ^1H NMR spectra of **1** and **2** was the absence of a C-1 methoxy group in **1**, which instead contained a methylene group [δ_{H} : 4.50 (1H, dt, $J=19.0, 2.0$ Hz, H-1), 4.86 (1H, dd, $J=19.0, 2.4$ Hz, H-1)]. Correlations from H-1 to C-3 and C-4a in the HMBC spectrum of **2** confirmed that a methylene moiety was located at C-1. In addition, the specific optical rotation of **2** exhibited a positive specific rotation value [$[\alpha]_{\text{D}} +100^\circ$ (c 0.3, MeOH)] similar to that of the




Figure 1. Chemical structures of 1–5.

previously described synthetic compound, (+)-(3S)-7,9-dimethoxy-3-methyl-3,4-dihydro-1H-benzo[*g*]isochromene-5,10-dione $[[\alpha]_D^{25} +112.5^\circ$ (c 1.3, CH_2Cl_2)] (Bulbule et al. 2007). Thus, the structure of **2** was established and given the trivial name astropaquinone D.

The structure of **3** was identified as 3-*O*-methyl-9-*O*-methylfusarubin by comparing our physical and spectroscopic data with the reported data (Tatum et al. 1985).

Naphthoquinones isolated from fungi have been reported to possess antimicrobial activity against bacteria (Wang et al. 2009). The antimicrobial mechanism involves the generation of superoxide anion and hydrogen peroxide in bacterial membranes. Consequently, superoxide anion and hydrogen peroxide can induce damage to bacterial membranes, proteins and DNA, which induces apoptosis (Ravichandiran et al. 2019). Compounds **1–3** were evaluated for antimicrobial activity against Gram-positive and Gram-negative bacteria, yeast, and other fungal strains. Compounds **1**, **2** and **3** showed moderate antibacterial activity against *S. aureus* NBRC 13276 and *P. aeruginosa* ATCC 15442 [MIC values ($\mu\text{g} \cdot \text{mL}^{-1}$): **1**: 6.3, **2**: 12.5, **3**: 6.3 for *S. aureus* NBRC 13276, and **1**: 6.3, **2**: 6.3, **3**: 6.3, for *P. aeruginosa* ATCC 15442]. In contrast, neither *Aspergillus clavatus* F 318a, nor *Candida albicans* ATCC 2019 were inhibited by **1**, **2** or **3** (at $25 \mu\text{g} \cdot \text{mL}^{-1}$). It has been reported that **4** and **5** exhibited antimicrobial activity against *S. aureus* YMF 3.17 with MIC values ($\mu\text{g} \cdot \text{mL}^{-1}$) of 20 and 10, respectively. A comparison of the published MIC values for *S. aureus* to those of the present study suggested that **1** and **2** might be more potent inhibitors than **4** and **5**. Higher activity was observed when the structure had the naphthoquinone unit with a hydroxy at C-6. Since some endophytes produce biological active mycotoxins in the plant tissues, and the accumulation of these mycotoxins often enhance the phytotoxicity or symptoms of plant diseases (Ismail and Papenbrock 2015), the phytotoxic activities of **1**, **2** and **3** were investigated by seed germination test on lettuce (*Lactuca sativa* L.) with 2,4-dichlorophenoxyacetic acid ($0.3 \mu\text{g} \cdot \text{mL}^{-1}$) as the positive control (Supplementary Information Figure S16). Compounds **1**, **2** and **3** each inhibited the growth of both roots and hypocotyls at $30 \mu\text{g} \cdot \text{mL}^{-1}$. Furthermore, **1** suppressed seed germination at $100 \mu\text{g} \cdot \text{mL}^{-1}$.

3. Experimental

3.1. General experimental procedures

See detailed experimental section in [supplementary material](#).

3.2 Fungal material and fermentation

See detailed experimental section in [supplementary material](#).

3.3. Extraction and purification

The extract was separated by silica gel column chromatography with a stepwise elution of *n*-hexane/EtOAc (100:0–0:100) to give 11 fractions (Frs.1-1 to 1-11). Frs. 1-4 and 1-5 (2.1 g) were further separated on a silica gel column with $\text{CHCl}_3/\text{EtOAc}$ to provide

11 fractions (Frs. 2-1 to 2-11). Fr. 2-4 was chromatographed on silica gel CC using a stepwise gradient of CHCl_3 -EtOAc to give fractions 1 – 12 (Frs. 2-4-1 to 2-4-12). Fr. 2-4-5 (150 mg) was subjected to ODS CC by eluting with H_2O -MeOH (20:80) to afford 6-hydroxy-astropaquinone B (**1**, 11.0 mg), and astropaquinone D (**2**, 13.5 mg). Fr. 1-6 (1.2 g) was combined and chromatographed on a silica gel column chromatography using a stepwise gradient of CHCl_3 /EtOAc (100:0–0:100) to give 11 fractions (Frs. 1-6-1 to 1-6-11). Fr. 1-6-4 (62.0 mg) was subjected to ODS column chromatography by eluting stepwise with 100% H_2O to 100% MeOH (100:0–0:100), which yielded 11 fractions (Frs. 1-6-4-1 to 1-6-4-11). Fr. 1-6-4-5 (30.5 mg) was subjected to flash silica gel CC (CHCl_3 /MeOH, 50:1, *v/v*) to afford 3-*O*-methyl-9-*O*-methylfusarubin (**3**, 3.2 mg).

6-Hydroxy-astropaquinone B (**1**): an orange amorphous powder; $[\alpha]_{\text{D}}^{20} +55^\circ$ (*c* 0.3, MeOH); UV (MeOH); λ_{max} (log ϵ) 226 (4.3), 283 (3.7), 491 (3.2) nm; HRESITOFMS *m/z* 357.0931 $[\text{M} + \text{Na}]^+$, (calcd for $\text{C}_{17}\text{H}_{18}\text{O}_7\text{Na}$, 357.0950); IR (KBr): 3397, 2927, 2923, 1650, 1616 1457, 1376, 1256, 1051 cm^{-1} ; $^1\text{H-NMR}$ (600 MHz, CDCl_3) δ_{H} (ppm, *J* Hz): 1.38 (3 H, d, *J* = 6.7 Hz, H-11), 2.22 (1 H, dd, *J* = 18.0, 10.0, H-4), 2.66 (1 H, dd, *J* = 18.0, 3.5, H-4), 3.57 (3 H, s, 1-OMe), 4.00 (3 H, s, 7-OMe), 4.03 (3 H, s, 9-OMe), 4.21 (1 H, m, H-3), 5.52 (1 H, s, H-1), 6.74 (1 H, s, H-8); $^{13}\text{C-NMR}$ (150 MHz, CDCl_3) δ_{C} (ppm): 21.1 (C-11), 28.6 (C-4), 56.4 (1-OMe), 56.5 (7-OMe), 56.8 (9-OMe), 62.0 (C-3), 93.8 (C-1), 102.8 (C-8), 110.0 (C-9a), 114.5 (C-5a) 140.8 (C-4a), 142.7 (C-10a), 148.4 (C-6), 155.5 (C-9) 156.0 (C-7), 179.5 (C-10), 190.3 (C-5).

Astropaquinone D. (**2**): an orange amorphous powder; $[\alpha]_{\text{D}}^{20} +100^\circ$ (*c* 0.3, MeOH); UV (MeOH): λ_{max} (log ϵ): 227 (4.3), 284 (3.6), 490 (3.2); HRESITOFMS *m/z* 327.0823 $[\text{M} + \text{Na}]^+$, (calcd for $\text{C}_{16}\text{H}_{16}\text{O}_6\text{Na}$, 327.0844); IR (KBr): 3502, 2923, 1631, 1616, 1434, 1469, 1265, 1051 cm^{-1} ; $^1\text{H-NMR}$ (600 MHz, CDCl_3) δ_{H} (ppm, *J* Hz): 1.37 (3 H, d, *J* = 6.7, Me-11), 2.28 (1 H, m, H-4) 2.69 (1 H, dt, *J* = 18.3, 3.6 Hz, H-4), 3.65 (1 H, m, H-3), 3.99 (3 H, s, 7-OMe), 4.01 (3 H, s, 9-OMe) 4.50 (1 H, dt, *J* = 19.0, 2.0 Hz, H-1), 4.86 (1 H, dd, *J* = 19.0, 2.4, H-1), 6.71 (1 H, s, H-8); $^{13}\text{C-NMR}$ (150 MHz, CDCl_3) δ_{C} (ppm): 21.3 (C-11), 28.8 (C-4), 56.5 (7-OMe), 56.9 (9-OMe), 64.1 (C-1), 70.0 (C-3), 102.3 (C-8), 111.3 (C-9a), 114.3 (C-5a), 139.2 (C-4a), 146.4 (C-10a), 148.5 (C-6), 155.7 (C-9), 156.0 (C-7), 180.5 (C-10), 189.5 (C-5).

3.4. Bioassays

3.4.1. Antimicrobial activity

The minimal inhibition concentration (MIC) was determined by the agar dilution method, using nutrient agar for bacteria (1% beef extract, 1% peptone and 0.5% NaCl at pH 7.2). This test was performed in Petri dishes (4 cm. id.) in duplicate. Each test compound was dissolved at $1 \text{ mg} \cdot \text{mL}^{-1}$ in 10% aqueous DMSO. A suitably quantified volume of the test solution was mixed with the appropriate agar medium (2 mL) to prepare a plate with a given concentration (0 – $100 \mu\text{g} \cdot \text{mL}^{-1}$) of a test compound. Each plate was subsequently inoculated with a test microorganism ($100 \mu\text{L}$ of approximately $10^6 \text{ CFU} \cdot \text{mL}^{-1}$) and incubated at 30°C for 18–24 h for bacteria. MIC is defined as the lowest concentration resulting in no visible growth after incubation. Chloramphenicol and kanamycin were used for positive control against *S. aureus* and *P. aeruginosa* (each $1 \mu\text{g} \cdot \text{mL}^{-1}$), respectively.

3.4.2. Phytotoxic assay

Lettuce seeds (*Lactuca sativa* L.) were used for the bioassay. Fifteen seeds were sown on filter paper containing a defined concentration of the test compound in a Petri dish (4 cm id.). Distilled water (1 mL) was added to the Petri dish, and incubation was carried out at 25 °C under continuous light for 7 days. The elongation of the roots and shoots was measured to compare with those of the control. The negative control was using only distilled water and the positive control was using 2,4-dichlorophenoxyacetic acid.

4. Conclusion

As a result, the chemical investigation of an endophytic fungus *F. napiforme* IP-28 led to the discovery of three compounds **1–3**. Their structures were established on the basis of NMR spectroscopic data. Compounds **1–3** showed the moderate antibacterial activity and phytotoxicity on lettuce seeding at a concentration of 30 µg · mL⁻¹. The results in this article implied that endophytes living special environment such as mangrove area might be able to have survival strategy including production of the metabolites with unique skeletons displaying a wide range of biological effects.

Acknowledgements

The authors thank Dr. Takako Aboshi in Yamagata University for MS experimental supports.

Funding

This investigation was financially supported by RISTEKDIKTI, Indonesia (World Class Professor Grant, T/83/ D2.3/KK.04.05/2019 by U.S.).

References

- Bulbule VJ, Koranne PS, Deshpande VH, Borate HB. 2007. Simple and efficient synthesis of thysanone methyl ether. *Tetrahedron*. 63(1):166–170.
- Geng WL, Wang XY, Kurtán T, Mándi A, Tang H, Schulz B, Sun P, Zhang W. 2012. Herbarone, a rearranged heptaketide derivative from the sea hare associated fungus *Torula herbarum*. *J Nat Prod*. 75(10):1828–1832.
- Ismaiel AA, Papenbrock J. 2015. Mycotoxins: producing fungi and mechanisms of phytotoxicity. *Agriculture*. 5(3):492–537.
- Kornsakulkarn J, Dolsophon K, Boonyuen N, Boonruangprapa T, Rachtawee P, Prabpai S, Kongsaree P, Thongpanchang C. 2011. Dihydronaphthalenones from endophytic fungus *Fusarium* sp. BCC14842. *Tetrahedron*. 67(39):7540–7547.
- Li SJ, Zhang X, Wang XH, Zhao CQ. 2018. Novel natural compounds from endophytic fungi with anticancer activity. *Eur J Med Chem*. 56:316–343.
- Masi M, Nocera P, Zonno MC, Tuzi A, Pescitelli G, Cimmino A, Boari A, Infantino A, Vurro M, Evidente A. 2018. Lentiquinones A, B, and C, Phytotoxic Anthraquinone Derivatives Isolated from *Ascochyta lentis*, a Pathogen of Lentil. *J. Nat. Prod*. 81:2700–2709.
- Marasas WFO, Rabie CJ, Lübben A, Nelson PE, Toussoun TA, Wyk PSV. 1987. *Fusarium Napiforme*, A new species from millet and sorghum in Southern Africa. *Mycologia*. 79(6):910–914.

- Ravichandiran P, Sheet S, Premnath D, Kim AR, Yoo DJ. 2019. 1,4-Naphthoquinone analogues: potent antibacterial agents and mode of action evaluation. *Molecules*. 24(7):1437.
- Shiono Y, Miyazaki N, Murayama T, Koseki T, Harizon Katja DG, Supratman U, Nakata J, Kakihara Y, Saeki M, et al. 2016. GSK-3 β inhibitory activities of novel dichlororesorcinol derivatives from *Cosmospora vilior* isolated from a mangrove plant. *Phytochem Lett*. 18:122–127.
- Tatum JH, Baker RA, Berry RE. 1985. Naphthoquinones produced by *Fusarium solani* isolated from citrus. *Phytochemistry*. 24(3):457–459.
- Venugopalan A, Srivastava S. 2015. Endophytes as in vitro production platforms of high value plant secondary metabolites. *Biotechnol Adv*. 33(6 Pt 1):873–887.
- Wang L, Dong JY, Song HC, Shen KZ, Wang LM, Sun R, Wang CR, Gao YX, Li GH, Li L, Zhang KQ. 2009. Three new naphthoquinone pigments isolated from the freshwater fungus, *Astrosphaeriella papuana*. *Planta Med*. 75(12):1339–1343.
- Zakaria L. 2017. Mycotoxigenic *Fusarium* species from agricultural crops in Malaysia. *Proc Jpn Assoc Mycotoxicol*. 67(2):67–75.