

PAPER NAME

**k. Indonesia Journal of Geography\_Biom  
ass carbon stock assessment...2022.pdf**

AUTHOR

**Abdul Malik**

WORD COUNT

**5287 Words**

CHARACTER COUNT

**25010 Characters**

PAGE COUNT

**9 Pages**

FILE SIZE

**579.5KB**

SUBMISSION DATE

**Sep 18, 2022 1:29 AM GMT+8**

REPORT DATE

**Sep 18, 2022 1:29 AM GMT+8**

### ● 6% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.

- 5% Internet database
- 4% Publications database
- Crossref database
- Crossref Posted Content database
- 3% Submitted Works database

### ● Excluded from Similarity Report

- Bibliographic material
- Quoted material
- Cited material
- Small Matches (Less than 10 words)
- Manually excluded sources
- Manually excluded text blocks

# Biomass Carbon Stock Assessment of Mangrove Ecosystem in Pannikiang Island South Sulawesi Indonesia

Abdul Malik<sup>1\*</sup>, Uca Sideng<sup>1</sup>, and Jaelani<sup>2</sup>

<sup>1</sup>Department of Geography, Faculty of Mathematics and Natural Sciences, Universitas Negeri Makassar (UNM), Jl. Malengkeri Raya, Kampus UNM Parangtambung, Makassar, Indonesia 90224, <sup>2</sup>Study Program of Geography Education, Postgraduate Program of Universitas Negeri Makassar, Jalan Bontolangkasa, Makassar.

Received: 2019-07-28  
Accepted: 2022-03-08

**Keywords:**  
biomass carbon stock; climate change; South Sulawesi

**Correspondent email:**  
abdulmalik@unm.ac.id

**Abstract.** One of the essential services provided by mangroves is carbon sequestration, and therefore climate change mitigation. While previous assessments of mangrove carbon stocks and sequestrations have focused on the estuarine and deltaic mangrove setting, there are still limited studies carried out at small island mangroves. The study aims to assess mangrove biomass carbon stocks in Pannikiang, a small island in South Sulawesi, Indonesia, which occupies 91.64 ha of species-rich pristine mangrove forests. A field-based data collection survey was performed using a circular plot approach, while above-ground tree carbon (AGC) and below-ground root carbon (BGC) stocks were estimated using available species-specific allometric equations. The mean AGC and BGC were  $5.34 \pm 0.17$  and  $1.68 \pm 0.04$  Mg C ha<sup>-1</sup>, respectively. *Bruguiera gymnorrhiza* mangrove species stored the greatest of carbon stocks, followed by *Scyphiphora hydrophyllacea*. Carbon stocks obtained from small island mangroves in this study were lower than stocks assessed from other mangrove locations across Indonesia and Southeast Asia. However, historical rates of deforestation in Pannikiang Island may generate emissions to approximately 82.17 Mg CO<sub>2</sub>-eq. Findings from this study will be beneficial in providing baseline data for policy decision-making on climate change mitigation in the region, specifically for improved land use management via a low carbon development agenda

©2022 by the authors. Licensee Indonesian Journal of Geography, Indonesia.  
This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY-NC) license <https://creativecommons.org/licenses/by-nc/4.0/>.

## 1. Introduction

Climate change is amongst the most challenging in this century, and the increase of greenhouse gasses (GHGs), mainly from carbon dioxide (CO<sub>2</sub>) emissions to the atmosphere, are the primary cause of climate change (Bindu dkk., 2020). Climate change mitigation is not only an effort to reduce the level of CO<sub>2</sub> emissions but also balance with managing ecosystem services as carbon sinks (Wahyudi et al., 2017).

Mangroves are one of the most important coastal ecosystems in tropical and subtropical regions (Giri et al., 2011). They play an essential role in climate change mitigation by sequestering CO<sub>2</sub> from the atmosphere via photosynthesis and carbon (C) from the ocean (Howard et al., 2014). Mangroves are among the most massive forest carbon sinks per hectare than other tropical forests (Donato dkk., 2011; Alongi, 2014). However, they are particularly vulnerable to detach CO<sub>2</sub> and other GHGs to the atmospheres that could impact global climate change if disturbed, as a result of logging and conversion to other land use such as agricultural land and aquaculture ponds (Murdiyarso et al., 2015; Sukarna and Syahid, 2015). Both Friess (2016) and Murdiyarso et al (2015) noted the continuous global demands for mangrove commodities and land-use change activities resulted in the increasing degradation and deforestation in recent decades.

Since 1980, almost half of the global mangrove forests have disappeared (FAO, 2007), and Southeast Asia was the highest loser region (Richards and Friess, 2016). The loss of mangroves has resulted in carbon emissions into the atmosphere of about 0.08-0.48 Pg CO<sub>2</sub> yr<sup>-1</sup> or 10% of the total global CO<sub>2</sub> emissions of the world (Donato et al., 2011; Murdiyarso et al., 2015). In particular, Indonesia, which has the most significant percentage of mangroves in the world (22%), has lost about 40% of its mangroves (FAO, 2007) and resulting in a 20% carbon emissions increase due to land-use change (Murdiyarso et al., 2015).

The coastal area of South Sulawesi is an important area for mangrove blue carbon storage in Indonesia (Cameron dkk., 2019; Malik dkk., 2020). Mangroves are found in the coastal area of Makassar city and regencies of Maros, Pangkep, Barru, Pinrang, East Luwu, Luwu, Bone, Sinjai, Takalar, Jeneponto, Bantaeng, and Bulukumba (Saputro, 2009). In the 1950s, South Sulawesi mangrove-covered area of approximately 100,000 hectares Giesen et al (1991), which decreased around 89.6% to 10,412 hectares by 2017 (Susanto et al., 2018; Rahadian et al., 2019). The main driver of mangrove deforestation is conversion to aquaculture pond (Malik et al., 2017; 2020 ; Jalil et al., 2020), with annual deforestation rates between one and five percent during 1979 – 2012 (Malik et al., 2017). If current trends continue,

local mangroves could be lost in the next 20-150 years (Malik et al., 2017), and the carbon stocks stored in these areas will become a source of significant carbon emissions.

While most previous mangrove carbon studies focused on the estuarine and deltaic mangrove setting, there is still a lack of studies conducted at small island mangroves. Pannikiang Island is one of the small island mangrove hotspots in South Sulawesi. Mangroves occupied about 91.64 ha of the total area of the island (96.88 ha) in 2018. However, mangroves have disappeared by 3.19 ha since 1997 due to conversion into aquaculture ponds and settlements (Jaelani et al., 2021).

Although the historical rate of mangrove deforestation has occurred on a small scale during the period 1997-2018, it is critical to investigate the current carbon stock and carbon emissions from the mangrove deforested on this island. Therefore, this study's objectives are to assess spatial

variation by comparing carbon stocks between Pannikiang Island study sites and are to estimate carbon emissions generated by deforestation in this region between 1997 and 2018. Our findings contribute to providing baseline data for policy decision-making in climate change mitigation and land use management.

### 3. Methods

#### 4 Study Area

The research was carried out in the mangrove area of Pannikiang Island, Barru Regency. This island is located within the latitude of  $4^{\circ}20'00''$  -  $4^{\circ}22'00''$  and longitude of  $119^{\circ}35'28.38''$  -  $119^{\circ}36'30''$  (Figure 1), about 108 km from the capital of South Sulawesi, Makassar City, and 15 km from the center of Barru Regency.

Pannikiang island was selected as considered as one of the potential carbon sequestration projects based on the




Figure 1. Pannikiang Island, Barru Regency, South Sulawesi, and plot locations

extent of the mangrove cover area. The mangrove area is dominated by *Rhizophora stylosa* (Suwardi, 2014) and habitat for many faunae, mainly for thousands of bats (Dinas Pariwisata Kabupaten Barru, 2017). About 55 households occupy this island, and most of the head households work as fishermen.

**Data Collection**

Data on biomass carbon stocks were collected in April 2019. We implemented a circular plot approach by adopting a sampling protocol developed by Kauffman dan Donato (2012) to measure above-ground tree biomass (AGB) and below-ground root biomass (BGB) stocks at the eight plots selected (Figure 1). For each plot, we established five circular subplots with a radius of 7 m and 25 m distance to each subplot center (Figure 2) using a measuring tape and marked the position using the Global Positioning System/GPS. We also identified the species name and recorded the number of individual species of all mangrove trees using a tally counter. Finally, we measured stem diameter at breast height (DBH) 1.3 m above the ground or 30 cm above the highest prop root for *Rhizophora* sp. using measuring tape




Figure 2. Plot layout for biomass carbon stock assessment (adopted from Kauffman dan Donato (2012)).

Table 1. Wood density of mangroves

Species	Wood density (g cm <sup>-3</sup> )
<i>Bruguiera cylindrica</i>	0.72
<i>Bruguiera gymnorrhiza</i> (Bg)	0.81
<i>Ceriops decandra</i>	0.87
<i>Ceriops tagal</i>	0.85
<i>Hibiscuse tiliaceocus</i>	0.57
<i>Rhizophora apiculata</i>	0.87
<i>Rhizophora stylosa</i>	0.84
<i>Rhizophora mucronata</i>	0.83
<i>Sonneratia alba</i>	0.47
<i>Scyphiphora hydrophyllacea</i>	0.69
<i>Xylocarpus granatum</i>	0.61
<i>Xylocarpus mollucensis</i>	0.65

and tree heights using a clinometer (Malik et al., 2019; 2020).

**Data Analysis**

Species density, individual (ind.) tree basal area, and the total of tree basal area calculated using equations 1 - 3:

$$D = \frac{ni}{A} \quad (1); \quad bai = \left(\frac{1}{2}DBH\right)^2\pi \quad (2); \quad BA = \sum bai \quad (3)$$

where:

D: density of species i (ind. m<sup>-2</sup>). bai: ind. tree basal area (m<sup>2</sup> ha<sup>-1</sup>). BA: total tree basal area per hectare (m<sup>2</sup> ha<sup>-1</sup>). ni: number of standing species i. A: total area of the sample subplots per plot (769.3 m<sup>2</sup>). DBH: diameter at breast height (cm).

To calculate the AGB and BGB stocks of mangrove in this study, we used allometric equations 4 and 5 that were developed by Komiyama et al (2005) :

$$AGB = 0.251*r*DBH^{2.46} \quad (4)$$

$$BGB = 0.199*r*0.899*DBH^{2.22} \quad (5)$$

where, AGB and BGB: above- and below-ground biomass on mangrove tree and root (kg). r: species-specific wood density (see Table 1).

Furthermore, to estimate the Above-ground tree carbon (AGC) and Below-ground root carbon (BGC) stocks, we used equations 6 and 7 (Kauffman and Donato, 2012; Howard et al., 2014):

$$AGC = AGB \times 0.48 \quad (6)$$

$$BGC = BGB \times 0.39 \quad (7)$$

where, AGC tree and BGC root: above- and below-ground carbon on mangrove tree and root (kg C m<sup>2</sup>). 0.48 and 0.39: carbon conversion factor for AGB and BGB.

Moreover, to calculate the loss of AGC and BGC stocks, we multiplied the mean value of AGC and BGC stocks by the historical rate of mangrove deforestation during the period 1997-2018 (3.19 ha) as reported by Jaelani et al (2021), while to calculate carbon emissions, we multiplied the loss value of AGC and BGC stocks by the ratio of molecular weights between carbon dioxide (44) and carbon (12) (Kauffman and Donato, 2012).

**3. Results And Discussion**

**Species Composition of Mangrove**

A total of mangrove tree (392 trees) was identified in this study. Twelve mangrove species were recorded include *Bruguiera cylindrical* (Bc), *Bruguiera gymnorrhiza* (Bg), *Ceriops decandra* (Cd), *Ceriops tagal* (Ct), *Hibiscuse tiliaceocus* (Ht), *Rhizophora apiculata* (Ra), *Rhizophora*




Figure 3. AGC and BGC stocks for each plot include the standard error of the mean




Figure 4. AGC and BGC stocks of mangrove species include the standard error of the mean.

mucronata (Rm), *Rhizophora stylosa* (Rs), *Sonneratia alba* (Sa), *Scyphiphora hydrophyllacea* (Sh), *Xylocarpus granatum* (Xg), and *Xylocarpus mollucensis* (Xm).

The number of mangrove species in this study area was higher than in the similar areas of South Sulawesi, such as in Takalar Regency (10 species) (Malik et al., 2015) and Pangkep Regency (11 species) (Jalil et al., 2020), but lower than in Sinjai Regency (15 species) as reported by Suharti et al (2016).

Rs was the dominating species (162 trees) and found in all plots and followed by Bg (70 trees). Malik et al. (2015) reported that the dominance of *Rhizophora* sp. in South Sulawesi is similar to other areas in Indonesia and Southeast Asia, such as in Segara Anakan Lagoon in Central Java and Matang in Malaysia. Duke et al (1998) stated the dominance of *Rhizophora* sp. in Indonesia and other Southeast Asia countries is influenced by physical, environmental, and climate factors, such as soil characteristics, moderately high and good rainfall distribution, and the suitable temperature range for the species growths.

Plot 1 was registered as the largest number of the tree (97 trees). The mean value of the tree density ( $0.03 \pm 0.012$  ind. m<sup>-2</sup>) is higher compared to other plots. The mean value of tree DBH and height were  $11.05 \pm 1.54$  cm and  $7.23 \pm 0.74$  m, respectively, whereas the mean stand basal area was  $2.27 \pm 0.55$  m<sup>2</sup> ha<sup>-1</sup>. Although Rs was dominating in all plots, Bg and Sh have higher tree DBH than other species.

### Biomass Carbon Stocks of Mangrove

The mean concentration of AGC and BGC stocks at eight plots was  $5.34 \pm 0.17$  Mg C ha<sup>-1</sup> and  $1.68 \pm 0.04$  Mg C ha<sup>-1</sup>, respectively. The greatest AGC and BGC stocks were found at plot 8 ( $23.60 \pm 1.42$  Mg C ha<sup>-1</sup> and  $6.86 \pm 0.37$  Mg C ha<sup>-1</sup>), whereas the lowest figure was at plot 1 ( $0.21 \pm 0.02$  Mg C ha<sup>-1</sup> and  $0.09 \pm 0.01$  Mg C ha<sup>-1</sup>) (Figure 3).

Considering the total area of mangroves (91.64 ha) in 2018 (Jaelani et al., 2021), the total of AGC and BGC stocks were 489.35 Mg C and 153.95 Mg C respectively. However, the mean value of AGC and BGC stocks were much lower compared to other places in Indonesia, such as in Bunaken, North Sulawesi ( $69.2$  Mg C ha<sup>-1</sup> and  $14.9$  Mg C ha<sup>-1</sup>), Kubu Raya, West Kalimantan ( $134.8$  Mg C ha<sup>-1</sup> and  $14.3$  Mg C ha<sup>-1</sup>), Sembilang, South Sumatra ( $300.5$  Mg C ha<sup>-1</sup> and  $27.2$  Mg C ha<sup>-1</sup>), and Timika, Papua ( $323.6$  Mg C ha<sup>-1</sup> and  $43.6$  Mg C ha<sup>-1</sup>) as reported by Murdiyarso et al (2015), and also to other places in Southeast Asia, such as in Palawan, Philippine  $263.8$  Mg C ha<sup>-1</sup> and  $92.3$  Mg C ha<sup>-1</sup> as reported




Figure 5a. Relationship between DBH and AGC-BGC stocks




Figure 5b. Relationship between DBH and AGC-BGC stocks

by (Abino et al., 2014), and in Mekong Delta, Vietnam  $61.4$  Mg C ha<sup>-1</sup> and  $8.7$  Mg C ha<sup>-1</sup> as stated by Nam et al (2016). Low AGC and BGC stocks are likely due to the smaller size tree diameter (most of the tree diameter below 15 cm) compared to those places. Besides, this may be influenced by the high salinity that characteristic of the waters of small islands compared to the mainland estuary system. Therefore, it can affect mangrove growth.

Although Rs is the dominant species in this area, Bg represented the greatest AGC and BGC stocks ( $16.07 \pm 1.64$  Mg C ha<sup>-1</sup> and  $4.54 \pm 0.43$  Mg C ha<sup>-1</sup>), followed by Sh ( $6.18 \pm 0.71$  Mg C ha<sup>-1</sup> and  $1.90 \pm 0.20$  Mg C ha<sup>-1</sup>) (Figure 4). It is influenced by the larger tree diameter of this species compared to the other species.

The relationship between DBH and above-below ground carbon stocks is presented in Figure 5. The increased value of tree DBH influenced the trend concentration of AGC and BGC stocks. Plotted in linear regression showed that the correlation coefficient ( $r$ ) between DBH and AGC and BGC stocks were 0.86 and 0.89, respectively. This finding confirms a study from Alavaisha dan Mangora (2016) that noted DBH mangrove trees' significant effect on the AGC stock concentration. The value of BGC stock is positively correlated with the tree diameter. Hence, if the tree diameter is large, the BGC stock is large as well (Perera and Amarasinghe, 2013).

Furthermore, considering the conversion of mangrove areas to aquaculture ponds and settlements that reach 3.19 ha during the period 1997 – 2018 (Jaelani et al., 2021), the loss of AGC and BGC stocks was  $17.03$  Mg C and  $5.36$  Mg C, respectively, or it causes the carbon emissions of  $82.17$  Mg CO<sub>2</sub>-eq or  $3.91$  Mg CO<sub>2</sub>-eq per year into the atmosphere. It suggests that if deforestation is halted and mangrove is restored, historical emissions from the past 21 years can be effectively offset over the same period (4.76% each year over 21 years).

## Conclusion

This study has demonstrated the biomass carbon stocks in Pannikiang Island, Barru Regency, South Sulawesi, Indonesia. The mean value of AGC and BGC stocks is  $5.34 \pm 0.17 \text{ Mg C ha}^{-1}$  and  $1.68 \pm 0.04 \text{ Mg C ha}^{-1}$ , respectively. Although *Rs* is the dominant species, *Bruguiera gymnorrhiza* was the highest AGC and BGC stocks ( $16.07 \pm 1.64 \text{ Mg C ha}^{-1}$  and  $4.54 \pm 0.43 \text{ Mg C ha}^{-1}$ ), followed by *Scyphiphora hydrophyllacea* ( $6.18 \pm 0.71 \text{ Mg C ha}^{-1}$  and  $1.90 \pm 0.20 \text{ Mg C ha}^{-1}$ ). Tree DBH has a significant effect on the value of the AGC and BGC stocks. The mean value of AGC and BGC stocks is much lower than other places in Indonesia and Southeast Asia due to the lower tree diameter. However, mangrove clearing for different land uses has resulted in carbon emissions of  $82.17 \text{ Mg CO}_2\text{-eq}$  in the last two decades. Therefore, preventing mangroves from conversion to other land uses and conserving intact mangroves are essential actions to reduce  $\text{CO}_2$  emissions and help mitigate climate change.

## Acknowledgment

Our gratitude goes to the Ministry of Research, Technology, and Higher Education of the Republic of Indonesia for funding this research through the scheme of Penelitian Pascasarjana-Tesis Magister (contract number: 123/UN36.9/PL/2019). We thank the Institute of Research and Community Service Universitas Negeri Makassar, Department of Geography Universitas Negeri Makassar, and the Governments of South Sulawesi and Barru Regency for facilitating this research. We also thank the anonymous reviewer for the excellent comments and suggestions for this paper.

## References

- Abino, A. C., Castillo, J. A. A., & Lee, Y. J. (2014). Species Diversity, Biomass, and Carbon Stock Assessments of A Natural Mangrove Forest in Palawan, Philippines. *Pak. J. Bot.*, *46*(6), 1955–1962.
- Alavaisha, E., & Mangora, M. M. (2016). Carbon Stocks in the Small Estuarine Mangroves of Geza and Mtimbwani, Tanga, Tanzania. *International Journal of Forestry Research*, *2016*, 1–11. <https://doi.org/10.1155/2016/2068283>
- Alongi, D. M. (2014). Carbon Cycling and Storage in Mangrove Forests. *Annual Review of Marine Science*, *6*(1), 195–219. <https://doi.org/10.1146/annurev-marine-010213-135020>
- Bindu, G., Rajan, P., Jishnu, E. S., & Ajith Joseph, K. (2020). Carbon stock assessment of mangroves using remote sensing and geographic information system. *The Egyptian Journal of Remote Sensing and Space Science*, *23*(1), 1–9. <https://doi.org/10.1016/J.EJRS.2018.04.006>
- Dinas Pariwisata Kabupaten Barru (2017). Pulau Pannikiang. <http://dispar.barrukab.go.id/package/pulau-pannikiang/>. Accessed on 21 June 2019.
- Cameron, C., Hutley, L. B., Friess, D. A., & Brown, B. (2019). Community structure dynamics and carbon stock change of rehabilitated mangrove forests in Sulawesi, Indonesia. *Ecological Applications*, *29*(1). <https://doi.org/10.1002/eap.1810>
- Donato, D. C., Kauffman, J. B., Murdiyarso, D., Kurnianto, S., Stidham, M., & Kanninen, M. (2011). Mangroves among the most carbon-rich forests in the tropics. *Nature Geoscience*, *4* (5), 293–297. <https://doi.org/10.1038/ngeo1123>
- Duke, N. C., Ball, M., & Ellison, J. C. (1998). Factors Influencing Biodiversity and Distributional Gradients in Mangroves. *Global*

- Ecology and Biogeography Letter*, *7*(1), 27–47.
- FAO. (2007). *The World's Mangrove 1980-2005* (No. 153; FAO Forestry Paper).
- Friess, D. A. (2016). Ecosystem Services and Disservices of Mangrove Forests: Insights from Historical Colonial Observations. *Forests* *2016*, Vol. 7, Page 183, *7*(9), 183. <https://doi.org/10.3390/F7090183>
- Giesen, W., Baltzer, M., & Baruadi, R. (1991). *Integrating conservation with land-use development in wetlands of South Sulawesi*. Asian Wetland Bureau (AWB)-Indonesia.
- Giri, C., Ochieng, E., Tieszen, L. L., Zhu, Z., Singh, A., Loveland, T., Masek, J., & Duke, N. (2011). Status and distribution of mangrove forests of the world using earth observation satellite data. *Global Ecology and Biogeography*, *20*(1), 154–159. <https://doi.org/10.1111/j.1466-8238.2010.00584.x>
- Howard, J., Hoyt, S., Isensee, K., Telszewski, M., & Pidgeon, E. (2014). *Coastal blue carbon: methods for assessing carbon stocks and emissions factors in mangroves, tidal salt marshes, and seagrasses*. Conservation International, Intergovernmental Oceanographic Commission of UNESCO, International Union for Conservation of Nature. <https://cgspace.cgiar.org/handle/10568/95127>
- Jaelani, Malik, A., & Djalil, A. R. (2021). Mangrove Changes in Pannikiang Island Barru Regency South Sulawesi. *Journal of Physics: Conference Series*, *1752*(1), 012088. <https://doi.org/10.1088/1742-6596/1752/1/012088>
- Jalil, A., Malik, A., Nurdin, N., Saru, A., & Yunus, I. (2020). Assessment of seawater level, inundation duration and substrate elevation for mangrove rehabilitation program in the Spermonde Archipelago South Sulawesi Indonesia. *International Journal of Conservation Science. International Journal of Conservation Science*, *11*(4), 1115–1126.
- Kauffman, J. ., & Donato, D. . (2012). *Protocols for the measurement, monitoring and reporting of structure, biomass and carbon stocks in mangrove forests* (No. 86).
- Komiyama, A., Pongparn, S., & Kato, S. (2005). Common allometric equations for estimating the tree weight of mangroves. *Journal of Tropical Ecology*, *21*(4), 471–477. <https://doi.org/10.1017/S0266467405002476>
- Malik, A., Fensholt, R., & Mertz, O. (2015). Mangrove exploitation effects on biodiversity and ecosystem services. *Biodiversity and Conservation*, *24*(14), 3543–3557. <https://doi.org/10.1007/s10531-015-1015-4>
- Malik, A., Jalil, A. R., Arifuddin, A., & Syahmuddin, A. (2020). Biomass Carbon Stocks In The Mangrove Rehabilitated Area Of Sinjai District, South Sulawesi, Indonesia. *GEOGRAPHY, ENVIRONMENT, SUSTAINABILITY*, *13*(3), 32–38. <https://doi.org/10.24057/2071-9388-2019-131>
- Malik, A., Mertz, O., & Fensholt, R. (2017). Mangrove forest decline: consequences for livelihoods and environment in South Sulawesi. *Regional Environmental Change*, *17*(1), 157–169. <https://doi.org/10.1007/s10113-016-0989-0>
- Malik, A., Rahim, A., & Sideng, U. (2019). Biodiversity Assessment of Mangrove Vegetation for the Sustainability of Ecotourism in West Sulawesi, Indonesia. *Aquaculture, Aquarium, Conservation and Legislation*, *12*(4), 1458–1466.
- Murdiyarso, D., Purbopuspito, J., Kauffman, J. B., Warren, M. W., Sasmito, S. D., Donato, D. C., Manuri, S., Krisnawati, H., Taberima, S., & Kurnianto, S. (2015). The potential of Indonesian mangrove forests for global climate change

- mitigation. *Nature Climate Change*, 5(12), 1089–1092. <https://doi.org/10.1038/nclimate2734>
- Nam, V. N., Sasmito, S. D., Murdiyarso, D., Purbopuspito, J., & MacKenzie, R. A. (2016). Carbon stocks in artificially and naturally regenerated mangrove ecosystems in the Mekong Delta. *Wetlands Ecology and Management*, 24(2), 231–244. <https://doi.org/10.1007/s11273-015-9479-2>
- Perera, K. A. R. ., & Amarasinghe, M. . (2013). Carbon Partitioning and Allometric Relationships between Stem Diameter and Total Organic Carbon (TOC) in Plant Components of *Bruguiera gymnorrhiza* (L.) Lamk. and *Lumnitzera racemosa* Willd. in a Microtidal Basin Estuary in Sri Lanka. *International Journal of Marine Science*, 3(9), 72–78. <https://doi.org/10.5376/ijms.2013.03.0009>
- Rahadian, A., Prasetyo, L. B., Setiawan, Y., & Wikantika, K. (2019). Tinjauan Historis Data dan Informasi Luas Mangrove Indonesia. *Media Konservasi*, 24(2), 163–178.
- Richards, D. R., & Friess, D. A. (2016). Rates and drivers of mangrove deforestation in Southeast Asia, 2000–2012. *Proceedings of the National Academy of Sciences*, 113(2), 344–349. <https://doi.org/10.1073/pnas.1510272113>
- Saputro, G. B. (2009). *Peta Mangrove Indonesia*. Pusat Survey Sumberdaya Alam Laut, Badan Koordinasi Survei dan Pemetaan Nasional (Bakosurtanal).
- Suharti, S., Darusman, D., Nugroho, B., & Sundawati, L. (2016). Economic Valuation As a Basis for Sustainable Mangrove Resource Management: A Case in East Sinjai, South Sulawesi. *Jurnal Manajemen Hutan Tropika (Journal of Tropical Forest Management)*, 22(1), 13–23. <https://doi.org/10.7226/jtfm.22.1.13>
- Sukarna, R. M., and Syahid, Y. (2015). FCD Application of Landsat for Monitoring Mangrove in Central Kalimantan. *Indonesian Journal of Geography*, 47(2), 160–170.
- Susanto, A., Surbaya, C., Poniman, A. (2018). Kebijakan satu peta; Momentum reformasi penyelenggaraan informasi geospasial nasional. Seminar Nasional Peran Geospasial dalam Membingkai NKRI, 2016, Bogor, Indonesia Bogor (ID): BIG, 23-24.
- Suwardi, Tambaru, E., Ambeng, Priosambodo, D. (2014). Keanekaragaman jenis mangrove di Pulau Panikiang Kabupaten Barru Sulawesi Selatan. Jurusan Biologi Fakultas Matematika dan Ilmu Pengetahuan Alam, Universitas Hasanuddin, Makassar. <http://repository.unhas.ac.id/handle/123456789/10148>. Accessed on 21 June 2019.
- Wahyudi, J., Afdali, Adi, N.S., Rustam, A., Hadiyanto, Rahmawati, S., Irawan. A., Dharmawan, I.W.E., Prayudha, B., Hafizt, M., Prayitno, H.B., Rahayu, Y.P., et al. (2017). Potensi cadangan dan serapan karbon ekosistem mangrove dan padang lamun Indonesia (Intisari bagi pengambil kebijakan). Pusat Penelitian Oseanografi, Lembaga Ilmu Pengetahuan Indonesia (P2O-LIPI). <https://www.researchgate.net/publication/326649366>. Accessed on 21 June 2019.

## Supplementary Materials

Table S1. Species Composition of Mangrove in Pannikiang Island South Sulawesi

Plot	Species	n	D (ind. m <sup>-2</sup> )	DBH (cm)	H (m)	BA (m <sup>2</sup> ha <sup>-1</sup> )
1 (4° 20'23,73"S - 119°	Bc	4	0.01	2.87	2.75	0.09
	Bg	8	0.01	4.02	2.75	0.18
	Rm	8	0.01	2.11	1.88	0.05
	Rs	53	0.07	2.28	3.41	0.08
	Sa	24	0.03	8.44	5.54	2.01
Subtotal		97	0.13	-	-	-
Mean ± SE			0.03 ± 0.012	3.94 ± 1.17	3.27 ± 0.62	0.48 ± 0.38
2 (4°20'23,54"S - 119° 36'9,06"E)	Bc	4	0.01	4.70	8.00	0.34
	Bg	10	0.01	6.05	4.50	0.58
	Rm	4	0.01	5.57	9.75	0.34
	Rs	12	0.02	7.43	10.00	0.74
	Sa	21	0.03	6.85	6.24	0.52
Subtotal		51	0.07	-	-	-
Mean ± SE			0.01 ± 0.004	6.12 ± 0.48	7.70 ± 1.05	0.50 ± 0.08
3 (4°20'32.58"S - 119° 35'59.29"E)	Bg	13	0.02	14.80	7.23	2.24
	Ra	10	0.01	11.08	7.10	12.60
	Rm	4	0.01	11.07	7.75	1.26
	Rs	32	0.04	11.16	7.31	1.28
	Sa	4	0.01	19.35	8.00	4.28
Subtotal		63	0.08	-	-	-
Mean ± SE			0.02 ± 0.007	13.49 ± 1.63	7.48 ± 0.17	4.33 ± 2.14
4 (4°20'36.53"S - 119° 36'0.41"E)	Bg	22	0.03	6.46	4.32	0.45
	Ra	7	0.01	5.82	4.00	0.37
	Rs	16	0.02	6.27	4.31	0.42
	Sa	9	0.01	8.56	4.67	1.27
	Subtotal		54	0.07	-	-
Mean ± SE			0.02 ± 0.004	6.78 ± 0.61	4.32 ± 0.14	0.63 ± 0.21
5 (4°21'17.2"S - 119° 35'56.5"E)	Bg	6	0.01	10.46	8.17	1.25
	Ct	4	0.01	15.37	9.25	2.58
	Ra	9	0.01	12.28	8.44	1.63
	Rs	16	0.02	15.72	9.38	2.93
	Subtotal		35	0.05	-	-
Mean ± SE			0.01 ± 0.003	13.46 ± 1.26	8.81 ± 0.30	2.10 ± 0.39
6 (4°21'15.6"S - 119° 35'55.4"E)	Bg	6	0.01	11.57	9.00	1.51
	Cd	1	0.001	20.70	11.00	4.37
	Ct	2	0.00	9.39	8.00	0.91
	Ra	6	0.01	11.46	10.17	2.03
	Rs	17	0.02	10.55	9.24	1.22
Subtotal		32	0.04	-	-	-
Mean ± SE			0.01 ± 0.004	12.74 ± 2.03	9.48 ± 0.51	2.01 ± 0.62



Table S1. Species Composition of Mangrove in Pannikiang Island South Sulawesi

7 (4°21'38.25"S - 119°35'37.39"E)	Bg	4	0.01	10.75	4.38	1.31
	Ht	5	0.01	28.41	7.62	10.53
	Ra	2	0.003	9.08	9.35	0.93
	Rs	3	0.004	8.70	7.97	0.85
	Sh	5	0.01	22.87	9.88	6.65
	Xg	1	0.001	6.05	4.30	0.37
	Xm	1	0.001	5.41	4.40	0.30
Subtotal		21	0.03	-	-	-
Mean ± SE			0.004 ± 0.001	13.04 ± 3.38	6.84 ± 0.92	2.99 ± 1.51
8 (4°21'43.60"S - 119°35'37.24"E)	Bg	1	0.001	43.31	12.50	19.14
	Cd	1	0.001	14.33	9.00	2.10
	Ct	1	0.001	12.10	8.00	1.49
	Ht	2	0.003	14.17	8.90	2.06
	Ra	10	0.01	10.13	7.08	1.11
	Rs	13	0.02	8.50	7.23	0.75
	Sh	8	0.01	27.64	11.81	9.07
	Xg	2	0.003	26.27	13.40	8.66
Xm	1	0.001	12.74	11.80	1.66	
Subtotal		39	0.05	-	-	-
Mean ± SE			0.01 ± 0.002	18.80 ± 3.80	9.97 ± 0.81	5.12 ± 2.05
Total		392	-	-	-	-
Grand Mean ± SE			0.01 ± 0.002	11.05 ± 1.54	7.23 ± 0.74	2.27 ± 0.55

Bc: *Bruguiera cylindrica*. Bg: *Bruguiera gymnorhiza*. Cd: *Ceriops decandra*. Ct: *Ceriops tagal*. Ht: *Hibiscuse tiliaceocus*. Ra: *Rhizophora apiculata*. Rm: *Rhizophora mucronata*. Rs: *Rhizophora stylosa*. Sa: *Sonneratia alba*. Sh: *Scyphiphora hydrophyllacea*. Xg: *Xylocarpus granatum*. Xm: *Xylocarpus mollucensis*. n: individual number of species *i*. D: density of species *i*. DBH: diameter of breast height. H: height. BA: basal area. SE: the standard error of the mean.

Table S2. Biomass carbon stocks of mangroves for each plot in Pannikiang Island South

Plot	AGB	BGB	AGC	BGC
	(Mg ha <sup>-1</sup> )	(Mg ha <sup>-1</sup> )	(Mg C ha <sup>-1</sup> )	(Mg C ha <sup>-1</sup> )
1	0.44 ± 0.05	0.24 ± 0.03	0.21 ± 0.02	0.09 ± 0.01
2	1.07 ± 0.05	0.56 ± 0.02	0.51 ± 0.02	0.22 ± 0.01
3	7.33 ± 0.27	3.19 ± 0.11	3.52 ± 0.13	1.24 ± 0.04
4	1.03 ± 0.02	0.53 ± 0.01	0.49 ± 0.01	0.21 ± 0.003
5	6.92 ± 0.38	2.95 ± 0.15	3.32 ± 0.18	1.15 ± 0.06
6	8.73 ± 0.79	3.65 ± 0.30	4.19 ± 0.38	1.42 ± 0.12
7	14.38 ± 1.04	5.72 ± 0.39	6.90 ± 0.50	2.23 ± 0.15
8	49.17 ± 2.96	17.60 ± 0.96	23.60 ± 1.42	6.86 ± 0.37
Grand Mean ± SE	11.13 ± 0.35	4.31 ± 0.11	5.34 ± 0.17	1.68 ± 0.04
Max	49.17 ± 2.96	17.60 ± 0.96	23.60 ± 1.42	6.86 ± 0.37
Min	0.44 ± 0.05	0.24 ± 0.03	0.21 ± 0.02	0.09 ± 0.01

AGB tree: above-ground tree biomass. BGB: below-ground root biomass. AGC tree: above-ground tree carbon. BGC tree: below-ground tree carbon. SE: the standard error of the mean.

Table S3. Biomass carbon stocks of mangrove species in Pannikiang Island South Sulawesi

Species	AGC (Mg C ha <sup>-1</sup> )	BGC (Mg C ha <sup>-1</sup> )
Bc	0.07 ± 0.02	0.03 ± 0.01
Bg	16.07 ± 1.64	4.54 ± 0.43
Cd	3.31 ± 0.70	1.07 ± 0.21
Ct	2.05 ± 0.23	0.72 ± 0.07
Ht	3.97 ± 1.38	1.24 ± 0.40
Ra	2.53 ± 0.08	0.93 ± 0.03
Rm	0.58 ± 0.15	0.22 ± 0.05
Rs	2.92 ± 0.13	1.07 ± 0.04
Sa	1.45 ± 0.24	0.52 ± 0.08
Sh	6.18 ± 0.71	1.90 ± 0.20
Xg	3.05 ± 1.44	0.95 ± 0.44
Xm	0.60 ± 0.23	0.22 ± 0.08
Grand Mean ± SE	3.56 ± 0.17	1.12 ± 0.05
Max	16.07 ± 1.64	4.54 ± 0.43
Min	0.07 ± 0.02	0.03 ± 0.01

Bc: *Bruguiera cylindrica*. Bg: *Bruguiera gymnorrhiza*. Cd: *Ceriops decandra*. Ct: *Ceriops tagal*. Ht: *Hibiscuse tiliaceocus*. Ra: *Rhizophora apiculate*. Rm: *Rhizophora mucronata*. Rs: *Rhizophora stylosa*. Sa: *Sonneratia alba*. Sh: *Scyphiphora hydrophyllacea*. Xg: *Xylocarpus granatum*. Xm: *Xylocarpus mollucensis*. AGC: above-ground tree carbon. BGC: below-ground root carbon. SE:

● **6% Overall Similarity**

Top sources found in the following databases:

- 5% Internet database
- Crossref database
- 3% Submitted Works database
- 4% Publications database
- Crossref Posted Content database

TOP SOURCES

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

1	<b>ges.rgo.ru</b> Internet	2%
2	<b>core.ac.uk</b> Internet	<1%
3	<b>journal.ugm.ac.id</b> Internet	<1%
4	<b>Jaelani, A Malik, A R Djali. "Mangrove Changes in Pannikiang Island B...</b> Crossref	<1%
5	<b>eprints.unm.ac.id</b> Internet	<1%
6	<b>ijcs.ro</b> Internet	<1%
7	<b>Abdul Malik, Rasmus Fensholt, Ole Mertz. "Mangrove exploitation effe...</b> Crossref	<1%
8	<b>Phang, Valerie X. H., L. M. Chou, and Daniel A. Friess. "Ecosystem carb...</b> Crossref	<1%

**University of Melbourne on 2016-10-31**

Submitted works

&lt;1%

## ● Excluded from Similarity Report

- Bibliographic material
- Cited material
- Manually excluded sources
- Quoted material
- Small Matches (Less than 10 words)
- Manually excluded text blocks

---

### EXCLUDED SOURCES

**humaniora.journal.ugm.ac.id**

**35%**

Internet

### EXCLUDED TEXT BLOCKS

**ISSN 2354-9114 (online), ISSN 0024-9521 (print)Indonesian Journal of Geography ...**

humaniora.journal.ugm.ac.id

---

**of Geography, Faculty of Mathematics and Natural Sciences, Universitas Negeri M...**

humaniora.journal.ugm.ac.id

---

**focusedon the estuarine and deltaic mangrove setting, there**

journal.ugm.ac.id

---

**Indonesian Journal of Geography, Vol. 54, No.1, 2022**

Garuda.Kemdikbud.Go.Id

---

**providing baseline data forpolicy decision-making**

www.journal.ugm.ac.id