

PAPER NAME

**17638-Article Text-69136-3-10-2022110
9.pdf**

AUTHOR

Abdul Malik

WORD COUNT

6600 Words

CHARACTER COUNT

32483 Characters

PAGE COUNT

9 Pages

FILE SIZE

827.9KB

SUBMISSION DATE

Nov 11, 2022 4:30 AM GMT+8

REPORT DATE

Nov 11, 2022 4:31 AM GMT+8

● 7% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.

- 5% Internet database
- 3% Publications database
- Crossref database
- Crossref Posted Content database
- 3% Submitted Works database

● Excluded from Similarity Report

- Bibliographic material
- Quoted material
- Cited material
- Small Matches (Less than 10 words)
- Manually excluded sources
- Manually excluded text blocks

THE POTENTIAL SOIL ORGANIC CARBON STOCKS IN MANGROVE AREAS OF SINJAI DISTRICT, SOUTH SULAWESI, INDONESIA

Abdul MALIK^{1*}, Muhammad ICHSAN ALI², Suwardi ANNAS³, Abdul Rasyid JALIL⁴, Restu Utami MULYA¹, Konstantina GRAVANI⁵

¹Department of Geography, Faculty of Mathematics and Natural Sciences, Universitas Negeri Makassar, Makassar, South Sulawesi, Indonesia

²Department of Civil and Planning, Faculty of Engineering, Universitas Negeri Makassar, Makassar, South Sulawesi, Indonesia

³Department of Statistics, Faculty of Mathematics and Natural Sciences, Universitas Negeri Makassar, Makassar, South Sulawesi, Indonesia

⁴Department of Marine Science, Faculty of Marine Science and Fisheries, Universitas Hasanuddin, Makassar, South Sulawesi, Indonesia

⁵Department of Geosciences and Natural Resource Management, Faculty of Science, University of Copenhagen, Copenhagen K, Denmark

Received 03 October 2020; accepted 30 June 2022

Highlights

- ▶ Soil organic carbon (SOC) stocks were estimated in disturbed mangroves of South Sulawesi.
- ▶ The mean values of the SOC stock are $413.10 \pm 12.37 \text{ Mg C ha}^{-1}$.
- ▶ Promoting restoration may preserve remaining mangroves and increase SOC stocks.

Abstract. The soil pool is the primary sink for carbon in mangrove wetlands and plays a major role in mitigating climate change. However, aquaculture pond expansions go further to disrupt carbon storage in mangroves. The aim of this study is to estimate the stock of soil organic carbon (SOC) in the mangrove area of South Sulawesi, Indonesia. The mangroves of Sinjai District in South Sulawesi are a disturbed region with no previous study on SOC stock. We implemented a line transect method at five study sites, collected 15 soil cores at a depth of 0–15 cm, 15–30 cm, and 30–50 cm, and performed soil analysis using the Loss on Ignition method. We find that the mean value of SOC stock is $413.10 \pm 12.37 \text{ Mg C ha}^{-1}$. More attention to the conservation and restoration of lost mangrove areas is a high priority. It may also increase the stock of SOC to mitigate climate change. This study will help to preserve the remaining mangroves.

Keywords: coastal blue carbon, climate change mitigation, disturbed mangroves, mangrove soil properties.

Introduction

Climate change mitigation is one of the essential ecosystem services provided by mangroves (Duncan et al., 2016; Soper et al., 2019; Malik et al., 2020). Although mangroves cover only a small portion of the planet, they store considerable organic carbon (Hopkinson et al., 2012; Nóbrega et al., 2015). Mangroves are one of the most carbon-rich forests (Donato et al., 2011), with a global average total forest stock of $738.9 \text{ Mg C ha}^{-1}$ (Alongi, 2020). Therefore, they represent a critical component in carbon sequestration for climate change

mitigation (Intergovernmental Panel on Climate Change, 2014; Murdiyarso et al., 2015).

Mangroves can store carbon in plant materials and soil pools (Howard et al., 2014). However, most of the carbon is in soil pools, which account for up to 50–90% of the total carbon stock of mangroves (Donato et al., 2011; Kauffman et al., 2011; Sharma et al., 2020).

Hamilton and Friess (2018) demonstrated that 70.65% of global mangrove carbon is deposited in mangrove soils. Alongi et al. (2015) and Murdiyarso et al. (2015) both reported that the mean proportion of organic soil carbon

*Corresponding author. E-mail: abdulmalik@unm.ac.id

storage in several mangrove areas of Indonesia was about 78%. Abino et al. (2014) revealed that the sediment carbon stock in the natural mangrove forest in Palawan, Philippines was 50%. Besides, Nam et al. (2016) demonstrated that the percentage of soil organic carbon stores from two different mangrove areas (natural and restoration area) in Mekong Delta, Vietnam was similar and reached 90%, respectively. It is also confirmed by Hong Tinh et al. (2020) results in northern Vietnam, who found that the SOC stock of 20–25 years of restored and intact mangroves was not significantly different.

The significant carbon content in soil is due to rapid rates of net primary production from autochthonous (in the form of litterfall and belowground fine root growth) and sedimentation from allochthonous (supply from upstream rivers and sea) (Donato et al., 2011; Alongi, 2012). However, the autochthonous have a higher contribution than allochthonous in soil carbon production (Chen et al., 2017; Kusumaningtyas et al., 2019; Sasmito et al., 2020b; Jennerjahn, 2020).

Unfortunately, mangrove soils might become a significant source of CO₂ emissions if disturbed by land-use change activities, such as conversion into settlements, agricultural lands, and aquaculture ponds (Donato et al., 2011; Murdiyarso et al., 2015; Sharma et al., 2020). In Indonesia and other Southeast Asian countries, the conversion of mangrove forests into aquaculture ponds is the primary land-use activity within the mangrove area (Richards & Friess, 2016). It has occurred rapidly in recent decades (Pendleton et al., 2012) and contributed to about 0.08–0.48 Pg CO₂, translating to 10% of the total global emissions (Donato et al., 2011). Moreover, Hamilton and Friess (2018) have shown that the annual rate of mangrove deforestation (0.26%) since 2000 has emitted between 5.76 Tg CO₂e and 13.95 Tg CO₂e due to conversion to aquaculture ponds in Indonesia (Sidik & Lovelock, 2013).

When mangrove forests are being cleared, and the soil is being excavated, the soil carbon is exposed to air, and subsequently, the accelerated microbial activity releases large amounts of CO₂ and other GHGs into the atmosphere (Howard et al., 2014). Consequently, conversion to aquaculture has become one of the primary sources of CO₂ emissions (Sidik & Lovelock, 2013).

The mangrove area in South Sulawesi province is one of Indonesia's most important areas for the blue carbon program (Malik et al., 2020). Mangroves are distributed in the districts of East Luwu, Luwu, Bone, Sinjai, Takalar, Barru, Pangkep and Pinrang (Rahardian et al., 2019). However, aquaculture development has been the primary driver of mangrove deforestation in the past decades (Malik et al., 2017, 2020). It can be declined the potential carbon stock of the mangrove area and increase a significant CO₂ emission to the atmosphere. Giesen et al. (1991) estimated that the mangrove forests of South Sulawesi accounted for a total area of 100 thousand hectares during the 1950s. However, approximately 10 thousand mangroves were deforested in 2017 due to woodcutting, primarily for aquaculture pond expansion (Rahadian et al.,

2019), with annual deforestation rates between 1% and 5% (Malik et al., 2017). Furthermore, the policymakers related to climate change mitigation policies often overlook the significant amount of carbon stocks in this region's case of mangrove deforestation (Malik et al., 2020).

Global climate negotiations have promoted mangroves' potential contribution and conservation in mitigating GHG emissions (Sasmito et al., 2020a). Many countries that own mangroves are interesting in blue carbon programs or fund mangrove protection and restoration through forest carbon programs (Bukoski et al., 2020). One critical step to developing a blue carbon program is data availability and accurate estimates of baseline carbon stocks (Bukoski et al., 2020). This area's soil carbon stock will be lower than other studies done in the region or Indonesia because of the disturbance. Therefore, this research aims to estimate soil organic carbon stocks in the disturbance mangrove area of South Sulawesi, Indonesia, for helping mitigate climate change.

1. Materials and methods

1.1. Study area

The study area is situated in the mangrove area of Sinjai District, South Sulawesi, focusing in East Sinjai sub-District (latitudes 5°7'00"–5°14'00" and longitudes 120°15'00"–120°19'00", Figure 1).

The area of East Sinjai Sub-District covers 7,188 ha and is bordered by North Sinjai Sub-District to the north, Bone Bay to the east, Tellu Limpoe Sub-District to the south, and North Sinjai and Central Sinjai Sub-Districts to the west. The population was 30,550 people in 2016, and most of the inhabitants live in the coastal area and work as fishermen and shrimp farmers (BPS Kabupaten Sinjai, 2017).

The coastal area is covered by mangroves, seagrass beds, coral reefs, aquaculture ponds, settlements, and ecotourism areas (Malik et al., 2020). The area covered by mangroves in 2017 was 761 ha (which is about 77% of the total mangrove area in Sinjai District) and distributed among five villages, including Samatarang (StV), Tongketongke (TtV), Panaikang (PkV), Pasimarannu (PrV), and Sanjai (SjV) (Malik & Rahim, 2017). Mangroves in these areas grow in a fringing hydrogeomorphic environment, and the tidal regime in this area is semi-diurnal. On most days, there are two tidal cycles, often with not significantly different amplitudes), with the tidal range being about 122 cm (Malik & Rahim, 2017). Mangroves soils are mostly clayey with rich organic matter content and are associated with dominant mangrove vegetation from *Rhizophora* sp. (Suharti et al., 2016; Malik et al., 2020). Moreover, the climatic situation in these areas is characterized by the average annual rainfall of 2697 mm with 211 rainy days in 2017, whereas the mean daily air temperature is between 21 °C and 32 °C (BPS Kabupaten Sinjai, 2017).

However, the expansion of aquaculture ponds through the clearing mangroves in these areas began in the 1930s




Figure 1. Study area: Sinjai District, South Sulawesi, Indonesia
The distance to the capital of South Sulawesi, Makassar City, is 220 km

(Amri, 2008), but the most significant development occurred in the last three decades (Malik & Rahim, 2017). The mangrove reforestation began in 1984 at the initiative of the local community (Amri, 2008). Since the early 2000s, this action has flourished to protect the area from coastal erosion and storms and provide land for the ecotourism area. However, the disturbances to the mangrove area continue, mainly due to the expansion of aquaculture ponds (Malik & Rahim, 2017).

1.2. Data collection

Fieldwork was conducted in April 2017 in five mangrove areas (StV, TtV, PkV, PrV, and SjV). We used a transect line method (Malik et al., 2015), the length of which depended on the thickness of the mangrove forest from seaward to landward at each site. We established three 10×10 m sampling plots for each transect using a tape measure and plastic ropes (Malik et al., 2015). Within each plot, we collected 15 soil cores at depth of 0–15 cm, 15–30 cm, and 30–50 cm. We inserted the sediment core (stainless steel Eijkelkamp gouge auger) vertically into the soil until the top of the sampler was level with the soil surface. When

the soil core was extracted, we measured the depth of the gathered samples using a measuring tape and collected a sub-sample with a length of 5 cm from the midpoint each of intervals depth. Soil samples were extracted, stored in labeled plastic bags, and taken to the laboratory for soil analysis (Kauffman & Donato, 2012).

1.3. Sample analysis

Soil analysis was performed using the Loss on Ignition (% LOI) method by burning the soil sample at high temperatures (Kauffman & Donato, 2012). We placed a 118.73 cm³ soil sample from different depths in each plot in a pre-weighed ceramic crucible and put them in the drying oven at a temperature of 60 °C for 72 hours to maintain a constant dry matter. The soil samples were carefully broken into smaller pieces to accelerate the drying process. The value of oven-dried soil samples was weighed and subtracted from the net weight of the ceramic crucible. We also placed 20 g of oven-dried sub-sample from each sample in a muffle furnace at 540 °C for five hours to ignite it (Kauffman & Donato, 2012).

To calculate the bulk density of mangrove soil (SBD), we divided the mass of the oven-dry soil sample by the volume of the pre-dried sample (Equation (1)) (Kauffman & Donato, 2012):

$$SBD \left(g \text{ cm}^{-3} \right) = \frac{\text{oven-dry sample mass (g)}}{\text{pre-dried soil sample volume (cm}^3\text{)}}, \quad (1)$$

where: volume of pre-dried soil sample = 118.73 cm³.

The % LOI, which indicated the soil organic matter content, was estimated by using Equation (2) (Kauffman & Donato, 2012):

$$\% \text{ LOI} = \frac{[\text{dry mass before combustion (mg)} - \text{dry after combustion (mg)}]}{\text{dry mass before combustion (mg)}} \times 100. \quad (2)$$

To calculate the soil organic carbon concentration/SOCC (% C_{org}), we also used Equation (3) as proposed by (Kauffman et al., 2011):

$$\text{SOCC} = 0.415 \times \% \text{ LOI} + 2.89. \quad (3)$$

To determine the soil organic carbon density (SOCD) and the soil organic carbon (SOC) stocks at each sampled depth, we used Equations (4) and (5), respectively (Kauffman & Donato, 2012):

$$\text{SOCD (g C cm}^{-3}\text{)} = \text{SBD (g cm}^{-3}\text{)} \times (\% \text{ C}_{\text{org}}/100); \quad (4)$$

$$\text{SOC (Mg C ha}^{-1}\text{)} = \text{SBD (g cm}^{-3}\text{)} \times \text{soil depth interval (cm)} \times \text{SOCC}. \quad (5)$$

Finally, statistical tests using One-Way Analysis of Variance (ANOVA) were performed to compare different soil properties within depth intervals and carbon stock at different sites.




Figure 2. Soil properties: a) SBD, b) SOCC, c) SOCD, and d) SOC in the mangrove forests of Sinjai District, South Sulawesi

2. Results

Soil property values and trends (SBD, SOCC, SOCD, and SOC) of mangrove areas of Sinjai District, South Sulawesi, Indonesia, are summarized in Figures 2a–2d and Table S1.

We found that the SBD value across sites and soil depths ranged from $0.42 \pm 0.13 \text{ g cm}^{-3}$ at a depth of 0–15 cm at the SjV site to $0.77 \pm 0.06 \text{ g cm}^{-3}$ at a depth of 15–30 cm at the TtV site, or the mean value of all ranges was $0.60 \pm 0.05 \text{ g cm}^{-3}$. The SBD value with interval depth had an increasing trend at all sites and showed a significant difference ($p < 0.05$) (Figure 2a, Table S1).

The lowest SOCC value of $11.74 \pm 0.50\%$ was observed at a depth of 30–50 cm at site SjV, whereas the highest value of $14.54 \pm 0.20\%$ was observed at the 0–15 cm at site StV, resulting in the mean SOCC of all sites $13.70 \pm 0.22\%$. The SOCC value of each site tended to decrease with depth but did not differ significantly ($p = 0.18$) (Figure 2b, Table S1). The mean SOCD value of soil depth was significantly different at all sites ($p < 0.05$), ranging from the mean value of 0.06 ± 0.0004 at site SjV to 0.10 ± 0.0004 at site TtV (Figure 2c, Table S1).

In addition, we estimated the total SOC stock at the five study sites to be $2,065.50 \text{ Mg C}$, resulting in a mean SOC stock of $137.70 \pm 12.37 \text{ Mg C ha}^{-1}$, with the lowest value at 0–15 cm depth at site SjV ($88.28 \pm 26.14 \text{ Mg C ha}^{-1}$) and the highest SOC stock at a soil depth of 30–50 cm at site TtV ($210.35 \pm 18.80 \text{ Mg C ha}^{-1}$). The mean of SOC stock between sites was not significantly different ($p = 0.26$) (Figure 2d, Table S1).

3. Discussion

The mean value of SBD ($0.60 \pm 0.05 \text{ g cm}^{-3}$) in this area was lower than the SBD values of mangroves in many mangrove areas in the world (0.73 g cm^{-3} – 1.42 g cm^{-3} , Hossain & Nuruddin, 2016), but higher than in Indo-Pacific and several mangrove areas in Indonesia (0.35 – 0.55 g cm^{-3}) as reported by Donato et al. (2011) and in Can Gio Mangrove Biosphere Reserve (CGMBR), Mekong Delta, Vietnam (0.52 g cm^{-3}) as detected by Nam et al. (2016). The value of SBD increased with increasing soil depth at all sites and was significantly different ($p < 0.05$) (Figure 2a, Table S1). It is due to lower organic matter content, soil accumulation, and compaction due to the weight of the overlying layer. The trend showed similarities with other mangrove areas in Indonesia (Donato et al., 2011) and in the Mekong Delta, Vietnam (Nam et al., 2016), as well as in Shenzhen Bay, China (Lunstrum & Chen, 2014). However, the SOCC tended to decrease with increasing depth (Figure 2b), which attributes to the high SBD that indicating higher soil density and small soil pores (Lunstrum & Chen, 2014). A similar trend was also observed in the Indo-Pacific and several mangrove areas in Indonesia, as reported by Donato et al. (2011) and consistent with Kauffman et al. (2014) in the Dominican Republic.

SOCD was determined by multiplying the value of SBD and organic carbon content ($\% C_{\text{org}}$) at each soil depth. Dorji et al. (2014) stated that SOCD is required for carbon accounting, accumulation, budgeting, and developing carbon sequestration strategies. The mean SOCD ($0.08 \pm 0.01 \text{ g C cm}^{-3}$, Table S1) in this area was higher

compared to several mangrove areas in the world, including the western and eastern Atlantic and Pacific coasts, the Indian Ocean, the Mediterranean Sea, and the Gulf of Mexico ($0.055 \pm 0.004 \text{ g C cm}^{-3}$), as demonstrated by Chmura et al. (2003). However, this value was lower than the mangrove rehabilitation site in Bali (0.13 g C cm^{-3}), as Mahasani et al. (2016) reported. SOCD tended to increase along with soil depth (Figure 2c) and influenced SBD and SOCC content (Dariah et al., 2012). In contrast, Dorji et al. (2014) and Cuc et al. (2009) found that SOCD decreases with soil depth. The inconsistent trend of SOCD values with the soil depth is due to the fact that SBD and SOCC values may vary with soil depth and location (Howard et al., 2014).

The mean SOC stock was $413.10 \pm 12.37 \text{ Mg C ha}^{-1}$ in this area (Table S1). With a total mangrove area of 761 ha in 2017 (Malik & Rahim, 2017), the total stock of SOC in this area is estimated to be approximately 0.31 Tg C. Being a mangrove fringe area, the accumulation of autochthonous SOC in this mangrove area is influenced by tidal hydrodynamic which includes biogeochemical and physical processes. Tidal inundation brings abundant sediment, encouraging lateral deposition of mud due to reduced water flow within the mangroves (Rovai et al., 2018). Besides, warm temperatures and abundant rainfall in the area are associated with primary productivity and decomposition can affect the levels of SOC (Ontl & Schulte, 2012). However, the mean SOC stocks in this area were lower compared to other mangrove areas in Indonesia, such as Java and Kalimantan (Donato et al., 2011), Sumatra (Alongi et al., 2015), and West Papua and Papua (Taberima et al., 2014). The loss of mangroves, mainly due to aquaculture development, affects the decline of SOC in the area. Converting mangroves to aquaculture ponds can reduce biomass carbon stocks by an average of 83% and soil carbon stocks by 52% (Sasmito et al., 2019; Friess et al., 2020).

Based on the soil depth layer per site, the mean SOC stock tended to increase from the topsoil (Figure 2d). This pattern was similar to mangrove areas in West Papua and Papua, and Micronesian mangrove forests, as demonstrated by Taberima et al. (2014) and Kauffman et al. (2011), respectively. The mean SOC stock value difference between soil layers was continuously high throughout the soil layer (Kauffman et al., 2011).

Almost half of the total values of SOC at each site are at the 30–50 cm depth (Figure 2d, Table S1). Donato et al. (2011) showed that 49–98% of the carbon is stored at a depth of 0.5 m to 3 m. Taberima et al. (2014) found that most SOC in West Papua and Papua were found at depth 10 cm to 200 cm, but the highest value was above 100 cm. Mangrove root growth at a depth of 50 cm plays a significant role in accumulating SOC (Nguyen et al., 2004). Ontl and Schulte (2012) found that the levels of SOC are primarily derived from root biomass and litter deposited by plants. Plant roots contribute SOC directly and indirectly through root growth and death and the transfer of carbon-rich compounds from roots to the soil.

Conclusions

The study has demonstrated the potential stock of soil organic carbon (SOC) from the mangrove area in Sinjai District, South Sulawesi, Indonesia. The mean SOC stock was $413.10 \pm 12.37 \text{ Mg C ha}^{-1}$. However, the observed mean stock of SOC from this area was lower than that studied in several mangrove areas in Indonesia due to human-induced exploitation, mainly for aquaculture expansion.

Promoting wise use and restoration of mangroves is a top priority for sustainable management, preserving SOC presently held, and rebuilding SOC lost during exploitation occurs. In addition, a better understanding of mangrove carbon stocks and human-induced pressures by stakeholders and decision-makers in this region will help them conserve and manage this vital ecosystem and mitigate climate change.

Acknowledgements

This research was funded by PNPB Pascasarjana Universitas Negeri Makassar under contract grant number: SP DIPA-042.01: 2.400964/2017, December 7, 2016. We sincerely thank the laboratory of the Department of Geography, Faculty of Mathematics and Natural Sciences Universitas Negeri Makassar for the analysis of mangrove soil. We thank the Government of South Sulawesi and Sinjai District for the research permit and other support for this work.

Conflicts of interest

The authors declare no conflicts of interest.

Author contributions

All authors wrote the manuscript.

References

- Abino, A. C., Castillo, J. A. A., & Lee, Y. J. (2014). Species diversity, biomass, and carbon stock assessment of natural mangrove forest in Palawan, Philippines. *Pakistan Journal of Botany*, *46*, 1955–1962.
- Alongi, D. M. (2012). Carbon sequestration in mangrove forests. *Carbon Management*, *3*, 313–322. <https://doi.org/10.4155/cmt.12.20>
- Alongi, D. M. (2020). Global significance of mangrove blue carbon in climate change mitigation. *Sci*, *2*(3), 67. <https://doi.org/10.3390/sci2030067>
- Alongi, D. M., Murdiyarso, D., Fourqurean, J. W., Kauffman, J. B., Hutahaen, A., Crooks, S., Lovelock, C. E., Howard, J., Herr, D., Fortes, M., Pidgeon, E., & Wagey, T. (2015). Indonesia's blue carbon: a globally significant and vulnerable sink for seagrass and mangrove carbon. *Wetland Ecology Management*, *24*, 3–13. <https://doi.org/10.1007/s11273-015-9446-y>
- Amri, A. (2008). Land property rights and coastal resource management: A perspective of community based mangrove conservation in Indonesia. In *12th Biennial Global Conference of the International Association for the Study of the Commons*

- (pp. 1–17), University of Gloucestershire. https://dlc.dlib-indiana.edu/dlc/bitstream/handle/10535/2247/Amri_201101.pdf?sequence=1&isAllowed=y
- BPS Kabupaten Sinjai. (2017). *Kabupaten Sinjai dalam angka 2016*. Badan Pusat Statistik Kabupaten Sinjai. <https://sinjaikab.bps.go.id/frontend/index.php/publikasi/128>
- Bukoski, J. J., Elwin, A., MacKenzie, R. A., Sharma, S., Purbopuspito, J., Kopania, B., Apwong, M., Poolsiri, R., & Potts, M. D. (2020). The role of predictive model data in designing mangrove forest carbon programs. *Environmental Research Letters*, 15(8), 084019. <https://doi.org/10.1088/1748-9326/ab7e4e>
- Chen, G., Azkab, M. H., Chmura, G. L., Chen, S., Sastrosuwondo, P., Ma, Z., Dharmawan, I. W. E., Yin, X., & Chen, B. (2017). Mangroves as a major source of soil carbon storage in adjacent seagrass meadows. *Scientific Reports*, 7(1), 1–10. <https://doi.org/10.1038/srep42406>
- Chmura, G. L., Anisfeld, S. C., Cahoon, D. R., & Lynch, J. C. (2003). Global carbon sequestration in tidal, saline wetland soils. *Global Biogeochemical Cycles*, 17(4), 1111. <https://doi.org/10.1029/2002GB001917>
- Cuc, N. T. K., Ninomiya, I., Long, N. T., Tri, N. H., Tuan, M. S., & Hong, P. N. (2009). Belowground carbon accumulation in young *Kandelia candel* (L.) Blanco plantations in Thai Binh River Mouth, Northern Vietnam. *International Journal of Ecology & Development*, 12(W09), 107–117.
- Dariah, A., Maftuah, E., & Maswar. (2012). Karakteristik lahan gambut. In *Panduan pengelolaan berkelanjutan lahan gambut terdegradasi* (pp. 17–29). Balai Penelitian Tanah (Balittanah), Badan Penelitian dan Pengembangan Pertanian, Kementerian Pertanian. http://balittanah.litbang.pertanian.go.id/ind/dokumentasi/panduan%20gambut/03ai_karakteristik.pdf
- Donato, D. C., Kauffman, J. B., Murdiyarso, D., Kurnianto, S., Stidham, M., & Kanninen, M. (2011). Mangroves among the most carbon-rich forests in the tropics. *Nature Geoscience*, 4, 293–297. <https://doi.org/10.1038/ngeo1123>
- Dorji, T., Odeh, I. O. A., & Field, D. J. (2014). Vertical distribution of soil organic carbon density in relation to land use/cover, altitude and slope aspect in the Eastern Himalayas. *Land*, 3, 1234–1250. <https://doi.org/10.3390/land3041232>
- Duncan, C., Primavera, J. H., Petteorelli, N., Thompson, J. R., Loma, R. J. A., & Koldewey, H. J. (2016). Rehabilitating mangrove ecosystem services: A case study on the relative benefits of abandoned pond reversion from Panay Island, Philippines. *Marine Pollution Bulletin*, 109, 772–782. <https://doi.org/10.1016/j.marpolbul.2016.05.049>
- Friess, D. A., Krauss, K. W., Taillardat, P., Adame, M. F., Yando, E. S., Cameron, C., Sasmito, S. D., & Sillanpää, M. (2020). Mangrove blue carbon in the face of deforestation, climate change, and restoration. *Annual Plant Reviews online*, 427–456. <https://doi.org/10.1002/9781119312994.apr0752>
- Giesen, W., Baltzer, M., & Baruadi, R. (Eds.). (1991). *Integrating conservation with land-use development in wetlands of South Sulawesi*. PHPA/AWB (Asian Wetland Bureau), Bogor.
- Hamilton, S. E., & Friess, D. (2018). Global carbon stocks and potential emissions due to mangrove deforestation from 2000 to 2012. *Nature Climate Change*, 4, 240–244. <https://doi.org/10.1038/s41558-018-0090-4>
- Hong Tinh, P., Thi Hong Hanh, N., Van Thanh, V., Sy Tuan, M., Van Quang, P., Sharma, S., & MacKenzie, R. A. (2020). A comparison of soil carbon stocks of intact and restored mangrove forests in northern Vietnam. *Forests*, 11(6), 660. <https://doi.org/10.3390/f11060660>
- Hopkinson, C. S., Cai, W. J., & Hu, X. (2012). Carbon sequestration in wetland dominated coastal systems – a global sink of rapidly diminishing magnitude. *Current Opinion in Environmental Sustainability*, 4, 186–194. <https://doi.org/10.1016/j.cosust.2012.03.005>
- Hossain, M. D., & Nuruddin, A. A. (2016). Soil and mangrove: A review. *Journal of Environmental Science and Technology*, 9, 198–207. <https://doi.org/10.3923/jest.2016.198.207>
- Howard, J., Hoyt, S., Isensee, K., Telszewski, M., & Pidgeon, E. (Eds.). (2014). *Coastal blue carbon: Methods for assessing carbon stocks and emissions factors in mangroves, tidal salt marshes, and seagrasses*. Conservation International, Intergovernmental Oceanographic Commission of UNESCO, International Union for Conservation of Nature, Arlington, Virginia, USA.
- Intergovernmental Panel on Climate Change. (2014). *2013 Supplement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories: Wetlands*. <http://www.ipcc-nggip.iges.or.jp/public/wetlands/>
- Jennerjahn, T. C. (2020). Relevance and magnitude of ‘Blue Carbon’ storage in mangrove sediments: Carbon accumulation rates vs. stocks, sources vs. sinks. *Estuarine, Coastal and Shelf Science*, 247, 107027. <https://doi.org/10.1016/j.ecss.2020.107027>
- Kauffman, J. B., & Donato, D. C. (2012). *Protocols for the measurement, monitoring and reporting of structure, biomass and carbon stocks in mangrove forests* (Working Paper No. 86). CIFOR, Bogor, Indonesia.
- Kauffman, J. B., Heider, C., Cole, T. G., Dwire, K. A., & Donato D. C. (2011). Ecosystem carbon stocks of Micronesia mangrove forests. *Wetlands*, 31, 343–352. <https://doi.org/10.1007/s13157-011-0148-9>
- Kauffman, J. B., Heider, C., Norfolk, J., & Payton, F. (2014). Carbon stocks of intact mangroves and carbon emissions arising from their conversion in Dominican Republic. *Ecological Application*, 24, 518–527. <https://doi.org/10.1890/13-0640.1>
- Kusumaningtyas, M. A., Hutahaean, A. A., Fischer, H. W., Pérez-Mayo, M., Ransby, D., & Jennerjahn, T. C. (2019). Variability in the organic carbon stocks, sources, and accumulation rates of Indonesian mangrove ecosystems. *Estuarine, Coastal and Shelf Science*, 218, 310–323. <https://doi.org/10.1016/j.ecss.2018.12.007>
- Lunstrum, A., & Chen, L. (2014). Soil carbon stocks and accumulation in young mangrove forests. *Soil Biology and Biochemistry*, 75, 223–232. <https://doi.org/10.1016/j.soilbio.2014.04.008>
- Mahasani, I. G. A. I., Karang, I. W. G., & Hendrawan, I. G. (2016, Juli 27). Karbon organik di bawah permukaan tanah pada kawasan rehabilitasi mangrove, taman hutan rakyat Nguurah Rai, Bali. In *Prosiding Seminar Nasional Kelautan 2016* (pp. 33–42), Universitas Trunojoyo Madura. http://ilmuke-lautan.trunojoyo.ac.id/wp-content/uploads/2016/08/6_Prosiding_semnaskel_2016.pdf
- Malik, A., & Rahim, A. (2017). *Assessment of potentials for payment for mangrove ecosystem services in South Sulawesi Indonesia* (Final Report DIPA Biotrop 2017). Southeast Asian Regional Centre for Tropical Biology (SEAMEO BIOTROP), Bogor, West Java.
- Malik, A., Fensholt, R., & Mertz, O. (2015). Mangrove exploitation effects on biodiversity and ecosystem services. *Biodiversity & Conservation*, 24, 3543–3557. <https://doi.org/10.1007/s10531-015-1015-4>
- Malik, A., Jalil, A. R., Arifuddin, A., & Syahmuddin, A. (2020). Biomass carbon stocks estimation in the mangrove rehabilitated area of Sinjai District, South Sulawesi, Indonesia. *Geography, Environment, Sustainability*, 13(3), 32–38. <https://doi.org/10.24057/2071-9388-2019-131>

- Malik, A., Mertz, O., & Fensholt, R. (2017). Mangrove forest decline: Consequences for livelihoods and environment in South Sulawesi. *Regional Environmental Change*, 17, 157–169. <https://doi.org/10.1007/s10113-016-0989-0>
- Murdiyarso, D., Purbopuspito, J., Kauffman, J. B., Warren, M., Sasmito, S., Donato, D., Manuri, S., Krisnawati, H., Taberima, S., & Kurnianto, S. (2015). The potential of Indonesian mangrove forests for global climate change mitigation. *Nature Climate Change*, 5, 1089–1092. <https://doi.org/10.1038/nclimate2734>
- Nam, V. N., Sasmito, S. D., Murdiyarso, D., Purbopuspito, J., & MacKenzie, R. A. (2016). Carbon stocks in artificially and naturally regenerated mangrove ecosystems in the Mekong Delta. *Wetlands Ecology Management*, 24, 231–244. <https://doi.org/10.1007/s11273-015-9479-2>
- Nguyen, H. T., Yoneda, R., Ninomiya, I., Harada, K., Van Dao, T., Sy, T. M., & Phan, H. N. (2004). The effects of stand-age and inundation on carbon accumulation in mangrove plantation soil in Namdinh, Northern Vietnam. *Tropics*, 14(1), 21–37. <https://doi.org/10.3759/tropics.14.21>
- Nóbrega, G. N., Ferreira, T. O., Artur, A. G., de Mondonça, E. S., de O. Leão, R. A., Teixeira, A. S., & Otero, X. L. (2015). Evaluation of methods for quantifying organic carbon in mangrove soils from semi-arid region. *Journal of Soils and Sediments*, 15, 282–291. <https://doi.org/10.1007/s11368-014-1019-9>
- Ontl, T. A., & Schulte, L. A. (2012). Soil carbon storage. *Nature Education Knowledge*, 3(10).
- Pendleton, L., Donato, D. C., Murray, B. C., Crooks, S., Jenkins, W. A., Siflet, S., Craft, C., Fourqurean, J. W., Kauffman, J. B., Marba, N., Megonigal, P., Pidgeon, E., Herr, D., Gordon, D., & Baldera, A. (2012). Estimating global “blue carbon” emissions from conversion and degradation of vegetated coastal ecosystems. *PLoS ONE*, 7, e43542. <https://doi.org/10.1371/journal.pone.0043542>
- Rahadian, A., Prasetyo, L. B., Setiawan, Y., & Wikantika, K. (2019). Tinjauan historis data dan informasi luas mangrove Indonesia (A historical review of data and information of Indonesian Mangroves Area). *Media Konservasi*, 24(2), 163–178. <https://doi.org/10.29244/medkon.24.2.163-178>
- Richards, D. R., & Friess, D. A. (2016). Rates and drivers of mangrove deforestation in Southeast Asia, 2000–2012. *Proceedings of the National Academy of Sciences of the United States of America*, 113, 344–349. <https://doi.org/10.1073/pnas.1510272113>
- Rovai, A. S., Twilley, R. R., Castañeda-Moya, E., Riul, P., Cifuentes-Jara, M., Manrow-Villalobos, M., Horta, P. A., Simonassi, J. C., Fonseca, A. L., & Pagliosa, P. R. (2018). Global controls on carbon storage in mangrove soils. *Nature Climate Change*, 8(6), 534–538. <https://doi.org/10.1038/s41558-018-0162-5>
- Sasmito, S. D., Sillanpää, M., Hayes, M. A., Bachri, S., Saragi-Sasmito, M. F., Sidik, F., Hanggara, B. B., Mofu, W. Y., Rumbiak, V. I., Taberima, S., & Nugroho, J. D. (2020a). Mangrove blue carbon stocks and dynamics are controlled by hydrogeomorphic settings and land-use change. *Global Change Biology*, 26(5), 3028–3039. <https://doi.org/10.1111/gcb.15056>
- Sasmito, S. D., Kuzyakov, Y., Lubis, A. A., Murdiyarso, D., Hutley, L. B., Bachri, S., Friess, D. A., Martius, C., & Borchard, N. (2020b). Organic carbon burial and sources in soils of coastal mudflat and mangrove ecosystems. *Catena*, 187, 104414. <https://doi.org/10.1016/j.catena.2019.104414>
- Sasmito, S. D., Taillardat, P., Clendenning, J. N., Cameron, C., Friess, D. A., Murdiyarso, D., & Hutley, L. B. (2019). Effect of land-use and land-cover change on mangrove blue carbon: A systematic review. *Global Change Biology*, 25(12), 4291–4302.
- Sharma, S., MacKenzie, R. A., Tieng, T., Soben, K., Tulyaswan, N., Resanond, A., Blate, G., & Litton, C. M. (2020). The impacts of degradation, deforestation and restoration on mangrove ecosystem carbon stocks across Cambodia. *Science of the Total Environment*, 706, 135416. <https://doi.org/10.1016/j.scitotenv.2019.135416>
- Sidik, F., & Lovelock, C. E. (2013). CO₂ efflux from shrimp ponds in Indonesia. *PLoS ONE*, 8, e66329. <https://doi.org/10.1371/journal.pone.0066329>
- Soper, F. M., MacKenzie, R. A., Sharma, S., Cole, T. G., Litton, C. M., & Sparks, J. P. (2019). Non-native mangroves support carbon storage, sediment carbon burial, and accretion of coastal ecosystems. *Global Change Biology*, 25(12), 4315–4326. <https://doi.org/10.1111/gcb.14813>
- Suharti, S., Darusman, D., Nugroho, B., & Sundawati, L. (2016). Economic valuation as basis for sustainable mangrove resource management. A case in East Sinjai, South Sulawesi. *Journal of Tropical Forest Management*, 22, 12–23. <https://doi.org/10.7226/jtfm.22.1.13>
- Taberima, S., Nugroho, Y. D., & Murdiyarso, D. (2014, September 17–18). *The distribution of carbon stock in selected mangrove ecosystem of wetlands Papua: Bintuni, Teminabuan, and Timika Eastern Indonesia* [Conference presentation]. International Conference on Chemical, Environment & Biological Sciences (CEBS-2014), Kuala Lumpur, Malaysia.

Supplementary information

**THE POTENTIAL SOIL ORGANIC CARBON STOCKS IN MANGROVE
AREAS OF SINJAI DISTRICT, SOUTH SULAWESI, INDONESIA**

Abdul MALIK, Muhammad ICHSAN ALI, Suwardi ANNAS, Abdul Rasyid JALIL,
Restu Utami MULYA, Konstantina GRAVANI

Table S1. Soil properties and carbon stock in the mangrove area of Sinjai District South Sulawesi Indonesia

Site	Depth	n	SBD (g cm ⁻³)		SOCC (%)		SOCD (g C cm ⁻³)		SOC (Mg C ha ⁻¹)	
			Mean	SE	Mean	SE	Mean	SE	Mean	SE
StV	0–15	3	0.62	0.07	14.54	0.20	0.09	0.01	134.87	16.11
	15–30	3	0.67	0.04	13.92	0.40	0.09	0.005	139.56	7.32
	30–50	3	0.73	0.04	13.74	0.39	0.10	0.003	199.00	5.30
	Mean	–	0.67	0.03	14.06	0.24	0.09	0.003	157.81	20.64
	Total	9	–		–		–		473.43	
	p-value		0.34		0.29		0.64		0.01	
TtV	0–15	3	0.65	0.14	14.34	0.47	0.09	0.02	139.65	23.48
	15–30	3	0.75	0.12	14.12	0.37	0.11	0.01	159.13	19.77
	30–50	3	0.77	0.06	13.61	0.35	0.11	0.01	210.35	18.80
	Mean	–	0.72	0.04	14.02	0.22	0.10	0.004	169.71	21.08
	Total	9	–		v		–		509.13	
	p-value		0.62		0.46		0.74		0.12	
PkV	0–15	3	0.48	0.10	13.74	1.36	0.06	0.01	96.20	11.85
	15–30	3	0.54	0.14	13.36	1.38	0.07	0.02	112.08	37.06
	30–50	3	0.75	0.08	13.05	1.45	0.10	0.02	199.37	38.79
	Mean	–	0.59	0.08	13.38	0.20	0.08	0.01	135.88	32.07
	Total	9	–		–		–		407.65	
	p-value		0.28		0.94		0.44		0.12	
PmV	0–15	3	0.44	0.10	14.03	0.37	0.09	0.00	91.53	18.93
	15–30	3	0.56	0.05	13.96	0.68	0.08	0.01	119.15	16.68
	30–50	3	0.59	0.07	14.10	0.60	0.08	0.01	167.36	24.90
	Mean	–	0.53	0.05	14.03	0.04	0.08	0.002	126.01	22.16
	Total	9	–		–		–		378.04	
	p-value		0.40		0.99		0.89		0.10	

End of Table S1

Site	Depth	n	SBD (g cm ⁻³)		SOCC (%)		SOCD (g C cm ⁻³)		SOC (Mg C ha ⁻¹)	
			Mean	SE	Mean	SE	Mean	SE	Mean	SE
SjV	0–15	3	0.42	0.13	14.10	0.12	0.06	0.02	88.28	26.14
	15–30	3	0.46	0.04	13.13	0.25	0.06	0.005	90.22	7.28
	30–50	3	0.51	0.02	11.74	0.50	0.06	0.002	118.75	3.27
	Mean	–	0.46	0.03	12.99	0.68	0.06	0.0004	99.08	9.85
	Total	9	–		–		–		297.25	
	p-value		0.74		0.01		1.00		0.37	
p-value			0.02		0.18		0.002		0.26	
Grand mean		–	0.60	0.05	13.70	0.22	0.08	0.01	413.10	12.37
Grand total		45							2065.50	

Note: StV: Samataring village; TtV: Tongke-tongke village; PkV: Panaikang village; PmV: Pasimarannu village; SjV: Sanjai village; n: number of soil sample; SBD: soil bulk density; SOCC: soil organic carbon concentration; SOCD: soil organic carbon density; SOC: soil organic carbon.

● **7% Overall Similarity**

Top sources found in the following databases:

- 5% Internet database
- 3% Publications database
- Crossref database
- Crossref Posted Content database
- 3% Submitted Works database

TOP SOURCES

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

1	ges.rgo.ru Internet	<1%
2	Vien Ngoc Nam, Sigit D. Sasmito, Daniel Murdiyarto, Joko Purbopuspito... Crossref	<1%
3	cifor.org Internet	<1%
4	Murdiyarto, Daniel, Joko Purbopuspito, J. Boone Kauffman, Matthew ... Crossref	<1%
5	Alexandru Ioan Cuza University of Iasi on 2020-10-26 Submitted works	<1%
6	thepharmajournal.com Internet	<1%
7	Abdul Malik, Rasmus Fensholt, Ole Mertz. "Mangrove exploitation effe... Crossref	<1%
8	Oregon State University on 2017-04-27 Submitted works	<1%

9	bibliotekanauki.pl Internet	<1%
10	pdf.usaid.gov Internet	<1%
11	suaire.suanet.ac.tz Internet	<1%
12	nature.com Internet	<1%
13	pmcarbono.org Internet	<1%
14	tropicalplantresearch.com Internet	<1%
15	University of Melbourne on 2019-11-08 Submitted works	<1%
16	wrc.org.za Internet	<1%
17	eprints.ulm.ac.id Internet	<1%
18	pdfs.semanticscholar.org Internet	<1%
19	scholarspace.manoa.hawaii.edu Internet	<1%

● Excluded from Similarity Report

- Bibliographic material
- Cited material
- Manually excluded sources
- Quoted material
- Small Matches (Less than 10 words)
- Manually excluded text blocks

EXCLUDED SOURCES

journals.vilniustech.lt	7%
Internet	
doaj.org	7%
Internet	
jau.vgtu.lt	<1%
Internet	
jest.vgtu.lt	<1%
Internet	

EXCLUDED TEXT BLOCKS

of Civil and Planning, Faculty of Engineering, Universitas Negeri Makassar, Makassar
www.ijsr.net

South Sulawesi
e-journal.unair.ac.id

of Geosciences and Natural Resource Management, Faculty of Science, University ...
curis.ku.dk

CARBON STOCKS IN MANGROVE
ges.rgo.ru

E-mail

Universitas Negeri Semarang on 2022-08-31

1. Materials and methods1.1. Study areaThe study area is

egijournals.researchcommons.org

carbon stocks in mangrove

ges.rgo.ru

LOI = $\frac{\text{dry mass before combustion}(\text{mg}) - \text{dry after combustion}(\text{mg})}{\text{dry mass before combustion}(\text{mg})}$

King's College on 2022-05-16

carbon stocks in mangrove areas of Sinjai District, South Sulawesi

ges.rgo.ru

the Department of

ges.rgo.ru

Conflicts of interestThe authors declare no conflicts of interest.Author contributi...

onlinelibrary.wiley.com

Journal of Environmental Engineering and Landscape ManagementISSN 1648-689...

curis.ku.dk

1Department of

Universiti Teknologi Malaysia on 2018-09-04
