

The abundance and diversity of phosphatesolubilizing bacteria in integrated recirculating aquaculture systems

by Dahlia, Ardiansyah, Indrayani, Firmansyah Bin A. Jabbar


Submission date: 11-Apr-2023 06:10AM (UTC+0800)

Submission ID: 2060933620

File name: Indrayani-AACL_Bioflux_14_5_3055-3067_2021.pdf (120.84K)

Word count: 7480

Character count: 41420



The abundance and diversity of phosphate-solubilizing bacteria in integrated recirculating aquaculture systems as a function of harvesting regime of duckweed (*Lemna minor* L.)

¹Dahlia, ¹Ardiansyah, ²Indrayani, ³Firmansyah Bin A. Jabbar

12

¹ Pangkep State Polytechnic of Agriculture, South Sulawesi, Indonesia; ² Makassar State University, Makassar, South Sulawesi, Indonesia; ³ Universitas Sulawesi Barat, West Sulawesi, Indonesia. Corresponding author: Ardiansyah, ardi_kimsan@yahoo.com

43

Abstract. The present study aimed to determine the effects of different harvesting regimes of duckweed (*Lemna minor* Linnaeus) on the abundance and diversity of phosphate-solubilizing bacteria (PSB) in integrated recirculating aquaculture systems (IRAS). The study was conducted in twelve independent IRAS, where four duckweed harvesting regimes of two, four, and six-day harvest intervals tested the abundance and diversity of PSB. After 36 days, ten selected isolates belonging to five different genera were biochemically characterized. These isolates and their respective genera were: PSB1, PSB3, and PSB7 belonging to *Bacillus*, PSB17, and PSB26 to *Pseudomonas*, and PSB15, PSB29, PSB30, PSB32, and PSB35 to *Chryseobacterium*, *Micrococcus*, *Azotobacter*, *Acinetobacter*, and *Achromobacter*, respectively. *In vitro*, *Bacillus cereus* had higher solubilization efficiency and phosphatase activity. Duckweed harvested once every two days produced the highest duckweed biomass, however, harbored fewer bacteria, while six days of harvesting generated the highest duckweed growth index but lower bacterial diversity than the duckweed harvested every four days. The highest Shannon diversity index was obtained with a four-day harvest regime, while the lowest was when the duckweed was not harvested. A positive correlation ($R^2 > 0.65$) between the specific growth rate (SGR) and biomass of harvested duckweed and the diversity of PSB was found with a four-day harvest regime. Harvesting duckweed every four days provides desirable conditions for the attachment and growth of bacteria, thus increasing the abundance and diversity of PSB. These findings suggest the efficiency of removing excessive phosphorus and the associated PSB by duckweed and increase the productivity of IRAS.

Key Words: barramundi, biofilter, duckweed, integrated recirculating aquaculture system, phosphate-solubilizing bacteria.

Introduction. Aquaculture effluents with a high concentration of phosphorus (P) and nitrogen (N) contribute to eutrophication (Kawasaki et al 2016) and environmental degradation (Herath & Satoh 2015). Although recirculating aquaculture systems (RAS), considerably reduce the effluent load, significant amounts of N and P are still discharged through the protein-skimming mechanism (Martins et al 2010). Excessive P is often found in RAS effluents due to uneaten feed (Verdegem 2013). The removal of P to an acceptable level by various chemical methods is expensive (Jorgensen et al 2012) and requires high chemical dosages (Sun et al 2019). Even though the chemical removal of P rapidly improves the effluent quality (Zhao & Zhang 2017) more than the multi-step biological process, the chemical removal of P produces more sludge than its biological removal (Ruzhitskaya & Gogina 2017).

In any integrated RAS (IRAS), a substantial amount of organic and inorganic matter from uneaten feed and faecal matter accumulates and may subsequently be used by macrophytes. The phosphates could be reduced in an IRAS if the target species is cultivated with plants that utilised phosphates into valuable biomass (Turcois & Papenbrock 2014). The combination of fish culture with phototrophic and herbivorous organisms improves nutrient retention in the system (Turcois & Papenbrock 2014). The roots of various plant species incorporating phosphate-solubilizing bacteria (PSB) play ecophysiological role (Chen et al 2012). The PSB convert insoluble phosphates into

soluble forms (Alori et al 2017) and increase their availability to plants. The plants, in turn, may supply root-borne carbon (C) compounds - mainly sugars - that can be metabolized for bacterial growth (Gomes et al 2017). The microorganisms with phosphate-solubilizing potential rise the soluble phosphate availability and enhance plant growth by increasing the mineralization capacities of organic P and solubilizing precipitated phosphates (Sharma et al 2013), by increasing the supply of other trace elements such as iron, zinc, and by the production of plant growth-promoting regulators (Muszyńska & Labudda 2019).

Duckweed (*Lemna minor*) is known as a small free-floating aquatic plant belonging to the Lemnaceae family and it has been used for wastewater treatments due to its rapid multiplication (Nassar et al 2015; Paolacci et al 2018). A total of 70-80% total phosphate can be removed from wastewaters and orthophosphate removal rates between 83-95% in different types of wastewaters (Ozengin & Elmaci 2007). The use of duckweed species as a biofilter media in RAS decreases a significant amount of orthophosphate and total phosphorus in water and increases the growth of cultivated fish species (Velichkova & Sirakov 2013).

Since PSB are ubiquitous in the whole water column, they play an essential function in regulating the available concentration of inorganic phosphate in the water column for plant uptake (Shardendu et al 2012; Sushanta et al 2012; Cheng et al 2019). Thus, the abundance and diversity of the bacterial communities' act as PSB in RAS are affected by the type of biofilter and the media's surface area per unit volume, as biofilters affect the growth and structure of bacterial communities (Omri et al 2013).

Even though all intensive fish culture systems are a major source of P pollution, relatively few studies have been conducted on the removal of P from these systems (Dauda et al 2019). Previous research demonstrated that the harvesting frequency of duckweed affects the abundance and diversity of heterotrophic bacteria communities, which are involved in nitrogen removal from IRAS (Ardiansyah & Fotedar 2016). Hence, the use of appropriate management strategies including the manipulation of the abundance and diversity of the PSB communities in any IRAS are required to enhance P absorption by rhizoplants (Maitra et al 2015). This can be controlled by the quantity of biofilter media including the amount of duckweed present in real time; therefore, the present study aimed to evaluate the effects of different amounts and different harvest regimes of duckweed in real-time on the composition, abundance, and diversity of phosphate solubilizing, non-pathogenic bacteria that are involved in the removal of phosphates in IRAS.

Material and Method. This experiment was conducted in Curtin Aquatic Research Laboratory (CARL), Curtin University, Perth, Western Australia, from April to May 2017. The experimental setup and design were based on Ardiansyah & Fotedar (2016). As a source of phosphate, 185 mg of di-ammonium phosphate ((NH₄)₂ HPO₄) was added to the fish-rearing tanks. Twelve independent recirculating systems were set up, each consisting of three tanks: a fish-rearing tank (400-L circular tank), a biofilter tank (100-L circular tank), and a waste-collection tank (100-L circular plastic drum). Four duckweed harvesting regimes were employed during the 36-day experimental period. The four harvesting regimes were applied randomly by harvesting duckweed at three biofilter tanks at once (in 3 replications) after 2, 4, and 6 days, whereas the remaining three tanks were not harvested (control). Harvested duckweed was replaced with fresh duckweed that had been previously washed. The reduction in water volume caused by evaporation was compensated by adding new freshwater. The observations of duckweed health followed the standard method (Daud et al 2018). The increase in biomass of duckweed on the surface of the water was determined using the growth index (GI) on alternate days. GI of duckweed on the water surface was calculated every second day using the equation $GI = W_t/W_0$, where W_t = biomass at time t , W_0 = biomass at time 0, and t = number of days. While, the increase in biomass of duckweed per unit time of harvest regime was indicated by specific growth rates (SGR). $SGR = (\ln(W_f) - \ln(W_i) \times 100)/t$, where: $\ln(W_f)$ = the natural logarithm of the final weight, $\ln(W_i)$ = the natural logarithm of the initial weight, and t = time (days).

Isolation. Isolation, enumeration, and evaluation of the bacteria were performed following the procedure described in Ardiansyah & Fotedar (2016). A 10 g fresh weight of

duckweed was randomly collected using 40 mL sample jars containing 10 mL distilled water, and then kept at 4°C before analysis. The samples were homogenized by centrifugation at 17,000 g for five min. A 10 mL sample of the homogenate water was diluted with 90 mL of phosphate-buffered saline (PBS) and homogenized again for two minutes at 1900 g. The resulting homogenate was diluted ten times with PBS for 10 min. Then 0.1 mL of the diluted samples were plated onto Pikovskaya agar (PVK) medium in triplicate by the spread plate method (Elias et al 2016) and a conventional pour plate technique. Bacterial colonies were selected according to their morphological appearance and maximum surrounding halo zone as an indicator of phosphate solubilization after 120 h of incubation at 30°C (Zhang et al 2017). Colonies with distinct clear zones and different morphologies were picked and re-streaked onto fresh PVK medium containing 0.5% insoluble tricalcium phosphate and re-incubated at 30°C for 10 days.

Evaluation of phosphate solubilizing efficiency. Analysis of the PSB trait was made by determining the solubilization efficiency (%) of the selected isolates following the method of Pande et al (2017). The selected colonies were then removed from the plates and purified in triplicate on LB agar.

Enumeration. Enumeration of the PSB communities was determined using the total plate count method; the colonies were counted, and the concentration of bacteria present was expressed as CFU per mL for the attached bacteria, as previously reported by Blanco-Vargas et al (2020).

Phosphatase activity. Phosphatase activity was determined using 0.1 M disodium p-nitrophenyl as a substrate. For the assay, 3 mL of PSB culture on the PVK medium, 1 mL universal buffer (5x) adjusted to pH 6.5 (Scherer et al 2017), and 0.5 mL of the substrate were pipetted into a 20 mL reagent vial and incubated at 30°C for 90 min. The reaction was stopped by cooling at 2°C for 18 min. Then, 20 mL of 0.5 N NaOH was added, and the mixture was transferred to a 50 mL volumetric flask, where the volume was increased to 50 mL with distilled water. The p-nitrophenol (PNP) formed was determined by spectrophotometry at 398 nm (Schoebitz et al 2013). Controls were made in the same way using 20 µg mL⁻¹ disodium p-nitrophenyl.

Quantitative estimation of indole acetic acid (IAA). The procedures were performed according to Chaiharu & Lumyong (2011). The PSB isolates were inoculated in Czapek's aux broth and incubated at 28±2°C for 72 h. The culture was centrifuged at 3000 rpm for 15 min, and 2 mL of the supernatant was mixed with 1 mL of Salkowsky's reagent. The optical density was determined using a spectrophotometer at 535 nm.

Putative PSB identification. The putative PSB was identified by the biochemical method described in Bergey's Manual of Determinative Bacteriology (Garrity et al 2005). A variety of biochemical tests including Gram stain, catalase, oxidase, hydrogen sulfide production, gelatinase test, nitrate reduction, starch hydrolysis, citrate utilization, ornithine decarboxylase, indole, tyrosine hydrolysis, arginine dihydrolase, methyl-red and Voges-Proskauer, urease, glucose, fructose, lactose, maltose, mannose, sucrose, xylose, TSI agar test, and motility test, were conducted according to the standard procedure (Aditi et al 2017). API 20E system was used to confirm the identification of the PSB.

Bacteria diversity index. The bacteria diversity index assumes that an individual bacterium is randomly sampled from a large population so that all species are represented in the sample. The Shannon diversity index was used to determine the species diversity with the following equation: $H = - [\sum P_i * \ln(P_i)]$, where H = Diversity index, P_i = the number of each species in the sample/total number of samples, and $\ln(P_i)$ = the natural logarithm of this proportion.

Data analysis. A one-way analysis of variance (ANOVA) was used to examine the significance of any differences between mean concentrations of data obtained, followed by the Tukey HSD test, while the relationship between the growth of duckweed and bacterial abundance and diversity was analyzed using regression analysis. Values of $R^2 <$

0.65 were considered to have poor correlations, whereas $R^2 > 0.65$ indicated good correlations. A $p < 0.05$ was considered significant for all analyses.

Results and Discussion. Forty-seven isolates of the putative PSB were screened from duckweed in the biofilter tank. Among the isolates, ten morphologically different bacterial isolates were observed (Table 1). Different harvest regimes caused significant differences ($p < 0.05$) in the bacterial abundance. The harvesting regime significantly impacted the abundance of six types of PSB (PSB1, PSB3, PSB15, PSB17, PSB26, and PSB35). Duckweed harvested every 2 days had the lowest abundance of the putative PSB, while the highest PSB abundance was found in duckweed harvested every 4 days. However, no significant difference was found in PSB abundance between duckweed harvested every 4 days and those harvested every 6 days.

Table 1

The abundance of PSB attached to the duckweed in the biofilter tank

Putative PSB bacteria (cfu x 10 ³ /mL)	Harvest regimes			
	6 days	4 days	2 days	No harvesting
PSB1	5.23±0.21 ^b	5.34±0.26 ^b	4.76±0.34 ^a	4.70±0.32 ^a
PSB3	3.74±0.03 ^b	3.85±0.06 ^b	3.29±0.03 ^a	3.37±0.01 ^a
PSB7	1.79±0.08 ^a	1.75±0.09 ^a	1.64±0.05 ^a	1.59±0.05 ^a
PSB15	4.73±0.12 ^b	4.85±0.19 ^b	4.12±0.13 ^a	4.24±0.15 ^a
PSB17	5.17±0.11 ^b	5.09±0.26 ^b	4.53±0.26 ^a	4.98±0.08 ^b
PSB26	4.60±0.14 ^b	4.52±0.17 ^b	3.85±0.48 ^a	3.97±0.03 ^a
PSB29	1.84±0.21 ^a	1.79±0.27 ^a	1.65±0.35 ^a	1.71±0.46 ^a
PSB30	2.16±0.13 ^a	2.01±0.16 ^a	1.87±0.10 ^a	1.95±0.10 ^a
PSB32	2.89±0.02 ^a	2.94±0.04 ^a	2.75±0.02 ^a	2.80±0.08 ^a
PSB35	3.47±0.08 ^b	3.52±0.17 ^b	3.05±0.07 ^a	3.13±0.05 ^a
Total	35.62±0.43 ^b	35.66±0.44 ^b	31.51±0.37 ^a	32.44±0.39 ^a

Note: values in the same row with the same superscript letter are not significantly different ($p > 0.05$).

Harvesting duckweed every four days resulted in a higher Shannon diversity index value ($H = 2.2910$) than the two-day harvest (2.2365), the six-day harvest (2.2745), and the unharvested (control) ($H = 2.2257$) (Table 2). PSB abundance and duckweed biomass harvest were poorly correlated ($R^2 < 0.65$) with the harvest regime but at the four-day harvest, a significantly stronger correlation between the bacterial diversity and the harvested duckweed biomass was observed. Similarly, the SGR harvest of duckweed was significantly correlated with the abundance and diversity of PSB at the four-day harvest regime ($R^2 > 0.65$; $p < 0.05$), whereas the SGR and biomass harvest of the duckweed harvested every six days had a poor relation to the abundance and diversity of PSB ($R^2 < 0.65$).

Table 2

Correlation between SGR and total biomass harvest of duckweed with abundance and diversity of PSB over the 36-day trial

Parameters	Harvest regimes (days)			
	6	4	2	0
R^2 (biomass vs abundance)	0.5971±0.02 ^a	0.6192±0.06 ^b	0.5941±0.04 ^a	0.6185±0.00 ^b
R^2 (biomass vs diversity)	0.6141±0.01 ^a	0.6502±0.01 ^b	0.6193±0.05 ^a	0.6835±0.03 ^b
R^2 (SGR vs abundance)	0.5852±0.06 ^a	0.6984±0.06 ^b	0.6248±0.03 ^a	0.6894±0.03 ^b
R^2 (SGR vs diversity)	0.6242±0.01 ^a	0.6623±0.01 ^b	0.6193±0.05 ^a	0.6835±0.03 ^b
Shannon diversity index (H)	2.2745±0.01 ^{bc}	2.2910±0.01 ^c	2.2365±0.01 ^b	2.2257±0.00 ^a
GI	3.15±0.02 ^c	3.09±0.03 ^{bc}	3.15±0.04 ^a	3.04±0.03 ^{ab}
Total biomass harvest	674.45±0.54 ^b	1003.45±1.29 ^c	1880.2±0.67 ^d	99.75±0.11 ^a
SGR	8.14±0.00 ^b	33.31±0.00 ^c	11.05±0.00 ^d	3.04±0.00 ^a

Values in the same row with the same superscript letter are not significantly different ($p > 0.05$).

Harvest of duckweed biomass and PSB abundance were poorly correlated ($R^2 < 0.65$) to harvest regime. Significantly stronger correlations between duckweed biomass at harvest and PSB diversity were observed at the 4-day harvest regime. A 4-day harvest regime may provide the desirable duckweed biomass on the water surface and optimal availability of organic substrates as carbon and energy sources for the bacteria (Ishizawa

et al 2017). Likewise, the diversity of PSB was significantly correlated ($R^2 > 0.65$) with SGR of duckweed at the 4-day harvest regime, and the unharvested (control). Furthermore, harvesting duckweed every two and six days resulted in a poor relationship between SGR and biomass harvest with the abundance and diversity of PSB. Different harvest regimes also had a significant influence on the GI of duckweed. Duckweed harvested every two and six days had the highest GI of 3.15, whereas those unharvested control had the lowest GI of 3.04 (Table 2). However, higher GI of the 2 and six-day harvest regime do not have stronger relationship with the abundance and diversity of PSB communities.

Phosphate solubilization efficiency, IAA, and phosphatase activity. The results of the solubilization efficiency of the selected PSB are shown in Table 3. Among the ten bacterial isolates, the highest efficiency was observed for PSB1 (E = 83.33%), followed by PSB17 (E = 75.00%), and then PSB15 (E = 62.50%), whereas PSB29 (E = 33.33%) and PSB32 (E = 34.47) had comparatively lower efficiencies. All putative PSB isolates produced IAA.

Table 3
The phosphate solubilizing efficiency and phosphatase activity recorded for the putative PSB attached to duckweed in the biofilter tank

Putative PSB	Phosphate solubilization efficiency (E, %)	IAA production ($\mu\text{g L}^{-1}$)	Phosphatase activity ($\mu\text{mol g}^{-1}\text{h}^{-1}$)	R^2 (Phosphate solubilization efficiency vs Phosphatase activity)
PSB1	83.33±0.01	7.35±0.01	33.68±0.01	0.921
PSB3	53.59±0.01	7.45±0.02	28.92±0.01	0.921
PSB7	43.93±0.01	7.00±0.04	23.92±0.02	0.921
PSB15	62.50±0.02	7.60±0.01	30.85±0.02	0.921
PSB17	75.00±0.01	7.23±0.01	31.25±0.03	0.921
PSB26	60.83±0.01	7.00±0.02	29.84±0.03	0.921
PSB29	33.33±0.02	7.08±0.02	17.65±0.01	0.921
PSB30	44.82±0.05	6.55±0.03	18.85±0.01	0.921
PSB32	34.47±0.02	7.08±0.02	15.84±0.06	0.921
PSB35	50.00±0.01	6.07±0.01	25.32±0.01	0.921

Data are shown as the mean±SE.

The highest IAA ($7.60 \mu\text{g L}^{-1}$) was produced by PSB15, followed by PSB3 ($7.45 \mu\text{g L}^{-1}$), while the lowest IAA ($6.07 \mu\text{g L}^{-1}$) was produced by PSB35. Furthermore, the maximum phosphatase activity was observed in PSB1 ($33.68 \mu\text{mol g}^{-1} \text{h}^{-1}$), followed by PSB17 ($31.25 \mu\text{mol g}^{-1} \text{h}^{-1}$), while the minimum phosphatase activity was observed in PSB32 ($15.84 \mu\text{mol g}^{-1} \text{h}^{-1}$). The remaining bacterial isolates secreted phosphatase enzyme in range of $17.65\text{--}30.85 \mu\text{mol g}^{-1} \text{h}^{-1}$. The correlation coefficient values ($R^2 = 0.921$) showed a significant relationship ($p < 0.05$) between the phosphate solubilization efficiency of all PSB bacteria and the ability of these bacteria to produce phosphatase enzymes.

Morphology and biochemical characteristics of PSB. Ten bacterial isolates were determined by colony morphology, gram staining, and pigmentation (Table 4). Gram-negative bacteria dominated the bacterial communities. Specifically, 40.41% of gram-negative bacilli isolates were identified as *Pseudomonas* and *Chryseobacterium*; 29.79% of isolates of gram-positive bacilli belonged to *Bacillus*; 23.41% of isolates of gram-negative ovoid, cocci, or coccobacilli were identified as *Azotobacter*, *Acinetobacter*, and *Achromobacter*, and 6.39% of isolates of gram-positive cocci were identified as *Micrococcus* (Table 4). Non-pigmented isolates belonged to *Bacillus*, *Achromobacter*, and *Acinetobacter*, whereas non-motile isolates were characterized as *Chryseobacterium*, *Acinetobacter*, *Azotobacter*, and *Micrococcus*.

Ten species of putative PSB were characterized by both biochemical tests and confirmed with the API 20E (Table 4). All isolates were characterized as the same species by both conventional biochemical tests and API 20E. Out of these ten PSB isolates, as shown in Table 4, *Bacillus* was represented by three species, PSB1, PSB3, and PSB7, *Pseudomonas* was represented by 2 species, PSB17 and PSB26, whereas *Chryseobacterium*, *Micrococcus*, *Azotobacter*, *Acinetobacter*, and *Achromobacter* had only one species each, PSB15, PSB29, PSB30, PSB32, and PSB35 respectively.

Table 4

The biochemical characteristics of PSB isolated from the biofilter tanks

Test	The PSB isolate										
	PSB1	PSB3	PSB7	PSB15	PSB17	PSB26	PSB29	PSB30	PSB32	PSB35	
Colony morphology	Bacilli	Bacilli	Bacilli	Bacilli	Bacilli	Bacilli	Cocci	Ovoid	Cocci	Cocci	
No. clones (%)	19.15	6.39	4.25	12.76	14.89	12.76	6.39	4.45	6.39	10.54	
Pigmentation	Yellowish-green	Translucent	White	Yellow	Yellowish-green	Brownish-yellow	Yellowish-orange	Yellow	White	White	
Spore	+	-	+	-	-	-	-	-	-	-	
Motility	-	+	+	-	+	+	-	+	-	-	
Gram stain	+	+	+	-	-	-	+	-	-	+	
Catalase	+	+	+	+	+	+	+	+	+	+	
Oxidase	+	-	-	+	-	+	+	-	-	+	
H ₂ S production	+	-	-	+	+	-	-	+	-	-	
Gelatinase	-	-	+	+	-	+	+	+	-	+	
Nitrate reduction	+	+	+	+	-	-	-	+	-	-	
Starch hydrolysis	+	+	+	+	+	+	-	+	+	-	
Citrate utilization	+	-	+	+	+	+	+	+	+	+	
Ornithine	-	-	-	-	-	-	-	+	-	-	
Indole	-	-	-	+	+	-	-	+	-	-	
Tyrosine hydrolysis	+	-	-	-	+	-	-	-	+	+	
Arginine dihydrolase	-	+	-	-	+	+	-	+	-	+	
Methyl Red	-	-	+	-	+	+	-	-	-	+	
Voges-Proskauer	-	-	+	+	+	+	-	-	+	-	
Urease	+	-	-	-	+	+	-	+	+	-	
Glucose	+	+	+	+	+	+	+	+	+	-	
Fructose	+	+	+	+	+	+	+	+	+	+	
Lactose	-	-	-	-	+	+	-	-	-	-	
Maltose	+	+	+	+	+	-	-	+	-	-	
Mannose	+	+	+	+	+	+	-	+	-	-	
Sucrose	+	+	+	+	+	+	-	+	-	-	
Xylose	-	-	+	-	-	+	+	+	-	+	

Where: PSB1 = *Bacillus cereus*; PSB3 = *Bacillus licheniformis*; PSB7 = *Bacillus subtilis*; PSB15 = *Chryseobacterium indologenes*; PSB17 = *Pseudomonas fluorescens*; PSB26 = *Pseudomonas mendocina*; PSB29 = *Micrococcus luteus*; PSB30 = *Azotobacter vinelandii*; PSB32 = *Acinetobacter calcoaceticus*; PSB35 = *Achromobacter xylosoxidans*.

Abundance and diversity of PSB. Different harvest regimes significantly affected the abundance of PSB (Table 1), in which six of ten PSB communities, *Bacillus cereus*, *Bacillus licheniformis*, *Chrysobacterium indologenes*, *Pseudomonas fluorescens*, *Pseudomonas mendocina* and *Achromobacter xylosoxidans* were significantly affected ($p < 0.05$) by different harvest regimes. Duckweed harvested with the shortest interval of 2 days had lower abundance than those with longer harvest intervals of 4 and 6 days. Similar results had been previously observed in the abundance of the heterotrophic bacteria associated with the different harvesting regimes in barramundi (*Lates calcarifer*) RAS (Ardiansyah & Fotedar 2016). Therefore, the four-day harvest regime may provide the desirable duckweed biomass and optimal availability of organic substrates as carbon and energy sources for the PSB.

Only ten isolates from forty-seven showed the formation of halo zones around the growing colonies on PVK media plates. Halo zones are formed because of the transformation of glucose into organic acids, in which glucose is the carbon source used in the test of PSB isolated from the biofilter to dissolve the bonded phosphor marked by halo zones (Pande et al 2017). However, the production of organic acids was not determined in the present study; thus, the authors cannot confirm whether all PSB isolates are capable of producing organic acids.

There is increasing evidence that PSB improves plant growth due to the biosynthesis of plant growth substances rather than to their action in releasing available P. In the present study, all PSB isolated from duckweed were capable of producing IAA. Majda & Robert (2018) stated that IAA is the biologically active form of auxin that stimulates radical system growth and increases the uptake of nutrients by plants. Thus, it has prominent effects on plant growth and development (Egamberdieva et al 2017). The results indicated that all PSB isolates can secrete physiologically active auxins. Shahab et al (2009) reported that PSB isolated from mung bean (*Vigna radiata*) excreted phytohormones, including auxins. Other researchers found similar results; that isolated bacterial species from rice fields have the potential to produce IAA (Etesami et al 2015). These results were also similar to the findings of Bal et al (2013); where the PSB isolated from the rhizosphere of Chinese cabbage (*Brassica rapa*) were found to solubilize P in the media, and besides this, they were able to produce IAA. However, the capacity to synthesize IAA is widespread among the PSB. Production of IAA greatly varies among different species and is influenced by culture conditions, growth stage, and substrate availability (Mohite 2013).

Phosphatase enzyme activity. Phosphatase enzyme activity showed that all the selected PSB isolates solubilized inorganic phosphates into various degrees in the PVK culture medium by producing phosphatase enzymes. The PSB have been shown to increase the solubility of the P-insoluble compound through the secretion of extracellular enzymes such as organic acid, phytase, and phosphatase enzymes (Khan et al 2014). These enzymes are found in all organisms but only bacteria, fungi, and some algae can secrete them outside their cells. However, the PSB are more effective in P solubilization than fungi (Nacoon et al 2020). Among the whole microbial population in the aquatic environment, PSB constituted 1-50%, while phosphorus-solubilizing fungi (PSF) constituted only 0.1-0.5% in P solubilization potential (Ingle & Padole 2017).

Furthermore, during the conversion process, P is partially assimilated by PSB but the amount made soluble and released is more than the requirement of the PSB. The excess amount released is made available to plants. During this conversion process, extracellular enzymes convert calcium phosphate to di- or monobasic phosphates and then become biologically available to plants (Kalayu 2019).

The stronger correlation between phosphate solubilizing efficiency and phosphatase enzyme activity may indicate an increase in the availability of P in the medium, which is assumed to be facilitated through increased phosphatase enzyme activity (Singh & Prakash 2012; Behera et al 2017). The results suggest that all selected PSB can produce phosphatase enzymes to convert insoluble phosphate into the soluble form, which leads to increased available phosphate for plants (Kalayu 2019). Microorganisms increase P availability to the macrophytes by mineralizing organic P and

by solubilizing precipitated phosphates (Alori et al 2017). Additionally, Behera et al (2017) found a positive correlation between phosphate solubilizing activity and phosphatase enzyme activity. These previous findings are in agreement with the present results, which showed that the increase in inorganic phosphate levels was due to phosphate solubilization, and did not repress phosphatase production by the PSB isolates found in the study.

Morphology and biochemical characteristics of PSB. The results found in this study are in line with those reported by Maitra et al (2015) that *Pseudomonas*, *Chryseobacterium*, *Aeromonas*, and *Acinetobacter* are common inhabitants of various aquatic environments. Furthermore, Maitra et al (2015) and Adeleke & Babalola (2021) reported that the bacterial genera with the capacity to solubilize insoluble inorganic phosphate compounds amongst aquatic environments are *Bacillus*, *Enterobacter*, *Agrobacterium*, *Pseudomonas*, *Acinetobacter*, *Achromobacter*, *Micrococcus*, *Chryseobacterium*, and *Aerobacter*.

The most common bacterial genera in this study were *Bacillus*. The majority of the *Bacillus* species showed high phosphatase activity but the proportion varied with species. The *Bacillus* consisted of obligate aerobes (*B. subtilis*) or facultative anaerobes (*B. cereus* and *B. licheniformis*). The genus is known to have more characteristics promoting plant growth: synthesis of phytohormones such as IAA (auxins); nitrogen fixation; antifungal activity by siderophore production; antibiotics; enzymes; and phosphorus solubilization (Nabti et al 2013).

Sharma et al (2013) reported that the genus *Pseudomonas* was a suitable genus of bacteria in dissolving bonded P from any source, and was dominant in the mineralization of organic phosphorus. This type of bacteria can also be found in extreme saline and alkaline water habitats (Poli et al 2017). In this study, *Pseudomonas* consisted of *P. fluorescens* and *P. mendocina*. Bacteria from the genera *Bacillus* and *Pseudomonas* are among the most powerful phosphate solubilizers (Kalayu 2019).

A considerable number of bacteria from different genera are capable of solubilizing phosphate including *Chryseobacterium*, which has been isolated from a wide range of habitats such as soil, plant roots, sludge, fish, sewage, fresh water, freshwater sediment, marine water, and marine sediment. The genus *Chryseobacterium* was first created for five species formerly classified as members of the genus *Flavobacterium* (Song et al 2021). Various *Chryseobacterium* species have been shown to solubilize phosphate by releasing organic acids that mobilize P, and by releasing phosphatase to release phosphate groups bound to organic matter (Kalayu 2019).

P availability depends on the degree of solubilization by various organic and inorganic acids produced by microorganisms, and the genus *Micrococcus* is one of the most important microorganisms that produce a substantial number of phosphates. *Micrococcus* inhabits a wide range of environments including soil, water, and dust, and can sustain well environments with high salt concentration and little water. This genus has been shown to possess multiple plant growth traits like P-solubilisation, and IAA, and siderophore production (Afzal et al 2019).

Various bacterial genera are involved in various biotic activities important to nutrient turnover dynamics (Ahemad & Kibret 2014). Thus, there is ongoing research to discover various rhizobacteria that possess novel traits such as phosphate solubilization (Ahemad & Khan 2012) capacities. Among the bacterial genera with this capability is *A. xylooxidans* (Ahemad & Kibret 2014). This species shows a considerable level of nitrogenase activity, IAA production, and P solubilization ability. In addition, this species may also increase copper uptake by plants and increase their shoot length, root length, and fresh weight, and dry weight of plants (Yang et al 2013).

Azotobacter is a genus of gamma-proteobacterium belonging to the family Pseudomonadaceae. It is an obligately aerobic, free-living Gram-negative bacterium that is broadly dispersed in various environments including water, soil, and sediments (Dangi 2015). *Azotobacter* has favorable effects on plant yields due to its capability in fixing nitrogen and solubilizing phosphates (Nosrati et al 2014). *A. vinelandii* produces metabolically dormant cysts, which are formed under unfavorable environmental

conditions; thus, it is suitable for use in diverse environments (Garcia³⁴ al 2014). Several *Azotobacter* species isolated from wheat rhizosphere has shown the ability to solubilize tricalcium phosphate, Mussoorie rock phosphate, and also to produce IAA. Additionally, the use of *Azotobacter* may increase seed yield, plant height, and microbial population in soil (Peng et al 2013).

A. calcoaceticus has demonstrated the potential²⁴ ability of P solubilization through the production of organic acids, and can provide plant growth-promoting factors such as the production of IAA and siderophores. This species also exhibits resistance to lead (Pb) and antibiotic³². The use of *A. calcoaceticus* effectively increases the available Pb in the rhizosphere soil and promotes the growth of the host plant, which leads to an increase in Pb uptake (Ren et al 2013).

All the putative PSB from the present study have been known to play a fundamental role in the phosphate solubilization process in a diverse range of ecosystems (Anand et al 2016). Our research suggests that duckweed biofilters in IRAS are suitable substrates for the attachment, survival, and growth of PSB.

A short two-day harvest regime reduced the PSB abundance in biofilter tanks, whereas a six-day harvest regime, or no harvest, decreased duckweed biomass due to chlorosis and disconnection of their fronds. The results showed that a four-day harvest regime maintains the optimum biomass of duckweed. Furthermore, the different harvesting regimes influenced the abundance and diversity of PSB in IRAS. The selected PSB isolates should be further evaluated in larger scale *in vivo* trials regarding the mechanism and efficiency of P solubilization, and their potential application in managing sustainable aquaculture systems.

Conclusions. Duckweed in the biofilter of RAS is a suitable substrate for the attachment, survival, and growth of phosphate solubilizing bacteria. A short 2-day harvesting regime reduced the abundance of phosphate solubilizing bacteria in the biofilter tank, whereas the 6-day or no-harvesting regime decreased duckweed biomass. The results suggested that the 4-day harvesting regime maintained optimum duckweed biomass in the biofilter tank. Harvesting duckweed every four days provides desirable conditions for the attachment and growth of bacteria, thus increasing the abundance and diversity of PSB. These findings suggest the efficiency of removing excessive phosphorus and the associated PSB by duckweed and increase the productivity of IRAS.

Acknowledgements. This study was funded by the Indonesian Directorate General of Higher Education (DIKTI, Grant No. 1287/E4.4/K).

References

- Adeleke B. S., Babalola O. O., 2021 Roles of plant endosphere microbes in agriculture - a review. *Journal of Plant Growth Regulation* 2021. Doi.org/10.1007/s00344-021-10406-2.
- Aditi F. Y., Rahman S. S., Hossain M., 2017 A study on the microbiological status of mineral drinking water. *Open Microbiology Journal* 11:31-44.
- Afzal I., Shinwari Z. K., Sikandar S., Shahzad S., 2019 Plant beneficial endophytic bacteria: mechanisms, diversity, host range and genetic determinants. *Microbiological Research* 221:36-49.
- Ahemad M., Khan M. S., 2012 Effect of fungicides on plant growth promoting activities of phosphate solubilizing *Pseudomonas putida* isolated from mustard (*Brassica campestris*) rhizosphere. *Chemosphere* 86(9):945-950.
- Ahemad M., Kibret M., 2014 Mechanisms and applications of plant growth promoting rhizobacteria: current perspective. *Journal of King Saud University - Science* 26(1):1-20.
- Alori E. T., Glick B. R., Babalola O. O., 2017 Microbial phosphorus solubilization and its potential for use in sustainable agriculture. *Frontiers in Microbiology* 8:971.

- Anand K., Kumar B., Mallick M. A., 2016 Phosphate solubilizing microbes: an effective and alternative approach as biofertilizers. *International Journal of Pharmacy and Pharmaceutical Sciences* 8(2):37-40.
- Ardiansyah A., Fotedar R., 2016 The abundance and diversity of heterotrophic bacteria as a function of harvesting frequency of duckweed (*Lemna minor* L.) in recirculating aquaculture systems. *Letters in Applied Microbiology* 63(1):53-59.
- Bal H., Das S., Dangar T., Adhya T., 2013 ACC deaminase and IAA producing growth promoting bacteria from the rhizosphere soil of tropical rice plants. *Journal of Basic Microbiology* 53(12):972-984.
- Behera B. C., Yadav H., Singh S. K., Mishra R. R., Sethi B. K., Dutta S. K., Thatoi H. N., 2017 Phosphate solubilization and acid phosphatase activity of *Serratia* sp. isolated from mangrove soil of Mahanadi river delta, Odisha, India. *Journal of Genetic Engineering and Biotechnology* 15(1):169-178.
- Blanco-Vargas A., Rodríguez-Gacha L. M., Sánchez-Castro N., Garzón-Jaramillo R., Pedroza-Camacho L. D., Poutou-Piñales R. A., Rivera-Hoyos C. M., Díaz-Ariza L. A., Pedroza-Rodríguez A. M., 2020 Phosphate-solubilizing *Pseudomonas* sp., and *Serratia* sp., co-culture for *Allium cepa* L. growth promotion. *Heliyon* 6(10):e05218.
- Chaiham M., Lumyong S., 2011 Screening and optimization of indole-3-acetic acid production and phosphate solubilization from rhizobacteria aimed at improving plant growth. *Current Microbiology* 62(1):173-181.
- Chen W. M., Tang Y. Q., Mori K., Wu X. L., 2012 Distribution of culturable endophytic bacteria in aquatic plants and their potential for bioremediation in polluted waters. *Aquatic Biology* 15:99-110.
- Cheng Y., Feng G., Moraru C. I., 2019 Micro- and nanotopography sensitive bacterial attachment mechanisms: a review. *Frontiers in Microbiology* 10:191.
- Dangi S. R., Gerik J. S., Tirado-Corbala R., Ajwa H., 2015 Soil microbial community structure and target organisms under different fumigation treatments. *Applied and Environmental Soil Science* 2015:673264.
- Daud M. K., Ali S., Abbas Z., Zaheer I. E., Riaz M. A., Malik A., Hussain A., Rizwan M., Zia-ur-Rehman M., Zhu S. J., 2018 Potential of duckweed (*Lemna minor*) for the phytoremediation of landfill leachate. *Journal of Chemistry* 2018:3951540.
- Dauda A. B., Ajadi A., Tola-Fabunmi A. S., Akinwale A. O., 2019 Waste production in aquaculture: sources, components and managements in different culture systems. *Aquaculture and Fisheries* 4(3):81-88.
- Egamberdieva D., Wirth S. J., Alqarawi A. A., Abd-Allah E. F., Hashem A., 2017 Phytohormones and beneficial microbes: essential components for plants to balance stress and fitness. *Frontiers in Microbiology* 8:2104.
- Elias F., Woyessa D., Muleta D., 2016 Phosphate solubilization potential of rhizosphere fungi isolated from plants in Jimma zone, Southwest Ethiopia. *International Journal of Microbiology* 2016:5472601.
- Etesami H., Alikhani H. A., Hosseini H. M., 2015 Indole-3-acetic acid (IAA) production trait, a useful screening to select endophytic and rhizosphere competent bacteria for rice growth promoting agents. *MethodsX* 2:72-78.
- García A., Segura D., Espín G., Galindo E., Castillo T., Peña C., 2014 High production of poly- β -hydroxybutyrate (PHB) by an *Azotobacter vinelandii* mutant altered in PHB regulation using a fed-batch fermentation process. *Biochemical Engineering Journal* 82:117-123.
- Garrity G. M., Brenner D. J., Krieg N. R., 2005 *Bergey's manual of systematic bacteriology*. Vol II: The Proteobacteria. 2nd edition. Springer, Berlin, 1106 pp.
- Gomes M. P., Gonçalves C. A., Moreira de Brito J. C., Souza A. M., da Silva Cruz F. V., Bicalho E. M., Figueredo C. C., Garcia Q. Z., 2017 Ciprofloxacin induces oxidative stress in duckweed (*Lemna minor* L.): implications for energy metabolism and antibiotic-uptake ability. *Journal of Hazardous Materials* 328:140-149.
- Herath S., Satoh S., 2015 Environmental impact of phosphorus and nitrogen from aquaculture. *Feed and Feeding Practices in Aquaculture*, Woodhead Publishing Series pp.369-386.

- Ingle K., Padole D., 2017 Phosphate solubilizing microbes: an overview. *International Journal of Current Microbiology and Applied Sciences* 6(1):844-852.
- Ishizawa H., Kuroda M., Morikawa M., Ike M., 2017 Evaluation of environmental bacterial communities as a factor affecting the growth of duckweed *Lemna minor*. *Biotechnology for Biofuels* 10:62.
- Jorgensen S. E., Tundisi J. G., Tundisi T. M., 2012 Handbook of inland aquatic ecosystem management. 1st edition. CRC Press, USA, 452 pp.
- Kalayu G., 2019 Phosphate solubilizing microorganisms: promising approach as biofertilizers. *International Journal of Agronomy* 2019:4917256.
- Kawasaki N., Kushairi M. R. M., Nagao N., Yusoff F., Imai A., Kohzu A., 2016 Release of nitrogen and phosphorus from aquaculture farms to Selangor River, Malaysia. *International Journal of Environmental Science and Development* 7(2):113-116.
- Khan M., Zaidi S. A., Ahmad E., 2014 Mechanism of phosphate solubilization and physiological functions of phosphate-solubilizing microorganisms. In: Phosphate solubilizing microorganisms. Khan M. S., Zaidi A., Musarrat J. (eds), Springer Publishers Switzerland, pp. 31-62.
- Maitra N., Manna S. K., Samanta S., Sarkar K., Debnath D., Bandopadhyay C., Sahu S. K., Sharma A. P., 2015 Ecological significance and phosphorus release potential of phosphate solubilizing bacteria in freshwater ecosystems. *Hydrobiologia* 745:69-83.
- Majda M., Robert S., 2018 The role of auxin in cell wall expansion. *International Journal of Molecular Sciences* 19(4):951.
- Martins C. I. M., Eding E. H., Verdegem M. C. J., Heinsbroek L. T. N., Schneider O., Blancheton J. P., Roque d'Orbcastel E., Verreth J. A. J., 2010 New developments in recirculating aquaculture systems in Europe: a perspective on environmental sustainability. *Aquaculture Engineering* 43(3):83-93.
- Mohite B., 2013 Isolation and characterization of indole acetic acid (IAA) producing bacteria from rhizospheric soil and its effect on plant growth. *Journal of Soil Science and Plant Nutrition* 13(3):638-649.
- Muszyńska E., Labudda M., 2019 Dual role of metallic trace elements in stress biology - from negative to beneficial impact on plants. *International Journal of Molecular Sciences* 20(13):3117.
- Nabti E. H., Mokrane N., Ghouli M., Manyani H., Dary M., Megias M. G., 2013 Isolation and characterization of two halophilic *Bacillus* (*B. licheniformis* and *Bacillus* sp) with antifungal activity. *Journal of Ecology of Health and Environment* 1(1):13-17.
- Nacoon S., Jogloy S., Riddech N., Mongkoltharuk W., Kuyper T. W., Boonlue S., 2020 Interaction between phosphate solubilizing bacteria and arbuscular mycorrhizal fungi on growth promotion and tuber inulin content of *Helianthus tuberosus* L. *Scientific Reports* 10:4916.
- Nassar H., Shaban M., Basseem S., Abdel-Gawad F., 2015 Utilization of duckweed (DW) in nutrient removal from agricultural waste water and producing alternative economic animal fodder. *Der Pharma Chemica* 7(12):280-285.
- Nosrati R., Owlia P., Saderi H., Rasooli I., Malboobi M. A., 2014 Phosphate solubilization characteristics of efficient nitrogen fixing soil *Azotobacter* strains. *Iranian Journal of Microbiology* 6(4):285-295.
- Omri I., Aouidi F., Bouallagui H., Godon J. J., Hamdi M., 2013 Performance study of biofilter developed to treat H₂S from wastewater odour. *Saudi Journal of Biological Sciences* 20(2):169-176.
- Ozengin N., Elmaci A., 2007 Performance of duckweed (*Lemna minor* L.) on different types of wastewater treatment. *Journal of Environmental Biology* 28(2):307-314.
- Pande A., Pandey P., Mehra S., Singh M., Kaushik S., 2017 Phenotypic and genotypic characterization of phosphate solubilizing bacteria and their efficiency on the growth of maize. *Journal of Genetic Engineering and Biotechnology* 15(2):379-391.
- Paolacci S., Harrison S., Jansen M., 2018 The invasive duckweed *Lemna minuta* Kunth displays a different light utilisation strategy than native *Lemna minor* Linnaeus. *Aquatic Botany* 146:8-14.

- Peng S. H., Wan-Azha W. M., Wong W. Z., Go W. Z., Chai E. W., Chin K. L., H'ng P. S., 2013 Effect of using agro-fertilizers and N-fixing *Azotobacter* enhanced biofertilizers on the growth and yield of corn. *Journal of Applied Sciences* 13(3):508-512.
- Poli A., Finore I., Romano I., Gioiello A., Lama L., Nicolaus B., 2017 Microbial diversity in extreme marine habitats and their biomolecules. *Microorganisms* 5(2):25.
- Ren Y. X., Zhu X. L., Fan D. D., Ma P., Liang L. H., 2013 Inoculation of phosphate solubilizing bacteria for the improvement of lead accumulation by *Brassica juncea*. *Environmental Technology* 34:463-469.
- Ruzhitskaya O., Gogina E., 2017 Methods for removing of phosphates from wastewater. *MATEC Web of Conferences* 106:07006.
- Schoebitz M., Ceballos C., Ciampi L., 2013 Effect of immobilized phosphate solubilizing bacteria on wheat growth and phosphate uptake. *Journal of Soil Science and Plant Nutrition* 13(1):1-10.
- Shahab S., Ahmed N., Khan N., 2009 Indole acetic acid production and enhanced plant growth promotion by indigenous PSBs. *African Journal of Agricultural Research* 4(11):1312-1316.
- Shardendu S., Sayantan D., Sharma D., Irfan S., 2012 Luxury uptake and removal of phosphorus from water column by representative aquatic plants and its implication for wetland management. *International Scholarly Research Notices* 2012:516947.
- Sharma S. B., Sayyed R. Z., Trivedi M. H., Gobi T. A., 2013 Phosphate solubilizing microbes: sustainable approach for managing phosphorus deficiency in agricultural soils. *SpringerPlus* 2:587.
- Singh M., Prakash N. T., 2012 Characterisation of phosphate solubilising bacteria in sandy loam soil under chickpea cropping system. *Indian Journal of Microbiology* 52(2):167-173.
- Song J., Min L. J., Wu J. R., He Q. F., Chen F. M., Wang Y., 2021 Response of the microbial community to phosphate-solubilizing bacterial inoculants on *Ulmus chenmoui* Cheng in Eastern China. *PLoS ONE* 16(2):e0247309.
- Sun G., Zhang C., Li W., Yuan L., He S., Wang L., 2019 Effect of chemical dose on phosphorus removal and membrane fouling control in a UCT-MBR. *Frontiers of Environmental Science and Engineering* 13:1.
- Sushanta B., Sunita P., Gonsalves M. J. B. D., Das A., Mascarenhas-Pereira M. B. L., Loka Barathi P. A., 2012 Phosphate solubilizing bacteria: comparison between coastal and deep-sea sediments. *Journal of Coastal Environment* 3(2):153-164.
- Turcois A. E., Papenbrock J., 2014 Sustainable treatment of aquaculture effluents - what can we learn from the past for the future? *Sustainability* 6(2):836-856.
- Velichkova K., Sirakov I., 2013 The usage of aquatic floating macrophytes (*Lemna* and *Wolffia*) as biofilter in recirculation aquaculture system (RAS). *Turkish Journal of Fisheries and Aquatic Sciences* 13:101-110.
- Verdegem M. C. J., 2013 Nutrient discharge from aquaculture operations in function of system design and production environment. *Reviews in Aquaculture* 5(3):158-171.
- Yang R., Luo C., Chen Y., Wang G., Xu Y., Shen Z., 2013 Copper-resistant bacteria enhance plant growth and copper phytoextraction. *International Journal of Phytoremediation* 15(6):573-584.
- Zhang J., Wang P., Fang L., Zhang Q. A., Yan C., Chen J., 2017 Isolation and characterization of phosphate-solubilizing bacteria from mushroom residues and their effect on tomato plant growth promotion. *Polish Journal of Microbiology* 66(1):57-65.
- Zhao Y., Zhang C., 2017 Purification of leach solution of zinc and lead in alkaline solutions. In: *Pollution control and resource reuse for alkaline hydrometallurgy of amphoteric metal hazardous wastes. Handbook of Environmental Engineering*, vol 18. Springer, Cham, pp. 133-170.

Received: 31 August 2021. Accepted: 12 October 2021. Published online: 29 October 2021.

Authors:

Dahlia, Department of Aquaculture, Pangkep State Polytechnic of Agriculture, Makassar–Pare Rd, Mandalle, Pangkep Regency 90655, South Sulawesi, Indonesia, e-mail: unga_dahlia@yahoo.co.id

Ardiansyah, Department of Aquaculture, Pangkep State Polytechnic of Agriculture, Makassar–Pare Rd, Mandalle, Pangkep Regency 90655, South Sulawesi, Indonesia, e-mail: ardi_kimsan@yahoo.com

Indrayani, Study Program Agricultural Technology, Faculty of Engineering, Makassar State University, South Sulawesi, Indonesia, e-mail: indrayani_tajudin@yahoo.com.au

Firmansyah Bin Abd Jabbar, Aquaculture Study Program, Universitas Sulawesi Barat, Majene, West Sulawesi, Indonesia, e-mail: firmansyahjabbar@unsulbar.ac.id

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Dahlia, Ardiansyah, Indrayani, Jabbar F. B. A., 2021 The abundance and diversity of phosphate-solubilizing bacteria in integrated recirculating aquaculture systems as a function of harvesting regime of duckweed (*Lemna minor* L.). *AAFL Bioflux* 14(5):3055-3067.

The abundance and diversity of phosphatesolubilizing bacteria in integrated recirculating aquaculture systems

ORIGINALITY REPORT

9%

SIMILARITY INDEX

5%

INTERNET SOURCES

8%

PUBLICATIONS

1%

STUDENT PAPERS

PRIMARY SOURCES

- 1** F Kamiludin, H Setiadi, W Sumadio. "Livelihoods sustainability pattern in Sangrawayang Village, Simpenan District, Sukabumi Regency", IOP Conference Series: Earth and Environmental Science, 2020
Publication 1%
 - 2** www.hindawi.com
Internet Source <1%
 - 3** Ahemad, Munees. "Phosphate-solubilizing bacteria-assisted phytoremediation of metalliferous soils: a review", 3 Biotech, 2014.
Publication <1%
 - 4** Carlos R. Fernandez Pulido, Jonathan Caballero, Mary Ann Bruns, Rachel A. Brennan. "Recovery of waste nutrients by duckweed for reuse in sustainable agriculture: Second-year results of a field pilot study with sorghum", Ecological Engineering, 2021
Publication <1%
-

5

Kumar Adesh, Kumar Shukla Umesh, Singh Anshu, Kumar Poonam Arun, Prasad Shambhoo, Kumar Singh Sushil, Kumar Dharmendra. "Evaluation of Pseudomonas isolates from wheat for some important plant growth promoting traits", African Journal of Microbiology Research, 2014

Publication

<1 %

6

Submitted to University of KwaZulu-Natal

Student Paper

<1 %

7

vtechworks.lib.vt.edu

Internet Source

<1 %

8

Submitted to Midlands State University

Student Paper

<1 %

9

ir.lib.ruh.ac.lk

Internet Source

<1 %

10

Narendra Babu, Anupama, Sudisha Jogaiah, Shin-ichi Ito, Amruthesh Kestur Nagaraj, and Lam-Son Phan Tran. "Improvement of growth, fruit weight and early blight disease protection of tomato plants by rhizosphere bacteria is correlated with their beneficial traits and induced biosynthesis of antioxidant peroxidase and polyphenol oxidase", Plant Science, 2015.

Publication

<1 %

11

Nilanjan Maitra, Sanjib Kumar Manna, Srikanta Samanta, Keka Sarkar et al. "Ecological significance and phosphorus release potential of phosphate solubilizing bacteria in freshwater ecosystems", *Hydrobiologia*, 2014

Publication

<1 %

12

Selvi Panggua, Basri Wello, Baso' Jabu, Doris Macdonald. "Self-Assessed Professional Competence of the High School EFL Teachers in Toraja Indonesia", *International Journal of English Linguistics*, 2017

Publication

<1 %

13

Syeda Huma Bokhari, Iftikhar Ahmad, Muhammad Mahmood-Ul-Hassan, Ashiq Mohammad. " Phytoremediation potential of L. for heavy metals ", *International Journal of Phytoremediation*, 2015

Publication

<1 %

14

VallverdÀ-Queralt, Anna, Gemma Oms-Oliu, Isabel Odriozola-Serrano, Rosa MÀ^a Lamuela-RaventÀ³s, Olga MartÀn-Belloso, and Pedro Elez-MartÀnez. "Metabolite profiling of phenolic and carotenoid contents in tomatoes after moderate-intensity pulsed electric field treatments", *Food Chemistry*, 2013.

Publication

<1 %

15

Internet Source

<1 %

16

journals.plos.org

Internet Source

<1 %

17

kuojs.lib.ku.ac.th

Internet Source

<1 %

18

www.peertechz.com

Internet Source

<1 %

19

Abdur Rahim Khan, Ihsan Ullah, Muhammad Waqas, Raheem Shahzad et al. "Plant growth-promoting potential of endophytic fungi isolated from Solanum nigrum leaves", World Journal of Microbiology and Biotechnology, 2015

Publication

<1 %

20

Aleksandra Wdowczyk, Agata Szymańska-Pulikowska, Magdalena Domańska. "Analysis of the Bacterial Biocenosis of Activated Sludge Treated with Leachate from Municipal Landfills", International Journal of Environmental Research and Public Health, 2022

Publication

<1 %

21

Oluwaseyi A. Akpor, Victoria O. Aina, Oghenerobor B. Akpor. "Occupational Stress Levels and Coping Strategies among Nurses

<1 %

Working in an Urban Metropolis in North
Central Nigeria", The Open Pain Journal, 2023

Publication

22

www.jove.com

Internet Source

<1 %

23

www.tlsr.usm.my

Internet Source

<1 %

24

Jay Prakash Verma, Durgesh Kumar Jaiswal, Punkaj Kumar Maurya. "Screening of bacterial strains for developing effective pesticide-tolerant plant growth-promoting microbial consortia from rhizosphere soils of vegetable fields of eastern Uttar Pradesh, India", Energy, Ecology and Environment, 2016

Publication

<1 %

25

Lingling Zhang, Yuping Zhang, Lizao Liu. "Effect of submerged macrophytes *Vallisneria spiralis* L. on restoring the sediment contaminated by enrofloxacin in aquaculture ponds", Ecological Engineering, 2019

Publication

<1 %

26

Setu Bazie Tagele, Sang Woo Kim, Hyun Gu Lee, Hyun Seung Kim, Youn Su Lee. "Effectiveness of multi-trait *Burkholderia* contaminans KNU17BI1 in growth promotion and management of banded leaf and sheath blight in maize seedling", Microbiological Research, 2018

<1 %

27	api.intechopen.com Internet Source	<1 %
28	bnrc.springeropen.com Internet Source	<1 %
29	dspace.unitus.it Internet Source	<1 %
30	etda.libraries.psu.edu Internet Source	<1 %
31	www.trjfas.org Internet Source	<1 %
32	Asfa Rizvi, Bilal Ahmed, Mohammad Saghir Khan, Shahid Umar, Jintae Lee. "Sorghum-Phosphate Solubilizers Interactions: Crop Nutrition, Biotic Stress Alleviation, and Yield Optimization", <i>Frontiers in Plant Science</i> , 2021 Publication	<1 %
33	Submitted to Cleveland State University Student Paper	<1 %
34	D P Krishna Samal, Lala Behari Sukla. "Isolation of phosphate solubilizing fungus from rice fields of Odisha for agro - ecological sustainable agriculture", <i>Environmental Quality Management</i> , 2022 Publication	<1 %

35

Daria Chlebek, Artur Pinski, Joanna Żur, Justyna Michalska, Katarzyna Hupert-Kocurek. "Genome Mining and Evaluation of the Biocontrol Potential of *Pseudomonas fluorescens* BRZ63, a New Endophyte of Oilseed Rape (*Brassica napus* L.) against Fungal Pathogens", *International Journal of Molecular Sciences*, 2020

Publication

<1 %

36

Elena Bartkiene, Vadims Bartkevics, Janis Rusko, Vytaute Starkute, Egle Bendoraitiene, Daiva Zadeike, Grazina Juodeikiene. "The effect of *Pediococcus acidilactici* and *Lactobacillus sakei* on biogenic amines formation and free amino acid profile in different lupin during fermentation", *LWT*, 2016

Publication

<1 %

37

H. M. Abdel-Rahman, A. A. Salem, Mahmoud M. A. Moustafa, Hoda A. S. El-Garhy. "A novice *Achromobacter* sp. EMCC1936 strain acts as a plant-growth-promoting agent", *Acta Physiologiae Plantarum*, 2017

Publication

<1 %

38

Madhurankhi Goswami, Chandana Malakar, Suresh Deka. "Chapter 2 Rhizosphere Microbes for Sustainable Maintenance of

<1 %

39

Mamta, Praveen Rahi, Vijaylata Pathania, Arvind Gulati, Bikram Singh, Ravinder Kumar Bhanwra, Rupinder Tewari. "Stimulatory effect of phosphate-solubilizing bacteria on plant growth, stevioside and rebaudioside-A contents of *Stevia rebaudiana* Bertoni", *Applied Soil Ecology*, 2010

Publication

<1 %

40

María Rebolleda - Gómez, Nicole J. Forrester, Avery L. Russell, Na Wei et al. "Gazing into the anthosphere: considering how microbes influence floral evolution", *New Phytologist*, 2019

Publication

<1 %

41

Małgorzata Pawlik, Zofia Piotrowska-Seget. "Endophytic Bacteria Associated with : Their Potential for Hydrocarbon-Utilizing and Plant Growth-Promotion ", *Journal of Toxicology and Environmental Health, Part A*, 2015

Publication

<1 %

42

ojs.uma.ac.id
Internet Source

<1 %

43

scialert.net
Internet Source

<1 %

44

www.pesca.agricultura.sp.gov.br

Internet Source

<1 %

45

Éamonn Walsh, Holger Kuehnhold, Seán O'Brien, Neil E. Coughlan, Marcel A.K. Jansen. "Light intensity alters the phytoremediation potential of *Lemna minor*", *Environmental Science and Pollution Research*, 2021

Publication

<1 %

46

Adnane Bargaz, Karim Lyamlouli, Mohamed Chtouki, Youssef Zeroual, Driss Dhiba. "Soil Microbial Resources for Improving Fertilizers Efficiency in an Integrated Plant Nutrient Management System", *Frontiers in Microbiology*, 2018

Publication

<1 %

47

"Indole-3-Acetic Acid and 1-Aminocyclopropane-1-Carboxylate Deaminase: Bacterial Traits Required in Rhizosphere, Rhizoplane and/or Endophytic Competence by Beneficial Bacteria", *Sustainable Development and Biodiversity*, 2015.

Publication

<1 %

48

Bhawna Dipta, Sonal Bhardwaj, Manoj Kaushal, Shruti Kirti, Ruchi Sharma. "Obliteration of phosphorus deficiency in plants by microbial interceded approach", *Symbiosis*, 2019

<1 %

Publication

Exclude quotes On

Exclude bibliography On

Exclude matches Off