

REVIEWER JURNAL INTERNASIONAL BEREPUTASI

Biodiversitas Journal of Biological Diversity

(<https://smujo.id/biodiv/index>)

BIODIVERSITAS

Volume 18, Number 2, April 2017

Pages: 659-665

ISSN: 1412-033X

E-ISSN: 2085-4722



DOI: 10.13057/biodiv/d180230

Dynamics of mangrove community in revegetation area of Karangsong, north coast of Indramayu District, West Java, Indonesia

HENDRA GUNAWAN*, SUGIARTI, SOFIAN ISKANDAR

Forest Research and Development Center, FORDA, Ministry of Environment and Forestry, Jl. Gunung Batu No. 5, Bogor 16118. P.O. Box 165, West Java, Indonesia. Tel. +62-251-8633234, 7520067; Fax. +62-251-8638111; *email: hendragunawan1964@yahoo.com

Indexing The journal has been indexed/registered in SCOPUS, DOAJ, Google Scholar, Crossref, EBSCO, Microsoft Academic Search, etc.





Abdul Malik <abdulmalik@unm.ac.id>

Fwd: Invitation to review

6 messages

Managing Editor <unsjournals@gmail.com>
To: malik@ign.ku.dk, abdulmalik@unm.ac.id

Fri, Aug 5, 2016 at 11:41 AM

Dear **Dr. Abdul Malik**,

We received a manuscript titled: **Dynamics of mangrove community in revegetation area of Karangsong, north coast of Indramayu District, West Java, Indonesia.**

We will be most grateful if you could find time to review the manuscript. The time is 2-3 weeks. Please find attached the full manuscript and Guidance for Author.

You may present your evaluations the way you deem fit. If you wish to correct parts of the manuscript itself, please indicate your corrections with a different color (for example, red for addition and blue for deletion) or use track change (for MS word program).

Kindly send me a mail to acknowledge the receipt.

Thank you,
Best Regards,

Ahmad Dwi Setyawan

Managing Editor,
- Biodiversitas, Journal of Biological Diversity (biodiversitas.mipa.uns.ac.id) (SCOPUS, DOAJ)
- Nusantara Bioscience (biosains.mipa.uns.ac.id/nusbioscience.htm) (Web of Science (ESCI), DOAJ)

Chairman/Co-Chairman
- National Seminar & International Conference on Biodiversity, <http://biodiversitas.mipa.uns.ac.id/snmbi.html>

Department of Biology,
Faculty of Mathematics and Natural Sciences, Sebelas Maret University,
Jl. Ir. Sutami 36A Solo 57126, Central Java, Indonesia,
Tel. & Fax. +62-271-663375,
e-mail: unsjournals@gmail.com

2 attachments

 **!Biodiversitas-Gudance for Authors.pdf**
208K

 **Dynamics of mangrove community in revegetation area of Karangsong.doc**
851K

Abdul Malik <abdulmalik@unm.ac.id>
To: Managing Editor <unsjournals@gmail.com>

Fri, Aug 5, 2016 at 3:47 PM

Dear Ahmad Dwi Setyawan
Managing Editor of Biodiversitas, Journal of Biological Diversity

Thank you for the invitation to review the manuscript. I will go through the manuscript and send back the result in the determined time.

I would like also inform to you for the next correspondence through this email address only.

Best regards,
Abdul Malik, Ph.D.

[Quoted text hidden]

--
Abdul Malik, Ph.D.

Lecturer at Department of Geography
Faculty of Mathematics and Natural Sciences
Universitas Negeri Makassar (UNM)
Jl. Malengkeri Raya, Kampus Parangtambung Makassar, 90224
South Sulawesi - INDONESIA

Managing Editor <unsjournals@gmail.com>
To: Abdul Malik <abdulmalik@unm.ac.id>

Fri, Aug 5, 2016 at 3:53 PM

Dear Sir,

Thank you so much for your willingness.

Thank you,
Regards,

Ahmad Dwi Setyawan

Managing Editor,
- Biodiversitas, Journal of Biological Diversity (biodiversitas.mipa.uns.ac.id) (SCOPUS, DOAJ)
- Nusantara Bioscience (biosains.mipa.uns.ac.id/N/index.htm) (Web of Science (ESCI), DOAJ)

Chairman/Co-Chairman
- National Seminar & International Conference on Biodiversity, <http://biodiversitas.mipa.uns.ac.id/snmbi.html>

Department of Biology,
Faculty of Mathematics and Natural Sciences, Sebelas Maret University,
Jl. Ir. Sutami 36A Solo 57126, Central Java, Indonesia,
Tel. & Fax. +62-271-663375,
e-mail: unsjournals@gmail.com

[Quoted text hidden]

Abdul Malik <abdulmalik@unm.ac.id>
To: Managing Editor <unsjournals@gmail.com>

Wed, Aug 24, 2016 at 11:47 AM

Dear Editor of Biodiversitas (Ahmad Dwi Setyawan)

Please find attached the reviewed manuscript.

Best regards,
Abdul Malik

[Quoted text hidden]

 **Dynamics of mangrove community in revegetation area of Karangsong_AM.doc**
851K

Managing Editor <unsjournals@gmail.com>
To: Abdul Malik <abdulmalik@unm.ac.id>

Wed, Aug 24, 2016 at 5:49 PM

Dear Sir,

Thank you so much for the evaluation.

Thank you,
Best Regards,

Ahmad Dwi Setyawan

Managing Editor,
- Biodiversitas, Journal of Biological Diversity (biodiversitas.mipa.uns.ac.id) (SCOPUS, DOAJ)
- Nusantara Bioscience (biosains.mipa.uns.ac.id/nusbioscience.htm) (Web of Science (ESCI), DOAJ)

Chairman/Co-Chairman
- National Seminar & International Conference on Biodiversity, <http://biodiversitas.mipa.uns.ac.id/snmbi.html>

Department of Biology,
Faculty of Mathematics and Natural Sciences, Sebelas Maret University,
Jl. Ir. Sutami 36A Solo 57126, Central Java, Indonesia,
Tel. & Fax. +62-271-663375,
e-mail: unsjournals@gmail.com

[Quoted text hidden]

Abdul Malik <abdulmalik@unm.ac.id>
To: reviews@webofscience.com

Sun, Aug 28, 2022 at 2:43 AM

Forwarded email corresponding review receipt from Biodiversitas, Journal of Biological Diversity, for your verification
(Review ID number: 16949578)

[Quoted text hidden]

--
Abdul Malik, Ph.D.

Department of Geography
Faculty of Mathematics and Natural Sciences
Universitas Negeri Makassar (UNM)
Kampus UNM Parangtambung, Jl.Malengkeri Raya, Makassar, 90224
South Sulawesi - INDONESIA
Phone: +62-853 9859 2785 Fax: +62-411-880568
E-mail: abdulmalik@unm.ac.id

Dynamics of mangrove community in revegetation area of Karangsong, north coast of Indramayu District, West Java, Indonesia

Mangrove along the north-North coast of Java is heavily degraded due to the conversion of land into fish ponds and human settlement areas. A revegetation program has been initiated by the local community of Karangsong Village, Indramayu District, supported by PT. Pertamina RU VI Indramayu. The research aimed to study the population dynamics of the mangrove revegetation in Karangsong. Secondary data was collected from the Fishery and Maritime Services of Indramayu and PT. Pertamina, while primary data was carried out on location by field observations and we interviewed key respondents. Data was analyzed to describe trends in the diversity index and population dynamics of the mangrove. The results showed that the revegetation effort in shoreline of Karangsong has covered ± 69.08 hectares which consisting of six species of mangrove and three tree species of coastal vegetation i.e. *Rhizophora mucronata* Lam., *Rhizophora stylosa* Griff., *Rhizophora apiculata* Blume, *Avicennia marina* (Forssk.) Vierh., *Avicennia alba* Blume, *Sonneratia caseolaris* (L.) Engl, *Terminalia catappa* L., *Casuarina equisetifolia* L., and *Ziziphus mauritiana* Lam. The mangrove population increased dramatically, from estimated 25,000 individuals in 2008 to 690,835 individuals in 2016. *Rhizophora mucronata* Lam. was the most dominant species (68.85%), followed by *Rhizophora stylosa* Griff. (18.33%) and *Rhizophora apiculata* Blume (9.53%). The Shannon diversity index was fluctuated but tends to be increase from 0.80 to 0.95.

Keywords: mangrove, re-vegetation, Indonesia, West Java, north coast, Karangsong.

INTRODUCTION

There are roughly 166,876 km² of mangrove habitat along the shorelines of the world, with the largest proportion of mangrove occurring in Asia (77,169 Km²) and the Americas (43,161 Km²) (Valiela et al. 2001). Countries with the largest area of mangroves are Indonesia (4.25 x 10⁴ km²) (Spalding et al. 1997), followed by Brazil (1.34 x 10⁴ km²) (Spalding et al. 1997), Nigeria (1.05 x 10⁴ km²) (Saenger & Bellan 1995), and Australia (1.00 x 10⁴ km²) (Robertson & Duke 1990).

Globally, the area of mangrove area is declining rapidly as it is cleared and converted to mariculture, agriculture, urban development, logged timber concessions, and fuel production areas (Fortes 1988; Marshall 1994; Primavera 1995; Twilley 1998; Polidoro et al. 2010). At least 35% of the world's mangrove forest area has been lost in the past two decades (Valiela et al. 2001). It is apparent that maricultural practices are responsible for the bulk of the increasing loss of mangrove worldwide. For example, pond culture has been reported to be responsible for 50%–80% of the loss of mangrove in Southeast Asia (Wolanski et al. 2000). Most of the damage is attributable to the direct loss of habitat from conversion of “cheap” mangrove land to “valuable” shrimp, prawn, and fish ponds (Valiela et al. 2001).

In 1999, Indonesia's mangrove forest covered 8.6 million hectares which consisted of 3.8 million hectares of forest area and 4.8 million hectares of non forest area. Degradation of mangrove in forest area is 1.7 million hectares (44.73%) and in non forest area is 4.2 million hectares (87.50%) (Gunawan & Anwar 2005). Indonesia has lost 40% of its mangrove in the last three decades (FAO 2007). The deforestation rate for mangrove in Indonesia is estimated to be 6% or 0.05 million hectares of the total annual forest loss (Margono et al 2014; Ministry of Forestry Republic of Indonesia 2014). The Ministry of Forestry has reported that only 31% of the remaining mangrove is in an intact condition and the rest (69%) is heavily degraded (Ministry of Forestry 2007). FAO (2007) reported that mangrove forest in Indonesia is 3,062,300 hectares or 19% of the world's mangrove and still the largest in the world, followed by Australia and Brazil.

Mangrove forest in Java Island is decreasing as the impact of conversion to mariculture, human settlement and other uses worsens. This impact is due to limited understanding and awareness by surrounding communities of the ecological importance of mangrove and to uncertainty about land status (Said & Smith, 1997). In 2011, mangrove in West Java Province was estimated as covering 40,130 hectares which was distributed between forest area 32,314 ha (80.52%) and non-forest area, 7,816 ha (19.48%), including 13 regencies-Districts (Forestry Service of West Java Province 2013). The degraded mangrove in this province is 15,276 hectares (38.06%), with the largest occurring in Karawang District 13,181ha (32.85%) followed by Bekasi 10,481ha, Indramayu 8,720ha, Subang 7,346 ha, Cirebon 190 ha, Ciamis 170 ha, Garut 32 ha and Sukabumi 9 ha (Ministry of Forestry 2012). The loss of mangrove in the Indramayu District has impacted on the disappearance of Ujung Gebang, Limbangan and Jatinyuat villages (Forestry Service of West Java Province 2016).

Mangrove forest in Indonesia provides benefits for local communities; supporting livelihoods by producing items of food, fuel wood, charcoal, construction materials, and furniture timber, as well as by generating income (Armitage 2002). Mangrove is also important in social-cultural terms in fulfilling various religious, spiritual, aesthetic, and recreational

Commented [AM1]: I suggest to use the latest data and references such as from Giri et al (2011), Richards and Friess (2016) and Bakosurtanal 2009).

Commented [AM2]: I suggest to use one of area units (km or hectare) in your manuscript for facilitate the reader to understand.

Commented [AM3]: I think this sentence or information could combine to the first paragraph (line 22-25).

Commented [AM4]: Please consistent to “Regency” or “District”

53 functions that benefit ecotourism (UNEP 2014). Mangrove ecosystems support essential ecological functions such as
 54 intercepting land-derived nutrients, pollutants, and suspended matter before these contaminants reach deeper water
 55 (Marshall 1994, Rivera-Monroy and Twilley 1996, Tam and Wong 1999). Mangroves also perform other important
 56 services, such as preventing coastal erosion by stabilizing sediments (Marshall 1994, Tam & Wong 1999), furnishing
 57 nursery and spawning areas for commercially important coastal fish and shellfish species (Rodelli et al. 1984, Sasekumar
 58 et al. 1992), and providing stopover sites for migratory birds, fish, and mammals (Saenger et al. 1983). Any loss of
 59 mangrove forest, therefore, means a loss of their important contributions to subsistence uses, and to ecological, economic,
 60 and conservation functions (Valiela et al. 2001).


61 Based on the essential functions of mangrove for human life, the Ministry of Environment and Forestry [Republic of](#)
 62 [Indonesia](#) has designated mangrove as an essential ecosystem which will be treated as a protected area or conservation area
 63 under the Directorate of Essential Ecosystems Management. The Ministry of Environment and Forestry [Republic of](#)
 64 [Indonesia](#) has also launched a National Movement on Forest and Land Rehabilitation (NMFLR) - a national initiative to
 65 plant trees in forest land and bare lands - including mangroves - as a commitment to improving the quality of environment
 66 for people's prosperity. The total extent of the national program for mangrove rehabilitation during 2010-2014 is 33,394
 67 hectares (Ministry of Environment and Forestry 2015). The rehabilitation program in West Java Province has planted 365
 68 hectares of mangrove in 2008, 50 hectares in 2009, 311 hectares in 2010, 480 hectares in 2011 and 270 hectares in 2012
 69 (Forestry Service of West Java 2013).

70 The local community of Karangsong village in Indramayu District which is supported by PT. Pertamina Refinery Unit
 71 VI Indramayu has initiated a mangrove re-vegetation program on private land (non- forest land) along the ~~north~~-North
 72 shoreline of Indramayu District. They started planting mangrove ~~species~~ in 2008 and have consistently extended the area of
 73 rehabilitation in a project that has involved many stakeholders. They are also developing an ecotourism program and
 74 are practicing sustainable utilization of non-timber mangrove products for generating income to raise local people's
 75 prosperity.

76 The research described here, aimed to study the population dynamics of this mangrove vegetation that has been planted
 77 in Karangsong village, Indramayu District.

78 **MATERIALS AND METHODS**

79 This research was conducted in Karangsong ~~village~~Village, Sub District of Indramayu, Indramayu District. The
 80 research site is located between 6°17'38.52"S - 6°18'17.52"S and 108°22'03.60"E - 108°22'17.94"E, on the ~~north~~-North
 81 coast of Indramayu District. The site is a mangrove habitat combined with coastal habitat and the estuary of the
 82 Prajagumiwang River which crosses the Karangsong Village and joins with the Java Sea. This area is part of the Cimanuk
 83 watershed, with the main Cimanuk River crossing the territory of Indramayu District. Secondary and primary data were
 84 collected on May to June 2016.



107 **Figure 1.** Research location.

Commented [AM5]: Due to your references for this information are too old for the publication 2016, I I guess you could find new references for this citation.

Commented [AM6]: I guess you can find new references for this citation.

Commented [AM7]: I guess you can find new references for this citation

Commented [AM8]: Due to protection and conservation mangrove area become International program, I suggest you to put broader information regarding it (not only in Indonesia). It also become international readers can be more interested and make your paper stronger.

Commented [AM9]: I suggest to re-draw your research location map rather than just cut and put images from Google map and other sources. In addition, please provide the map legend in English and the sign of North arrow as one of important map requirement.

109 ~~Data of about the mangrove~~ Mangrove plantation data was collected from the Fishery and Maritime Service of
110 Indramayu District and PT. Pertamina RU VI Indramayu. Information collected included the species, the number of
111 plants, the date of planting, an estimation of the site extent, and details of the institutions and community groups involved
112 in the plantation. In-depth interviews with resource persons and key respondents were needed to complete and confirm the
113 data and information collected. The point count method was applied for the bird survey (Hill et al. 2005) with an
114 observation radius of 50 m: the result was a list of bird species (van Lavieren 1982). Identification of birds referred to the
115 Field Guide for the Birds of Java and Bali (Mackinnon 1991) and Field Guide to the Birds of Borneo, Sumatra, Java and
116 Bali (MacKinnon et al. 1992).

117 All information and data were analyzed to describe the dynamics of the ecosystem, including the abundance trend,
118 species composition, and diversity and evenness indices of the mangrove population. The results will be used to evaluate
119 the program of mangrove rehabilitation and provide recommendations for future action.

Commented [AM10]: How many resource persons and key respondents did you interview? And what method did you use to select them? And Where they come from (what institution)?

Commented [AM11]: I suggest to move this sentence in the last of introduction part. "The results of this research will be used to..."


120 RESULTS AND DISCUSSION

121 National Program of Rehabilitation on Forest and Bare Land

122 The Ministry of Environment and Forestry has launched a National Movement on Forest and Land Rehabilitation
123 (NMFLR). It is a national initiative to plant trees in forest land and bare land, including mangroves, as a commitment to
124 improving the quality of environment for people's prosperity. In the period 2010-2014, the NMFLR program planted
125 2,279,380 hectares. The national program for rehabilitation of degraded mangrove in the period 2010 – 2014 planted
126 33,394 hectares (Ministry of Environment and Forestry 2015). In the decade 2003-2012, mangrove rehabilitation in West
127 Java Province through the NMFLR program covered 3,681 hectares (Figure 2) (Forestry Service of West Java Province
128 2012; 2013a; 2013b; 2016). Rehabilitation of mangrove along the north-North coast of Java is crucial, and not only for
129 ecological reasons; the socio-economic arguments are even more significant, due to the threat that mangrove degradation
130 poses to the surrounding communities who depend on mangrove and the fishery it sustains. Gunawan et al. (2007a)
131 reported that the presence of mangrove can improve the quality of water in fish ponds. Gunawan et al. (2007b) also found
132 that mangrove rehabilitation through a silvofishery program can increase the household income of the adjacent community
133 of Subang District, on the north-North coast of West Java Province.

Commented [AM12]: I suggest to change or delete this beginning statement due to has been written previously on the line 63-66.

Commented [AM13]: Very descriptive. I suggest to analyze and describe the trend of mangrove rehabilitation in West Java, and how is their contribution to the national program for mangrove rehabilitation and especially from the North Coast of Java.



137
138
139 Figure 2. Trend of mangrove rehabilitation in West Java Province.
140 Sources: Forestry Services of West Java Province (2012; 2013a,b; 2016);
141 Ministry of Forestry (2007; 2012; 2014); Ministry of Environment and Forestry (2015)
142

143 Mangrove Rehabilitation in Karangsong Village, Indramayu District

144 The revegetation program to rehabilitate degraded mangrove in Karangsong village-Village was initiated in 2003
145 through the planting of three species of mangrove (*Rhizophora mucronata*, *R. stylosa*, *R. apiculata*) covering 2.5 hectares
146 of shoreline in Karangsong Village. The plantation was initiated by a group of fishermen named "Kelompok Pantai
147 Lestari" who were supported by PT. Pertamina RU VI Indramayu through the Corporate Social Responsibility (CSR)
148 program. One of the crucial reasons for the involvement PT. Pertamina RU VI, is that the Java Sea to the north-North of
149 Indramayu is a route for oil tankers transporting refined oil from Balongan Indramayu to Jakarta. The sea has been
150 polluted by oil spills from tankers, which negatively impacts on the local sea-water quality and on the fishery. The
151 suspicion occurs based on the that this was happening was supported by the research results of Gunawan and Anwar
152 (2008) report, who detected the pollutants Lead (Pb), detergent (MBAS) and Mercury (Hg) in the waters of Subang
153 District; (a District on the north-North coast next to Indramayu District). In this area, Gunawan and Anwar They found

Commented [AM14]: If you use the taxonomic authority name after species name, please consistent throughout your paper text.

that the lead (Pb) content and detergent (MBAS) of waters were higher than the threshold for fishery culture. Furthermore, They-they also found that eight species of fishes and a species of shrimp in silvofishery ponds, and six species of fishes and a species of shrimp in common ponds without mangrove were contaminated with mercury (Hg). However, but the concentration of pollutants in the silvofishery ponds was lower than that-of-the common ponds.

It is believed that revegetation of mangrove can improve the quality of the coastal-sea water and in turn restore the habitat of the-biota living in the water. As a corporation who-that produces and transports oil through the Java Sea to the north-North of Indramayu, PT. Pertamina RU VI has a high commitment to restore the mangrove ecosystem and coastal sea waters along the north-North shoreline of Indramayu. The mangrove revegetation program which was initiated by the people of Karangsang and PT. Pertamina RU VI was-has then-followed-involved-by-37-other community groups (Table 1) and 13 institutions including-and-supported-by NGOs, private sectors, national-National government, provincial-Provincial government and district-District government (Table 2).-There are 37 community groups and 13 institutions involved in the development-and-plantation-of mangrove at Karangsang (Table 1 and Table 2). However, The-the private sector has a critical role in mangrove rehabilitation at Karangsang (31%) alongside the national government (23 %) (Figure 3). The facts demonstrate that the success of mangrove rehabilitation program depends upon the involvement of all stakeholders. Gunawan & Anwar (2005) similarly found that the A similar success of mangrove rehabilitation that on-the-north-coast-of Central-Java-Province-is determined by the participation of the local community has been reported on the North coast of Central Java Province (Gunawan & Anwar (2005) around-the-mangrove-area.

Commented [AM15]: Why? Please provide more information regarding it!

Table 1. Community groups involved in mangrove revegetation of north-North coast of Indramayu District.

Community Group	Planting Site		
	Block	Village	Sub District
Karang Taruna Putra Balongan	Sawah laut, Pertamina, Kesambi, Pesisir, Balongan	Majakerta, Balongan	Balongan
Kelompok Dwi Jaya	Waki	Brondong	Pasekan
Kelompok Karya Muda	Salkri	Lamarantarung	Cantigi
Kelompok Lamaran Jaya	Agus	Lamarantarung	Cantigi
Kelompok LMDH Bangsal Sari	Bangsang	Pagirikan	Pasekan
Kelompok Rapi Jaya Putra	Tiris	Pabean Ilir	Pasekan
Kelompok Tani Anugerah	Kastal	Cangkring	Cantigi
Kelompok Tani Bala Dewo	Blubbuk	Totoran	Pasekan
Kelompok Tani Blubuk Sejahtera	Blukbuk	Totoran	Pasekan
Kelompok Tani Brawijaya Putra	Bangsang & Keci	Pasekan	Pasekan
Kelompok Tani Jaka Kencana	Udik	Pabean Udik	Indramayu
Kelompok Tani Langgeng Jaya	Singkil Tanah Timbul, Langen	Singaraja	Indramayu
Kelompok Tani LMDH Paluh Adin Jaya	Blubbuk	Totoran	Pasekan
Kelompok Tani Loka Jaya	Muara	Lamarantarung	Cantigi
Kelompok Tani Makmur Jaya	Muara	Lamarantarung	Cantigi
Kelompok Tani Mulia Jaya	Bangsang & Keci	Pasekan	Pasekan
Kelompok Tani Muncul Jaya Mangrove	Bangsang & Keci	Pasekan	Pasekan
Kelompok Tani Pal Jaya	Toeni	Lamarantarung	Cantigi
Kelompok Tani Pancer Pindang Jaya	Kastal	Cangkring	Cantigi
Kelompok Tani Pelangi Mangrove	Bangsang & Keci	Pasekan	Pasekan
Kelompok Tani PH Pabean Hilir	Bangsang & Keci, Blukbuk	Pasekan, Pabean Ilir	Pasekan
Kelompok Tani Putra Kujang	Kastal, Toeni	Cangkring, Lamarantarung	Cantigi
Kelompok Tani Sea Green	Bangsang & Keci	Pasekan	Pasekan
Kelompok Tani Sejahtera Mangrove	Bangsang & Keci	Pasekan	Pasekan
Kelompok Tani Sidum Jaya	Bangsang & Keci	Pasekan	Pasekan
Kelompok Tani Sigra Mongso	Waki, Kiper	Brondong	Pasekan
Kelompok Tani Sinar Jaya Mangrove	Bangsang & Keci	Pasekan	Pasekan
Kelompok Tani Terumbu Karang	Jangin	Cangkring	Cantigi
Kelompok Tani Tumbuh Hijau	Jangin	Cangkring	Cantigi
Kelompok Tani Tunas Jaya	Bangsang	Pagirikan	Pasekan
Kelompok THP Nandur Jaya	Sawah Laut	Pabean Ilir	Pasekan
Kelompok THP Sumber Urip	Sawah Laut	Pabean Ilir	Pasekan
Kelompok THP Tumbuh Jaya	Perlat	Karanganyar	Pasekan
Kelompok Tunas Lestari	Payang	Pabean Ilir	Pasekan

Kelompok Usaha Bersama Tiris Berseri	Tegur Baru	Pabean Ilir	Pasekan
Kelompok Pantai Lestari*)	Karangsong**)	Karangsong	Indramayu
LSM Siklus	Pabean Udik	Pabean Udik	Indramayu

Source: PT. Pertamina RU VI, Indramayu (year?). *) Manager of Karangsong Mangrove area. **) Karangsong mangrove area

Table 2. Institutions involved in mangrove plantation in north-the North coast of Indramayu District and infrastructure development at Karangsong mangrove area.

Institutions	Contribution	Local Counterpart	Site
Kementerian Kelautan dan Perikanan	Planting, Infrastructure	Kelompok Pantai Lestari	Pabean Udik dan Brondong
Balai Besar Wilayah Sungai Cimanuk-Cisanggarung, Kementerian Pekerjaan Umum	Planting, Infrastructure	Kelompok Pantai Lestari	Karangsong
Balai Pengelolaan Hutan Mangrove Wilayah I Bali, Kementerian Lingkungan Hidup dan Kehutanan	Nursery	Kelompok Pantai Lestari	Karangsong
Dinas Perikanan dan Kelautan Prov. Jawa Barat	Planting	Kelompok Pantai Lestari, LSM SIKLUS	Karangsong, Pabean Udik
Dinas Perikanan dan Kelautan Kab. Indramayu	Planting	Kelompok Pantai Lestari	Karangsong
PSL IPB Program Magister Tahun 2014	Planting	Kelompok Pantai Lestari	Karangsong
Alumni Fahatan IPB (Angkatan E.27)	Planting	Kelompok Pantai Lestari	Karangsong
PT. BioFarma Bandung	Planting	LSM SIKLUS	Pabean Udik
PT. PLN	Planting	LSM SIKLUS	Pabean Udik
PT. Traktor Nusantara Jakarta	Planting	Kelompok Pantai Lestari	Karangsong
PT. Pertamina RU VI – Balongan, Indramayu	Planting, Infrastructure	Kelompok Pantai Lestari	Karangsong
MFF Indonesia (Mangrove For Future – UNDP)	Planting	Kelompok Pantai Lestari	Pabean Udik dan Karangsong
Yayasan KEHATI Jakarta	Planting	Kelompok Pantai Lestari	Karangsong

Source: PT. Pertamina RU VI, Indramayu (Year?).




Figure 3. Supporting institutions of mangrove rehabilitation at Karangsong Village, Indramayu District.

The extent of the rehabilitated mangrove area is has been increasing from year-2008 to 2016 year in Karangsong Village (Figure 4). There has been a particularly significant increase occurred from 2014 to 2016. In the first semester of 2016, the extent of mangrove revegetation in the north-North shoreline of Indramayu District has covered ±69.08 hectares. Karangsong Village in Sub District of Indramayu has the largest area of mangrove revegetation. Mangrove revegetation has been also implemented in other sub districts i.e. Balongan (...hectares), Cantigi (...hectares) and Pasekan (...hectares). The cumulative extent of green-belt-of mangrove revegetation in shoreline-of Indramayu District is estimated 103.19 hectares. This increase was triggered by national and international events such as the International Forest Day, Conservation Day, Environment Day, Tree Planting Day, One Billion Trees Program and One Man One Tree conducted at Karangsong and surroundings.

Commented [AM16]: Due to you provided data per year (2008-2016) (fig 4.) I think the significance increase occurred only from 2014 to 2015 not include 2016, right?

Commented [AM17]: It is confused! In fig 4. You showed the extent of mangrove revegetation in Karangsong from 2008-2016 and in first semester 2016 has covered 69.08 ha.

Commented [AM18]: Please provide data from other areas such as Balongan, Cantigi and Pasekan to support the statement that "Karangsong is the largest area of mangrove revegetation".




Figure 4. Cumulative extent of mangrove revegetation area at Karangsong Village, Indramayu District.

The increase in the extent of rehabilitated area has been accompanied by an increase in the population of mangrove species (Figure 5). *Rhizophora mucronata* is dominating the plantation (68.85%), followed by *R. stylosa* (18.33 %) and *R. apiculata* (9.53%) (Figure 6). The number of species has also increased from three species in 2008 to nine species in 2016, which consisted of six species of mangrove and three species of coastal vegetation trees (Table 3 and Figure 7). Compared with natural mangrove in Indonesia which consists of trees (at least 47 species), shrubs (5 species), herbs and grasses (9 species), and parasites (2 species) (Kusmana 2011), revegetation at Karangsong still needs to be diversified in terms of the range of species generated in the plantation areas. Increasing the species richness is critical to providing heterogeneous habitat for faunal diversity. The majority of studies have found a positive correlation between habitat heterogeneity/diversity and animal species diversity, although ecological effects of habitat heterogeneity may vary considerably between species groups depending on whether structural attributes are perceived as heterogeneity or fragmentation. Moreover, the effect of habitat heterogeneity for one species group may differ in relation to the spatial scale (Tews et al. 2004).

Although the population number of each species is increasing, the composition is not evenly distributed. This is indicated by the values for diversity and evenness indices. Figure 8 shows the change in values of diversity and evenness indices of mangrove during the period of 2008-2016. The indices have not been continuously increasing. In particular, the mass planting in 2014 – 2016 resulted in a decline in the diversity index as well as evenness index.




Figure 5. Population increase of mangrove and coastal species at Karangsong Village, Indramayu District.

Commented [AM19]: Please consistent to use the word "District"

This fig. just showed the extent of mangrove revegetation per year from 2008-2016, but not covered the cumulative extent. So please change the title of the fig.

Commented [AM20]: I suggest the discussion here provide the information why these *Rhizophora* sp is dominated here and even in Southeast Asia? You could use paper from Malik et al. (2015) Mangrove Exploitation Effects on Biodiversity and Ecosystem Services. *Biodiversity and Conservation* 24: 3543-3557 and Ellison AM (2000) Mangrove restoration: do we know enough? *Restor Ecol* 8:219-229 as your references.

195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214

215
216
217




Figure 6. Composition of plant species in the rehabilitation area of Karangsong, Indramayu.

Table 3. List of species at Karangsong rehabilitation area.

Local Name	Botanic Name	Family	IUCN Red List Category	Habitat
Bakau hitam	<i>Rhizophora mucronata</i> Lam.	Rhizophoraceae	LC (ver 3.1)	Mangrove
Bakau kecil	<i>Rhizophora stylosa</i> Griff.	Rhizophoraceae	LC (ver 3.1)	Mangrove
Bakau minyak	<i>Rhizophora apiculata</i> Blume	Rhizophoraceae	LC (ver 3.1)	Mangrove
Api-api	<i>Avicennia marina</i> (Forssk.) Vierh.	Acanthaceae	LC (ver 3.1)	Mangrove
Api-api	<i>Avicennia alba</i> Blume	Acanthaceae	LC (ver 3.1)	Mangrove
Pidada	<i>Sonneratia caseolaris</i> (L.) Engl.	Lythraceae	LC (ver 3.1)	Mangrove
Ketapang	<i>Terminalia catappa</i> L.	Combretaceae	LR/nt (ver 2.3)	Coastal
Cemara laut	<i>Casuarina equisetifolia</i> L.	Casuarinaceae	NE (ver 3.1)	Coastal
Bidara	<i>Ziziphus mauritiana</i> Lam.	Rhamnaceae	NE (ver 3.1)	Coastal

Enrichment planting is critically important to increase the diversity of mangrove in order to enhance quality and heterogeneity of habitats for promoting fauna diversity. Azlan et al. (2015) stressed the importance of diversity and quality of habitat in encouraging the diversity and density of birds in mangroves. Bird species composition in mangroves was closely associated with both plant species composition and configuration of the vegetation structure (Azlan et al 2015). Habitat structure and floristic characteristics is also closely related to species richness and diversity of birds. Larger areas tend to have more diverse habitats, both structurally and floristically, which bird species can occupy, resulting in greater bird diversity (MacArthur & Wilson 1967; Woinarski et al. 2001). Besides being an important factor in contributing to the increase of species richness and diversity, habitat structure is also an important determinant influencing habitat selection and distribution of species, especially in complex habitats such as tropical forest (Watson et al. 2004). Habitat heterogeneity in mangrove is less pronounced and may limit the number of coexisting species (Ford 1982).




Figure 7. Increase of species number of plant at Karangsong mangrove area, Indramayu.



Figure 8. Dynamics of species diversity index of vegetation at Karangsong mangrove area, Indramayu.

Enrichment planting should include an increase in species number and a balancing of the proportions among the species, so that the evenness index increases. Some species of mangrove that could be added to enrich the Karangsong mangrove area are Tanjung (*Bruguiera* sp.), Nyirih (*Xylocarpus* sp.), Tengar (*Ceriops* sp.) and Buta-butua (*Excoecaria* sp.). These have not been planted yet in Karangsong. Sandy coastal habitat should also be enriched with coastal species such as Butun (*Barringtonia asiatica* (L.) Kurz.), Nyamplung (*Calophyllum inophyllum* L.), Bintaro (*Cerbera manghas* L.), Ketapang (*Terminalia catappa* L.), Kampis cina (*Hernandia peltata* Meisn.), Waru (*Hibiscus tiliaceus* L.), Waru laut (*Thespesia populnea* (L.) Sol. Ex Correa), Kepuh (*Sterculia foetida* L.), Dungun (*Heritiera littoralis* Aiton), and Malapari (*Pongamia pinnata* (L.) Pierre).

Impact of Mangrove Revegetation

The presence of mangrove revegetation in the north-North coast of Karangsong Village has gradually encouraged birds and other faunas. Twelve families of birds consisting of twenty species were found in mangrove and coastal vegetation at Karangsong (Table 4). Eight species of water birds were found in the mangrove habitat. These birds are a very common presence in the mangrove of Karangsong and some of them are temporary residents. The area of the North coast of Java still having mangrove is essential habitat for migrant birds. In the North coast of Indramayu, Iskandar and Karlina (2004) reported 15 species of migrant birds and some of them migrant birds are consumed and sold by local people for additional income (Iskandar & Karlina 2004). The role of mangrove as habitat of wildlife was also demonstrated by Gunawan (2002). Gunawan (2002) He found 77 species of wildlife, consisting of three mammals, six reptiles and 68 birds that directly interacted with the mangrove vegetation in Rawa Aopa Watumohai National Park (RAWNP) Southeast Sulawesi. The mangrove of RAWNP is a secure home for endangered species that depend on mangroves and is a main transit habitat for many seasonally migrant birds.

Table 4. List of birds species found at Karangsong mangrove area.

Local Name	Latin Name	Family	IUCN Red List Category Ver.3.1
Cekakak sungai	<i>Todirhampus chloris</i> Boddaert	Alcedinidae	Least Concern (LC)
Meninting	<i>Alcedo meninting</i> Horsfield	Alcedinidae	Least Concern (LC)
Walet sapi	<i>Collocalia esculenta</i> Linnaeus	Apodidae	Least Concern (LC)
Walet linchi	<i>Collocalia linchi</i> Horsfield & Moore	Apodidae	Least Concern (LC)
Kuntul kerbau	<i>Bubulcus ibis</i> Linnaeus	Ardeidae	Least Concern (LC)
Kuntul karang	<i>Egretta sacra</i> Gmelin	Ardeidae	Least Concern (LC)
Kuntul kecil	<i>Egretta garzetta</i> Linnaeus	Ardeidae	Least Concern (LC)
Kuntul perak	<i>Egretta intermedia</i> Wagler	Ardeidae	Not Evaluated (NE)
Blekok sawah	<i>Ardeola speciosa</i> Horsfield	Ardeidae	Least Concern (LC)
Kokokan laut	<i>Butorides striata</i> Linnaeus	Ardeidae	Least Concern (LC)
Cinene pisang	<i>Orthotomus sutorius</i> Pennant	Cisticolide	Least Concern (LC)
Tekukur biasa	<i>Streptopelia chinensis</i> Scopoli	Columbidae	Least Concern (LC)
Wiwik kelabu	<i>Cacomantis merulinus</i> Scopoli	Cuculidae	Least Concern (LC)
Bondol peking	<i>Lonchura punctulata</i> Linnaeus	Estrildidae	Least Concern (LC)
Layang-layang	<i>Hirundo tahtica</i> Gmelin	Hirundinidae	Least Concern (LC)
Bentet kelabu	<i>Lanius schach</i> Linnaeus	Laniidae	Least Concern (LC)
Gereja erasia	<i>Passer montanus</i> Linnaeus	Passeridae	Least Concern (LC)
Cucak kutilang	<i>Pycnonotus aurigaster</i> Vieillot	Pycnonotidae	Least Concern (LC)
Kacamata biasa	<i>Zosterops palpebrosus</i> Temminck	Zosteropidae	Least Concern (LC)
Kacamata laut	<i>Zosterops chloris</i> Bonaparte	Zosteropidae	Least Concern (LC)

Commented [AM21]: I think this part is not covered in your research findings. Need more research to recommend this. So I suggest to delete it or otherwise you could keep it but put a reference.

Commented [AM22]: What's the impact of revegetation to social economic (such as income) from developing ecotourism and to utilization non-timber mangrove product as mentioned on the line 73-75 and 128-130?

238
239
240
241
242
243
244
245
246
247
248

249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264

265

266
267
268
269
270
271
272
273
274
275

The number of mangrove species planted in Karangsang has increased from three in 2008 to nine in 2016, consisting of six species of mangrove and three species of coastal vegetation. The population of each species is increasing, with the dominant species *Rhizophora mucronata* (68.85%), followed by *R. stylosa* (18.33 %) and *R. apiculata* (9.53%). The Shannon's species diversity index was fluctuated but tends to be increase from 0.80 to 0.95. The success of mangrove revegetation depends on the participation of local people and community groups as well as on contributions from the private sectors with support from government at national and local levels. The presence of mangrove has provided habitat for a diversity of fauna, especially for the bird community. Enrichment planting is still needed to increase the diversity of mangrove, which impacts on the diversity of fauna.

276

ACKNOWLEDGEMENTS

277
278
279
280
281
282
283

I gratefully acknowledge PT. Pertamina RU VI Indramayu, PT. Pertamina Training and Consulting, Forestry Service of Indramayu District and Pantai Lestari Group. Special thanks go to Mr. Agung Darmawan, Mr. Ragil Agus Saputra, Mr. Dudi Junaidi Permana, Mr. Cecep Supriyatna, Mr. Oni Ngo, Mr. Dulloh, Mr. Ali Sodikin, Mr. Makrus, Mr Carita, Mr. Eka Tarika, Mr. Roksikin, Mr. Riza and Ms Tri. I am thankful to Mr. Graham Eagleton for his invaluable support in reviewing the manuscript. I also thank the local people of "Pantai Lestari Group" for cooperating and assisting in the conduct of this research. Thanks are also due to the volunteers for their valuable time assistance and efforts.

284

REFERENCES

- 285 Anwar Ch, Gunawan H. 2007. Ecological and socio-economic roles of mangrove in development of coastal zone. Pp.23-
286 34 in Siran, S. et al. (Eds). Expose of Research Result in Conservation and Rehabilitation of Forest Resources.
287 Proceeding, Padang September 20th, 2006. [Indonesia]
- 288 Armitage D. 2002. Socio-institutional dynamics and political ecology of mangrove forest conservation in Central
289 Sulawesi, Indonesia. *Global Environmental Change* 12(3): 203-2017.
- 290 Azlan JM, Noske RA, Lawes MJ. 2015. The role of habitat heterogeneity in structuring mangrove bird assemblages.
291 *Diversity* 7:118-136; doi:10.3390/d7020118.
- 292 Berg, Å. 1997. Diversity and abundance of birds in relation to forest fragmentation, habitat quality and heterogeneity.
293 *Bird Stud.* 44: 355-366.
- 294 Campbell A, Brown B. 2015. Indonesia's vast mangroves are a treasure worth saving. The Conversation from [http://the](http://theconversation.com/Indonesia-vas-mangroves-are-atreasure-worth-saving-39367)
295 [conversation.com/Indonesia-vas-mangroves-are-atreasure-worth-saving-39367](http://theconversation.com/Indonesia-vas-mangroves-are-atreasure-worth-saving-39367).
- 296 Cousin JA, Phillips RD. 2008. Habitat complexity explains species-specific occupancy but not species richness in a
297 Western Australian woodland. *Aust. J. Zool.* 56: 95-102.
- 298 Donato DC, Kauffman JB, Murdiyoso D, Kurnianto S, Stidham M, Kanninen M. 2011. Mangroves among the most
299 carbon-rich forests in the tropics. *Nature geoscience* 4(5): 293-297.
- 300 FAO. 2007. The world's mangroves 1980-2005. Forest Resources Assessment Working Paper No. 153. Rome. Food
301 and Agriculture Organization of the United Nations.
- 302 Ford, J. 1982. Origin, evolution and speciation of birds specialized to mangroves in Australia. *Emu* 82: 12-23.
- 303 Forestry Services of West Java Province. 2013a. Base and strategy of sustainable mangrove management in West Java.
304 Bandung. Forestry Services of West Java Province. [Indonesia]
- 305 Forestry Services of West Java Province. 2016. Mangrove forest: the destruction of the abrasion protector.
306 www.dishut.jabarprov.go.id [Indonesia]
- 307 Forestry Services of West Java Province. 2012. West Java Forestry Statistic 2012. Bandung. Forestry Services of West
308 Java Province. [Indonesia]
- 309 Forestry Services of West Java Province. 2013b. Technical Guide for Mangrove and Coastal forest management in West
310 Java. Bandung. Forestry Services of West Java Province. [Indonesia]
- 311 Fortes M. 1988. Mangrove and sea grass beds of East Asia: Habitats under stress. *Ambio* 17: 207-213.
- 312 Giri C, Ochieng E, Tieszen L.L, Zhu Z, Singh A, et al. 2011. Status and distribution of mangroves forests of the world
313 using earth observation satellite data. *Global Ecology and Biogeography* 20(1): 154-159.
- 314 Gunawan H, Anwar Ch, Sawitri R, Karlina E. 2007a. Ecological status of silvofishery model of empang parit in sub forest
315 concession of Ciasem-Pamanukan, forest concession unit of Purwakarta. *Jurnal Penelitian Hutan dan Konservasi Alam*
316 Vol. IV (4): 429-439. [Indonesia]
- 317 Gunawan H, Anwar Ch, Sawitri R, Karlina E. 2007b. The role of silvofishery in increasing household income of the
318 adjacent community and conserving mangrove in the sub forest concession of Ciasem-Pamanukan, forest concession
319 unit of Purwakarta. *Info Hutan* Vol. IV (2): 153-163. [Indonesia]

Commented [AM23]: We don't find this reference in your text.
So please delete it.

Please make sure that all the references in the list used in the text.

Commented [AM24]: We don't find this reference in your text.
So please delete it.

Commented [AM25]: We don't find this reference in your text.
So please delete it.

Commented [AM26]: We don't find this reference in your text.
So please delete it.

320 Gunawan H, Anwar Ch. 2005. An Analyses on the Success of Mangrove Rehabilitation in the North Coast of Central
321 Java. *Info Hutan* Vol. II (4): 239-248. [Indonesia]

322 Gunawan H, Anwar Ch. 2008. The quality of waters and Mercury (Hg) content of fishes in silvofishery pond at Sub
323 Forest District of Ciasem-Pamanukan, Forest District of Purwakarta, Subang District, West Java. *Jurnal Penelitian*
324 *Hutan dan Konservasi Alam* Vol. V (1): 1-10. [Indonesia]

325 Gunawan H. 2002. The role of mangroves as habitat of wildlife in Rawa Aopa Watumohai National Park, South East
326 Sulawesi Province. *Buletin Penelitian Kehutanan* Vol.8 (2): 17 – 35. [Indonesia]

327 Hill D, Fasham M, Tucker G, Shewry M, Shaw P. (eds). 2005. Handbook of biodiversity methods: survey, evaluation and
328 monitoring. Cambridge University Press. Cambridge, UK.

329 Honkanen M, Roberge, JM, Rajaärkkä A, Mönkkönen M. 2009. Distinguishing the effects of area, energy and habitat
330 heterogeneity on boreal forest bird species richness in protected areas. *Glob. Ecol. Biogeogr.* 18: 61–71.

331 Iskandar S, Karlina E. 2004. Kajian pemanfaatan jenis burung air di pantai utara Indramayu, Jawa Barat. *Buletin Plasma*
332 *Nutfah* 10(1): 43-48. [Indonesia]

333 Kusmana C. 2011. Management of mangrove ecosystem in Indonesia. *JPSL* (1)2: 152- 157

334 MacArthur, R.H.; Wilson, E.O. 1967. *The Theory of Island Biogeography*. Princeton University Press: Princeton, NJ,
335 USA.

336 MacKinnon J, Phillips K, van Balen B. 1992. *A Field Guide to the Birds of Borneo, Sumatra, Java and Bali*. Birdlife
337 International – Indonesia Program. Bogor. [Indonesia]

338 MacKinnon J. 1991. *Field Guide to the Birds of Java and Bali*. Gadjah Mada University Press. Yogyakarta. [Indonesia]

339 Margono BA, Potapov PV, Turbanova S, Stolle F, Hansen MC. 2014. Primary forest cover loss in Indonesia over 2000-
340 2012. *Nature Climate Change*.

341 Marshall N. 1994. Mangrove conservation in relation to overall environmental consideration. *Hydrobiologia* 285: 303–
342 309.

343 Ministry of Environment and Forestry. 2015. *Ministry of Environment and Forestry Statistics 2014*. Jakarta. Ministry of
344 Environment and Forestry. [Indonesia]

345 Ministry of Forestry Republic of Indonesia. 2014. *Recalculation of Indonesia’s land cover in 2013*. Directorate General
346 of Planology, Ministry of Forestry Republic of Indonesia. [Indonesia]

347 Ministry of Forestry Republic of Indonesia. 2007. *Statistic of Indonesian Forestry*. Jakarta. Ministry of Forestry Republic
348 of Indonesia. [Indonesia]

349 Ministry of Forestry Republic of Indonesia. 2012. *Forestry profiles of 33 provinces*. Jakarta. Ministry of Forestry.
350 [Indonesia]

351 Polidoro BA, Carpenter KE, Collins L, Duke NC, Ellison AM, Ellison JC, et al. 2010. The lost of species: Mangrove
352 extinction risk and geographic areas of global concern. *PLoS ONE* 5(4): e10095. Doi:10.1371/journal.pone.0010095.

353 Primavera JH. 1995. Mangroves and brackish water pond culture in the Philippines. *Hydrobiologia* 295: 303–309.

354 Rivera-Monroy VH, Twilley RR. 1996. The relative role of denitrification and immobilization in the fate of inorganic
355 nitrogen in mangrove sediments. *Limnology and Oceanography* 41: 284–296.

356 Robertson AI, Duke NC. 1990. Recruitment, growth and residence time of fishes in a tropical Australian mangrove
357 system. *Estuarine, Coastal and Shelf Science* 31: 723–743.

358 Rodelli MR, Gearing JN, Gearing PJ, Marshall N, Sasekumar A. 1984. Stable isotope ratio as a tracer of mangrove carbon
359 in Malaysian ecosystems. *Oecologia* 61: 326–333.

360 Saenger P, Bellan MF. 1995. *The mangrove vegetation of the Atlantic Coast of Africa*. Toulouse (France): University of
361 Toulouse Press.

362 Saenger P, Hegerl EJ, Davie JDS. 1983. Global status of mangrove ecosystems. *The Environmentalist*, vol.3, 1983,
363 supplement no.3.

364 Said A, Smith MAK. 1997. *Proyek rehabilitasi dan pengelolaan mangrove di Sulawesi: ekonomi sumberdaya*. Laporan
365 Akhir. Direktorat Jenderal Reboisasi dan Rehabilitasi Lahan dan Asian Development Bank. Jakarta. [Indonesia]

366 Sasekumar A, Chong VC, Leh, MU, D’Cruz R. 1992. Mangroves as a habitat for fish prawns. *Hydrobiologia* 247: 195–207.

367 Spalding MD, Blasco F, Field CD. 1997. *World Mangrove Atlas*. Okinawa (Japan): International Society for Mangrove
368 Ecosystems.

369 Tam NFY, Wong YS. 1999. Mangrove soils in removing pollutants from municipal wastewater of different salinities.
370 *Journal of Environmental Quality* 28: 556–564.

371 Tews J, Brose U, Grimm V, Tielborger K, Wichmann MC, Schwager M, Jeltsch F. 2004. Animal species diversity
372 driven by habitat heterogeneity/ diversity: the importance of keystone structures. *J. Biogeogr.* 31: 79–92.

373 Twilley RR. 1998. Mangroves. Pages 445–473 in Messina MG, Conner WH (eds). *Southern Forested Wetlands: Ecology*
374 *and Management*. Boca Raton (FL): Lewis Publishers.

375 UNEP. 2014. *Importance of mangroves to people: a call to action*. United Nations Environment Programme World
376 Conservation monitoring Centre Cambridge.

377 Valiela I, Bowen JL, York JK. 2001. Mangrove forests: one of the world’s threatened major tropical environments.
378 *BioScience* 51(10): 807-815. <http://bioscience.oxfordjournals.org>.

Commented [AM27]: We don't find this reference in your text.
So please delete it.

- 379 van Lavieren LP. 1983. Wildlife management in the tropics, II. School of Environmental Conservation management,
380 Bogor.
- 381 Watson JEM, Whittaker RJ, Dawson TP. 2004. Avifaunal response to habitat fragmentation in the threatened littoral
382 forest of south-eastern Madagascar. *J. Biogeogr.* 31:1791–1807.
- 383 Woinarski JCZ, Fisher A, Brennan K, Morris I, Chatta R. 2001. Patterns of bird species richness and composition on
384 islands off Arnhem Land, Northern Territory, Australia. *Austral Ecol.* 26: 1–13.
- 385 Wolanski E, Spagnol S, Thomas S, Moore K, Alongi DM, Trott L, Davidson A. 2000. Modelling and visualizing the fate
386 of shrimp pond effluent in a mangrove-fringed tidal creek. *Estuarine, Coastal and Shelf Science* 50: 85–97.
- 387

Certificate of Appreciation

Granted with thanks to:

Abdul Malik

In recognition of his/her significant contribution as:

Peer Reviewer

of

Biodiversitas, Journal of Biological Diversity in 2016

Susoharta, Indonesia, 31st December 2016



Prof. Dr. Sutarna, M.Sc., Ph.D.
EDITOR-IN-CHIEF

[Contact Us](#)
[FAQ](#)
[Useful links](#)
[Home](#)
[Content and Archives](#)
[Aims and Scopes](#)
[Editorial Board](#)
[Guidance for Authors](#)
[Ethical Guidelines](#)
[Charges](#)
[Membership](#)
[Previously Reviewers](#)
[Conference Events](#)

LIST OF PREVIOUSLY REVIEWERS

2017 (January-June)

- **Abraham S. Khouw**, Department of Aquatic Resource Management, Faculty of Fishery and Marine Science, Universitas Pattimura, Ambon, Indonesia
- **Achmad Dinoto**, Research Center for Biology, Indonesian Institute of Sciences, Cibinong, Bogor 16911, West Java, Indonesia
- **Adel Ali Saeed Abduh Algeethi**, Universiti Tun Hussein Onn Malaysia, 86400 Parit Raja, Johor, Malaysia
- **Agus Nuryanto**, Faculty of Biology, Universitas Jenderal Soedirman, Purwokerto, Banyumas 53123, Central Java, Indonesia
- **Ahmad D. Setyawan**, Department of Biology, Faculty of Mathematics and Natural Sciences, Universitas Sebelas Maret, Surakarta 57126, Central Java, Indonesia
- **Ajay Kumar Gautam**, Faculty of Agriculture, Abhilashi University, Mandi, Himachal Pradesh-175045, India
- **Akhlaq Husain**, Vibhuti Khand, Gomti Nagar, Lucknow, Uttar Pradesh 226010, India
- **Akira Kikuchi**, Department of Biology Faculty of Mathematics and Natural Sciences, Universitas Brawijaya, Malang, 65145, Indonesia
- **Alberto Cuesta**, Department of Cell Biology and Histology, Faculty of Biology, University of Murcia, 30100 Murcia, Spain
- **Ali Akbar Hedayati**, Gorgan University of Agricultural Sciences and Natural Resources, Gorgan, Iran
- **Ali Sadough Niri**, Department of Fisheries, Faculty of Marine Sciences, Chabahar Maritime University, Chabahar, Iran
- **Amal Kumar Mondal**, Department of Botany and Forestry, Vidyasagar University, Midnapore-721-102, West Bengal, India
- **Andrea Pereira, Silveira**, Faculdade de Educacao de Itapipoca, Universidade Estadual do Ceara, CEP 60.500-000, Itapipoca - CE, Brasil
- **Andreas Otterbeck**, Marine Biology (MB), Department of Biosciences, University of Oslo, NO-0316 Oslo, Norway
- **Andrew Baird**, ARC Centre of Excellence for Coral Reef Studies, James Cook University, Townsville, Qld, Australia
- **Anna Cristina Lanna**, Fisiologia/Bioquimica Vegetal, Embrapa Arroz e Feijao, Empresa Brasileira de Pesquisa Agropecuaria (Embrapa), Santo Antonio de Goias, GO 75375-000, Brasil
- **Anne Russon**, Department of Psychology, Glendon College, York University, Toronto, Canada
- **Antonio Maria Luis Crespi**, UTAD Botanical Garden, CITAB, Vila Real, Portugal
- **Arshad Naji Alhasnawi**, School of Biosciences and Biotechnology, Faculty of Science and Biotechnology, Universiti Kebangsaan Malaysia, 43600 Bangi, Selangor, Malaysia
- **Ashok Shukla**, ICAR-Central Agroforestry Research Institute, Jhansi 284 003, Uttar Pradesh, India
- **Atsuko Takano**, Museum of Nature and Human Activities, Hyogo

- **Zhenghong Sui**, College of Marine Life Sciences, Ocean University of China, Qingdao 266003, China

2016 (July-December)

- **Aaron Moises Santana-Cordero**, Grupo de Investigacion Geografia Fisica y Medio Ambiente, IOCAG, Universidad de Las Palmas de Gran Canaria (ULPGC), 35001 Las Palmas de Gran Canaria, Las Palmas, Spanyol
- **Abdul Malik**, Department of Geography, Universitas Negeri Makassar, Kampus Parangtambung Makassar 90224, South Sulawesi, Indonesia
- **Abdullah Jaradat**, United States Department of Agriculture-Agricultural Research Service (USDA-ARS), The North Central Soil Conservation Research Laboratory, 803 Iowa Avenue Morris, MN 56267, USA
- **Abu Bakar Tawali**, Faculty of Agriculture, Universitas Hasanuddin, Makassar 90245, South Sulawesi, Indonesia
- **Abu Saleh Md. Golam Kibria**, Laboratory of Global Forest Environmental Studies, Department of Global Agricultural Sciences, Graduate School of Global Agriculture and Life Sciences, University of Tokyo, Bunkyo, Tokyo 1138657, Japan
- **Agus Sutanto**, Indonesian Tropical Fruit Research Institute, Solok 27301, West Sumatra, Indonesia
- **Agustin Sanguinetti**, Lab. de Biología del Desarrollo de las Plantas. Departamento de Biodiversidad y Biología Experimental. IBBEA, CONICET-UBA. Facultad de Ciencias Exactas y Naturales-Universidad de Buenos Aires. Lab. 17, 4to piso, Pabellon II, Ciudad Universitaria-C1428EHA. Ciudad Autonoma de Buenos Aires, Argentina
- **Ahmad D. Setyawan**, Department of Biology, Faculty of Mathematics and Natural Sciences, Universitas Sebelas Maret, Surakarta 57126, Central Java, Indonesia
- **Akari Watanabe**, Department of Oral Health Care and Rehabilitation, Subdivision of Oral Health and Welfare, Institute of Health Biosciences, University of Tokushima Graduate School, Tokushima 770-8504, Japan
- **Alain C. Tsobeng**, World Agroforestry Centre (ICRAF), Yaounde, Cameroon & World Agroforestry Centre, Nairobi, Kenya
- **Alejandro Casas**, Instituto de Investigaciones en Ecosistemas y Sustentabilidad, Universidad Nacional Autonoma de Mexico, Michoacan 58190, Mexico
- **Alejandro Martin Lopez**, Instituto de Ciencias Antropologicas, Universidad de Buenos Aires & Consejo Nacional de Investigaciones Cientificas y Tecnicas, Argentina
- **Alessandra La Notte**, University of Torino, 10153 Torino, Italy
- **Ali Mahmoud Muddathir**, Department of Horticulture, Faculty of Agriculture, University of Khartoum, Shambat 13314, Khartoum, Sudan
- **Amalia Moreno**, Department of Dental Materials and Prosthodontics Aracatuba Dental School, Universidade Estadual Paulista, Aracatuba, SP, Brazil
- **Ana Beatriz Pierri-Daunt**, Institute of Geosciences and Mathematical Sciences, UNESP - Sao Paulo State University, Rio Claro Campus, Department of Geography, Ecosystem Dynamic Observatory, Rio Claro, SP, Brazil
- **Ana Belen Guerrero Hinojosa**, Departamento de Ciencias de la Vida y de la Agricultura, Universidad de las Fuerzas Armadas, Sangolqui, Ecuador

Dynamics of mangrove community in revegetation area of Karangsong, north coast of Indramayu District, West Java, Indonesia

HENDRA GUNAWAN*, SUGIARTI, SOFIAN ISKANDAR

Forest Research and Development Center, FORDA, Ministry of Environment and Forestry, Jl. Gunung Batu No. 5, Bogor 16118, P.O. Box 165, West Java, Indonesia. Tel. +62-251-8633234, 7520067; Fax. +62-251-8638111; *email: hendragunawan1964@yahoo.com

Manuscript received: 15 July 2016. Revision accepted: 29 March 2017.

Abstract. Gunawan H, Sugiarti, Iskandar S. 2017. *Dynamics of mangrove community in revegetation area of Karangsong, north coast of Indramayu District, West Java, Indonesia. Biodiversitas 18: 659-665.* Mangrove along the north coast of Java is heavily degraded due to the conversion of land into fish ponds and human settlement areas. A revegetation program has been initiated by the local community of Karangsong Village, Indramayu District, West Java, Indonesia, supported by PT. Pertamina RU VI Indramayu. Our research aimed to study the population dynamics of the mangrove revegetation in Karangsong. Secondary data was collected from the Fishery and Maritime Services of Indramayu and PT. Pertamina. We carried out on-location field observations and we interviewed key respondents. Data was analyzed to describe trends in the diversity index and population dynamics of the mangrove. It was observed that the revegetation effort in the shoreline of Karangsong had covered ± 69.08 hectares which consisting of six species of mangrove and three tree species of coastal vegetation i.e. *Rhizophora mucronata* Lam., *Rhizophora stylosa* Griff., *Rhizophora apiculata* Blume, *Avicennia marina* (Forssk.) Vierh., *Avicennia alba* Blume, *Sonneratia caseolaris* (L.) Engl, *Terminalia catappa* L., *Casuarina equisetifolia* L., and *Ziziphus mauritiana* Lam. The mangrove population increased dramatically, from estimated 25,000 individuals in 2008 to 690,835 individuals in 2016. *Rhizophora mucronata* was the most dominant species (68.85%), followed by *Rhizophora stylosa* (18.33%) and *Rhizophora apiculata* (9.53%). The Shannon diversity index was fluctuated but tend to be increased from 0.80 to 0.95.

Keywords: Karangsong, mangrove, north coast, re-vegetation

INTRODUCTION

There are roughly 166,876 km² of mangrove along the shorelines of the world, with the largest proportion of mangrove occurring in Asia (77,169 km²) and the Americas (43,161 km²) (Valiela et al. 2001). Countries with the largest area of mangroves are Indonesia (4.25 x 10⁴ km²) (Spalding et al. 1997), followed by Brazil (1.34 x 10⁴ km²) (Spalding et al. 1997), Nigeria (1.05 x 10⁴ km²) (Saenger and Bellan 1995), and Australia (1.00 x 10⁴ km²) (Robertson and Duke 1990).

Globally, the area of mangrove area is declining rapidly as it is cleared and converted to mariculture, agriculture, urban development, logged timber concessions, and fuel production areas (Fortes 1988; Marshall 1994; Primavera 1995; Twilley 1998; Polidoro et al. 2010). At least 35% of the world's mangrove forest area has been lost in the past two decades (Valiela et al. 2001). It is apparent that maricultural practices are responsible for the bulk of the increasing loss of mangrove worldwide. For example, pond culture has been reported to be responsible for 50-80% of the loss of mangrove in Southeast Asia (Wolanski et al. 2000). Most of the damage is attributable to the direct loss of habitat from the conversion of "cheap" mangrove land to "valuable" shrimp, prawn, and fish ponds (Valiela et al. 2001).

In 1999, Indonesia's mangrove forest covered 8.6 million hectares which consisted of 3.8 million hectares of forest area and 4.8 million hectares of nonforest area.

Degradation of mangrove in forest area is 1.7 million hectares (44.73%) and in nonforest area is 4.2 million hectares (87.50%) (Gunawan and Anwar 2005). Indonesia has lost 40% of its mangrove in the last three decades (FAO 2007). The deforestation rate for mangrove in Indonesia is estimated to be 6% or 0.05 million hectares of the total annual forest loss (Margono et al 2014; Ministry of Forestry Republic of Indonesia 2014). The Ministry of Forestry has reported that only 31% of the remaining mangrove is in an intact condition and the rest (69%) is heavily degraded (Ministry of Forestry 2007). FAO (2007) reported that mangrove forest in Indonesia is 3,062,300 hectares or 19% of the world's mangrove and still the largest in the world, followed by Australia and Brazil.

Mangrove forest in Java Island is decreasing as the impact of conversion to mariculture, human settlement and other uses worsens. This impact is due to limited understanding and awareness by surrounding communities of the ecological importance of mangrove and uncertainty about land status (Said and Smith, 1997). In 2011, mangrove in West Java Province was estimated as covering 40,130 hectares which were distributed between forest area 32,314 ha (80.52%) and non-forest area, 7,816 ha (19.48%), including 13 regencies (Forestry Service of West Java Province 2013). The degraded mangrove in this province is 15,276 hectares (38.06%), with the largest occurring in Karawang District 13,181ha (32.85%) followed by Bekasi 10,481ha, Indramayu 8,720ha, Subang 7,346 ha, Cirebon 190 ha, Ciamis 170 ha, Garut 32 ha