

PAPER NAME

Combined effects of nitrification inhibitor and zeolite on greenhouse gas fluxes and corn growth.pdf

WORD COUNT

652 Words

CHARACTER COUNT

3481 Characters

PAGE COUNT

1 Page

FILE SIZE

111.5KB

SUBMISSION DATE

May 5, 2023 10:49 PM GMT+8

REPORT DATE

May 5, 2023 10:50 PM GMT+8

● **11% Overall Similarity**

The combined total of all matches, including overlapping sources, for each database.

- 9% Internet database
- 9% Publications database
- Crossref database
- Crossref Posted Content database
- 6% Submitted Works database

● **Excluded from Similarity Report**

- Quoted material
- Cited material
- Small Matches (Less than 10 words)
- Manually excluded sources



Combined effects of nitrification inhibitor and zeolite on greenhouse gas fluxes and corn growth

Oslan Jumadi¹ · Yusminah Hala¹ · R. Neni Iriany² · Andi Takdir Makkulawu² · Junja Baba³ · Hartono¹ · St. Fatmah Hiola¹ · Kazuyuki Inubushi³

Received: 11 September 2017 / Accepted: 15 October 2019 / Published online: 26 November 2019
© Springer-Verlag GmbH Germany, part of Springer Nature 2019

Abstract

Field and incubation experiments were conducted to determine the emission rate of greenhouse gases, nitrogen change, populations of AOB, NOB, and fungi as well as growth of corn in response to amendment of urea granulated with and without nitrification inhibitors and zeolite. The application of urea with neem, urea with zeolite, urea with zeolite + neem, urea with zeolite + dicyandiamide, and urea with dicyandiamide (UD) decreased the N₂O emissions by 16.3%, 59.6%, 66.8%, 81.9%, 16.3%, and 86.7%, respectively. Meanwhile, patterns of CH₄ fluxes were mostly determined by small emissions. Increase in corn height, weight of cobs, biomass, and chlorophyll leaf contents were not significantly different between urea alone and urea with NIs and zeolite. In the incubation experiment, the highest concentration of NH₄⁺ and N₂O production was detected during the first week and it remained high up to the second week of incubation in the combination of urea with NIs and zeolite treatments, although there was no significant difference compared with urea. During NH₄⁺ decrease, the concentration of NO₃⁻ started to accumulate from the second to the third weeks. Production of CO₂ showed no significant differences among treatments. The static production of CO₂ could also explain that NIs and zeolite additions did not change AOB, NOB, and fungi activities after the fourth week of incubation.

Keywords Emission of N₂O and CH₄ · CO₂ Production · Dicyandiamide · Neem · Nitrification inhibitor · Zeolite

Introduction

Urea (CO(NH₂)₂) has been widely used by farmers as a major source of nitrogen fertilizer to support corn production which is the second most important cereal after rice in Indonesia. The worldwide demand for urea was forecasted to increase from 148 Mt in 2010 to 171.7 Mt in 2015, representing a growth of 3.2% per annum (IFA 2011). Nitrogen (N) is more substantial in a plant and needs larger quantities than other nutrients and it is estimated that only 30–40% of the application of N fertilizer

is taken up by the crop. Most of it disappears by ammonia volatilization, nitrification, and denitrification. Therefore, the use of urea in the agriculture sector in order to increase the quantity and quality of agricultural food production can generate a negative impact on the environment, such as ozone layer depletion due to enhancement of greenhouse gas emissions, particularly nitrous oxide (N₂O) gas (Mosier and Kroeze 2000).

Emission of N₂O gas in agricultural land is determined by the nitrification process in aerobic soil conditions and formed nitrate (NO₃⁻) that has mobility capacity to leach as a pollutant to the environment. The NO₃⁻ is susceptible to denitrification loss in anaerobic conditions in soil or water. In addition, that process is also a cause of low use nitrogen fertilizer efficiency in the agricultural sector (Mosier and Kroeze 2000). The process of release of N₂O from the soil into the atmosphere is influenced by diffusion processes in the soil and the capacity of soil to consume N₂O, which is determined by several factors such as the production footprint in the soil, soil organic matter, soil texture, and soil water content (Zhang et al. 2017; Majumdar et al. 2002; Jumadi et al. 2014).

Responsible editor: Philippe Garrigues

✉ Oslan Jumadi
oslanj@unm.ac.id

¹ Biology Department, Faculty of Mathematics and Natural Sciences, Universitas Negeri Makassar, Makassar 90224, Indonesia

² Plant Breeding, Cereals Research Institute, Maros 90514, Indonesia

³ Graduate School of Horticulture, Chiba University, Matsudo, Chiba 271-8510, Japan

● **11% Overall Similarity**

Top sources found in the following databases:

- 9% Internet database
- 9% Publications database
- Crossref database
- Crossref Posted Content database
- 6% Submitted Works database

TOP SOURCES

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

1	repositorio.unb.br Internet	3%
2	Yasser Djawad, Hamza Rehman, Oslan Jumadi, Muhammad Tufail, Sha... Crossref	2%
3	intechopen.com Internet	2%
4	A R B Sofia, Y Hala, A T Makkulawu, S F Hiola, H Karim, R N Iriany, R Sj... Crossref	2%
5	repository.unimus.ac.id Internet	2%

● Excluded from Similarity Report

- Quoted material
- Small Matches (Less than 10 words)
- Cited material
- Manually excluded sources

EXCLUDED SOURCES

Oslan Jumadi, Yusminah Hala, R. Neni Iriany, Andi Takdir Makkulawu, Junja B...	70%
Crossref	
link.springer.com	64%
Internet	
researchgate.net	43%
Internet	
pubfacts.com	39%
Internet	
essentials.ebsco.com	36%
Internet	
pubag.nal.usda.gov	16%
Internet	
eprints.unm.ac.id	9%
Internet	
scribd.com	7%
Internet	
baixardoc.com	7%
Internet	
digilib.unila.ac.id	7%
Internet	

iieta.org

Internet

5%

iieta.org

Internet

5%