



FACULTY OF MATHEMATICS AND NATURAL SCIENCES
UNIVERSITAS NEGERI MAKASSAR
Makassar, Indonesia

PROCEEDING

*"The Role of Mathematics, Sciences, Technology,
Education towards ASEAN Economic Community and Global Challenges"*

ICMSTEA

ISBN: 978-602-99837-4-6



PROCEEDING

2nd ICMSTEA

INTERNATIONAL CONFERENCE ON MATHEMATICS,
SCIENCE, TECHNOLOGY, EDUCATION
AND THEIR APPLICATIONS

*"The Role of Mathematics, Sciences, Technology,
Education towards ASEAN Economic Community and Global Challenges"*

October 3rd - 4th, 2016
Makassar, South Sulawesi, Indonesia

FACULTY OF MATHEMATICS AND NATURAL SCIENCES
UNIVERSITAS NEGERI MAKASSAR

PROCEEDING

ICMSTEA 2016 :
The Role of Mathematics, Sciences, Technology, and Education Towards ASEAN
Economic Community and Global Challenges

Makassar, South Sulawesi, Indonesia
3rd – 4th October 2016

Fakultas Matematika dan Ilmu Pengetahuan Alam
Universitas Negeri Makassar
2016

ICMSTEA 2016: THE ROLE OF MATHEMATICS, SCIENCES, TECHNOLOGY, AND
EDUCATION TOWARDS ASEAN ECONOMIC COMMUNITY AND
GLOBAL CHALLENGES

- Editor** : Dr. Syafruddin Side
Dr. Ir. Wiharto Caronge, M.Si.
Uca, S.Si., M.Si., Ph.D.
Iwan Dini, S.Si., M.Si.
Ansari Saleh Ahmar, S.Si, M.Sc.
Bustang, S.Pd., M.Sc.
Muh. Husnul Khuluq, S.Pd., M.Sc.
Said Fachry Assagaf, S.Pd., M.Sc.
Sahlan Sidjara, S.Si., M.Si.
Ahmad Ansar, S.Pd., M.Sc.
Syamsuddin Mas'ud, S.Pd., M.Sc.
Muhammad Isbar, S.Si., M.Si.
Surahman, S.Pd., M.Pd.
- Editorial Assistant**
Layouter : Andi Ikhsan Maulana, S.Pd
Dimas Suharto
- Cover Administrator** : Nurul Kusuma Wardani, S.Pd.
- Reviewer Board** : Prof. Alfred Edward Szmidt, Ph.D.
Prof. Valerio Sbordoni, Ph.D.
Prof. Birgit Loch
Dr. Richard Randriantoamanana
Prof. Fahrul Zaman bin Huyop, Ph.D.
Prof. Mohd. Ekhwan Toriman, Ph.D.
Prof. Eko Hadi Sujiono, M.Si.
Prof. Oslan Jumadi, S.Si., M.Phil., Ph.D
Muh. Abdy, S.Si, M.Si., Ph.D
Prof. Dr. Jasruddin M, M.Si.
Prof. Dr. Muris, M.Si.
Prof. Dr. Ir. Yusminah Hala, M.Si.
Prof. Dr. Usman Mulbar, M.Pd.
Prof. Dr. Muh. Danial, M.Pd.

Copyright © October 2016
ISBN 978-602-99837-4-6

Diterbitkan oleh : Fakultas MIPA Universitas Negeri Makassar

WELCOME SPEECH

Forewords from the Chairman of Committee

Assalamu'alaykum wa Rahmatullahi wa Barakatuh

First of all, I wish to extend a warm welcome to fellow delegates from the various countries and regions. I realize that you are fully dedicated to the sessions that will follow but I do hope you will also take time to enjoy sparkling Makassar with its tropical setting, friendly people, and wonderful cuisine.

This 2nd International Conference on Mathematics, Science, Technology, Education, and Their Application 2016 is organized by the Faculty of Mathematics and Natural Sciences, Universitas Negeri Makassar to bring all experts and researches in these field sharing their important thoughts and findings. The conference will be held in two days from 3rd – 4th of October 2016 with two keynote speakers, seven invited speakers, and more than 80 parallel speakers from different backgrounds.

Let me take this opportunity, on behalf of the committee members, to express my gratitude and sincere thanks to the keynote, invited and all parallel speakers for spending their valuable time with us in this conference. I do hope that your time in Makassar will be valuable and memorable.

Finally, I would like to thank all steering and organizing committee for their hard work and dedication to the success of this conference. I would like also to apologize to all of you should you find any inconvenience during this event.

Thank you very much,

Wassalamu'alaykum Warahmatullahi Wa barakatuh

Chairman of Committee,

Dr. Drs. A. Mushawwir Tayeb, M.Kes.

The 2nd ICMSTEA Speech
By The Dean of Mathematics and Natural Sciences Faculty
Universitas Negeri Makassar

Your excellency Rector of Universitas Negeri Makassar
 Honorable Vice Rectors and Dean of All Faculties
 Honorable Keynote Speakers
 Distinguished all invited speakers from outstanding universities
 Distinguished all speakers and guests
 All participants,
 Ladies & Gentlement,

Assalamu'alaykum Warahmatullahi Wabarakatuh. My greetings for all of you. May peace and God's blessing be upon us all. *Alhamdulillah*, all praises be to the Almighty God, Allah *subhanahu wata'ala*.

It is my pleasure to welcome you all to the opening of The 2nd International Conference on Mathematics, Science, Technology, Education & their Applications (2nd ICMSTEA). I am delighted to see that the Mathematics and Natural Science Faculty has again organized the second conference that capitalize on our strength and built on our commitment to promoting Mathematics, Science, Technology and Education.

I do hope that this conference would bring a great opportunity for all of us to strengthen our contribution to the advancement of our nation.

I would like to take this opportunity to thank the conference organizing committee for their diligent work. I would also like to thank participants, especially those of you coming from abroad, for joining us and sharing your valuable experiences. Should you find any inconveniences and shortcomings, please accept our sincere apologies.

Finally, let me wish you fruitful discussion and a very pleasant stay in Makassar.

Thank you,
 Wassalamu'alaykum Warahmatullahi Wabarakatuh

Dean of Faculty of Mathematics and Natural Sciences
 Universitas Negeri Makassar

Prof. Dr. Abdul Rahman, M.Pd.

TABLE OF CONTENTS

Cover	i
Welcome Speech	iii
Table of Contents	v
Invited Speakers	1
Genetic Structure and Evolutionary History of a Diploid Hybrid Pine <i>Pinus densata</i> Inferred from the Nucleotide Variation at Seven Gene Loci <i>Alfred Edward Szmidt</i>	1 – 8
Biodiversity and Big Data In A Changing World: From Museum Specimens to Citizen Sciences <i>Valerio Sbordon</i>	9 – 15
Engaging Students with Tablet Technology in Science, Technology, and Mathematics Education <i>Birgit Loch</i>	16 – 38
Assesment Within Elpsa Lesson Design <i>Sitti Maesuri Patahuddin</i>	39 – 48
Why High Performance Computing (HPC) Matters? <i>Richard Randriantoamanana</i>	49 – 58
In Silico Study of L-specific dehalogenase from Rhizobial Species Strain RC1 <i>Fahrul Zaman bin Huyop</i>	59 – 68
Fostering Water Innovation: Securing the ASEAN Water Future <i>Mohd. Ekhwan Toriman</i>	69 – 78
Basic Research (in Physics) for Development of Nation Innovation <i>Eko Hadi Sujiono</i>	79 – 86
Parallel Speakers	
The Influence Of <i>Problem Based Learning Model</i> Toward Motivation and Physics Learning Outcomes Of Students SMA Negeri 1 Parangloe Gowa Regency <i>Muhajirin, Sidin Ali & Pariabti Palloan</i>	87 – 95
Analysis of Student’s Conceptual Understanding of Mathematics on Set at Class VII SMP Frater Palopo <i>Muhammad Ilyas & Fahrul Basir</i>	96 – 102

Pah Characteristics In Sediment Around Makassar Coast Using GC-FID <i>Muhammad Syahrir, Nurul Hidayat Aprilita, Nuryono, & Netti Herawati</i>	103 – 114
Perception of 21st Century Learners to The E-Books as A Learning Resources <i>Muhammad Takwin Machmud</i>	115 – 122
The Group Of Understory Herbaceous Vegetation Stand At Tropical Lower Mountain Forest Of Mount Salak, West Java <i>Muhammad Wiharto, Cecep Kusmana, Lilik Budi Prasetyo, Tukirin Partomiharjo, Hamka L, & Abd. Muis</i>	123 – 129
Antibacterial Compounds Characterization in Chloroform Extract Leaves of Tahiyam Plant (<i>Lantana Camara</i> Liin.) <i>Muharram, St. Nurzulaiha, Nurrahmania, Iwan Dini, Pince Salempa, & Maryono</i>	130 – 135
Effect of Mixture Inoculum of Lactic Acid Bacteria (LAB) and Mold Amylolytic In Various Concentration And Fermentation Time of Changing Protein And HCN Content Of Bitter Cassava Roots (<i>Manihot aipi</i> Phol.) <i>Nurhayani H. Muhiddin, Nur Arfa Yanti, & Hasanah</i>	136 – 143
The Analysis of Total Cholesterol Levels in Mice (<i>Mus musculus</i>) MALES who were Given Extracts of Methanol Leaf <i>Cemba</i> (<i>Acacia pennata</i>) <i>Nurul Muhliah, Irma Suryani Idris, Andi Mu'nisa, & Ernawati</i>	144 – 150
Plants in a Square: Explore Plants Description With QR Code Feature <i>Khalisha Azis, Sitti Saenab & Syamsiah</i>	151 – 155
Exploring the Correlation between Metacognitive Skills and Retention of Students in Different Learning Strategies in Biology Classroom <i>Arsad Bahri</i>	156 – 161
Physical Fitness For Futsal Referee Of Football Association In Thailand <i>Acting Sub Lt. Thaweesub Koeipakvaen</i>	162 – 170
Histological study of the respiratory system of Sulawesi Medaka fish (<i>Oryzias celebensis</i>): as a candidate of animal model <i>Dwi Kesuma Sari, Irma Andriani, & Khusnul Yakin</i>	171 – 176
Technological Innovation of Learning Mobile-Based Learning <i>Herlinah & Baso Habibi</i>	177 – 184
Evaluation of Antimicrobial Activity and Phytochemical Screening of Chloroform Extract of <i>Usnea</i> sp. <i>Iwan Dini, Maryono, Nurul Utami, Sitti Hajar, & Ahmad Hadani</i>	185 – 188
The effectiveness of Scientific-Inquiry Learning Model to Improve Scientific Thinking Skills of Grade X student of High School in Gowa Regency <i>Jusniar & Sumiati Side</i>	189 – 194

Pedagogical Content Knowledge: Teacher's Knowledge Of Students In Learning Mathematics On Limit Of Function Material <i>Ma'rufi, I Ketut Budayasa, & Dwi Juniati</i>	195 – 199
Simulation Infiltration Rate of Water on Sand Media by Finite Difference Method <i>Masitah Osman, Nasrul Ihsan, & Muhammad Arsyad</i>	200 – 210
The Influence Of Problem Based Learning Model Toward Motivation and Physics Learning Outcomes Of Students SMA Negeri 1 Parangloe Gowa Regency <i>Muhajirin, Sidin Ali & Pariabti Palloan</i>	211 – 218
The Influence of Experiment Methods on Science Process Skill and Cognitive Learning Outcomes of the X Grade Students of MIA SMAN 1 Soppeng Riaja 2015/2016 <i>A. Sri Sofializa, Muhammad Arsyad, & Muris</i>	219 – 226
The Influence Of Problem Based Learning Model Toward Motivation and Physics Learning Outcomes Of Students SMA Negeri 1 Parangloe Gowa Regency <i>Muhajirin, Sidin Ali & Pariabti Palloan</i>	227 – 233
The Development of Critical Thinking Inventory Instrument for Biology Department Students <i>Abd Muis, Adnan Gassing, Ismail Laumma & Nurjannah</i>	234 – 239
Unleash Students' Motivation with Blended Knowledge Transfer Instructional Model <i>Adnan, Sitti Saenab, & Andi Rahmat Saleh</i>	240 – 243
The Development of Student's Worksheet of Physics Based on Virtual Simulation and Its Influence on Physics Learning Outcomes of Students <i>Ahmad Swandi & Bunga Dara Amin</i>	244 – 251
Teaching Material Development with Challenge Based Learning (CBL) Basis to Improve Critical Thinking Ability on Human Reproduction System Material of Class XI IPA 4 Students at MAN Pinrang <i>Andi Asmawati Azis, Jasruddin, & Reni</i>	252 – 258
Implementation of Accreditation Policy and Quality of Public Primary School Education Service in South-Sulawesi <i>Andi Cudai Nur & Sumarlin Mus</i>	259 – 262
Plants in A Square: Explore Plants Description with QR Code Feature <i>Khalisha Azis, Wiharto, Siti Saenab, & Syamsiah</i>	263 – 272
Fostering Science Learning Quality in Frame of Ecosystem Topic through Lesson Study <i>Besse Aisyah, Nurfatima, Rina Kurnia, Andi Asmawati Aziz, & Andi Nurul Virninda Yusuf</i>	273 – 275
Fractionation Ethyl Acetate Extract of Stem Bark Soursop(<i>a. Muricata</i> . Linn) Potential Anticancer <i>Pince Salempa, Muharram, & Iwan Dini</i>	276 – 279

Entrepreneurship Education Development In Dealing Asean Economic Community <i>Mohammad Rakib</i>	280 – 285
The Development of Basic Chemistry Courses Program Based Problem Solving to Improve Student's Critical Thinking Skills <i>Ramlawati & Ratnawaty Mamin</i>	286 – 291
Increase The Production and Quality of Biogas from Waste of Cattle Rumen Content Through The Addition of Molasses and Zeolite <i>Ramli, Satria Aly, & Hartono</i>	292 – 296
Marica Goat's Response To The Provision of Superior Feed <i>Rosdiana Ngitung</i>	297 – 301
Students' Mathematics Achievement and Its Relationship with Parents' Education Level, and Socio-Economic Status In Turkey <i>Rusli</i>	302 – 308
The Abstraction Ability in Constructing Relation Within Triangles By The Seventh Grade Students of Junior High School <i>Sitti Mutmainna Hasma, Suwardi Annas, & Djadir</i>	309 – 313
Water Content Influence to Electrical Properties From Soil Volcanic <i>Sulistiawaty</i>	314 – 320
The Influence Of Using Destination Card Through Team Games Tournament (TGT) Type Of Cooperative Learning Toward Student's XII Grade Achievement at SMA Negeri 01 Unggulan Kamanre, Luwu Regency <i>Ulva Nilawaty Sudarmadi</i>	321 – 324
Applying SARAC Approach and The Effect in Learning Mathematics For Students Grade VIII <i>Usman Mulbar & Nasrullah</i>	325 – 331
Analysis Von Bertalanffy Equation With Variation Coefficient <i>Usman Pagalay, Budiawan, & Anisyah</i>	332 – 338
Results Increasing Student Learning Through The Use of Biology Learning Model Cooperative Think Pair Share (TPS) The Student Class XI IPA SMA Negeri 5 Makassar <i>Wiwik Wiji Astuti, Ismail & Musyafar</i>	339 – 346
Characterization of crude chitinase produced by <i>Trichoderma virens</i> in solid state fermentation <i>Rachmawaty & Madihah</i>	347 – 354

The Group Of Understory Herbaceous Vegetation Stand At Tropical Lower Mountain Forest Of Mount Salak, West Java

Muhammad Wiharto^{1*}, Cecep Kusmana², Lilik Budi Prasetyo²,

Tukirin Partomiharjo³, Hamka L¹, Abd. Muis¹

1. Dept. of Biology, State University of Makassar.

2. Bogor Agricultural University.

3. Research Center for Biology - Indonesian Institute of Sciences.

*Correspondence should be addressed to: Muhammad Wiharto; wiharto09@gmail.com; Tel: 08218846675

Abstract. This study aimed to determine the groups of herbaceous vegetation stands at tropical lower mountain forest of Mount Salak, Bogor, West Java. Hierarchical cluster analysis method was used for grouping herbaceous vegetation stands by Importance Value Index. The distance between stands was determined using Sørensen index while the grouping of stands made by the farthest distance method. The difference in stand's vegetation structure among the groups was tested with Kruskal-Wallis statistics. There were 4 stand groups (SG) formed at 80% distance of Sørensen index. The difference between SG was most visible through floristic composition. The most difference of SG was between SG 2 and SG 4, where the most dominant species that make up these SGs were *Nephrolepis exaltata*, *Isachne globosa* (family: Poaceae), and *Etilingera megalochelios* (family: Zingiberaceae) for SG 2 and *E. megalochelios*, *N. exaltata* and *Dinochloa scandens* (family: Poaceae) for SG 4. Species *N. exaltata*, *Scleria purpurascens* (family: Cyperaceae), and *Selaginella plana* (family: Selaginellaceae) were found throughout the entire SG. There were no significant differences among the SG in the case of Shannon-Wiener's Diversity Index, Pielou's Evenness Index, Simpson's dominance index and the number of species ($P \geq 0.05$). There was a decreasing trend of vegetation structure's value from SG 2 to SG 1.

Key words: Groups of herbaceous vegetation stands, Importance Value Index, Mount Salak, Sørensen index, tropical lower mountain forest,

INTRODUCTION

Mount Salak is part of the Mount Halimun Salak National Park (MHSNP) based on the Minister of Forestry's decree No. 175 / Kpts-II / 2003 (Ekayani, *et al.*, 2014). The MHSNP is the largest national park and contains largest mountainous tropical rainforest ecosystem on Java island, rich in biodiversity, beauty of the landscape and diverse of traditional culture (GHSNPMP-JICA, 2009). Administratively, the MHSNP covers two provinces of West Java and Banten as well as three districts, namely Bogor, Sukabumi and Lebak (Galudra *et al.*, 2005).

The Mount Halimun Salak National Park is vulnerable because its location is very close to the residential area. According to Tapadebu *et al.*, (2008), Java has experienced very fast rate of natural ecosystem conversion and it also is the most populated area in Indonesia.

Forests are the association of plants and animals communities which are dominated by trees and woody vegetation with a certain area that forms the micro-climate and specific ecological conditions (Simon, 1993). The quality and quantity of forest ecosystems should be preserved, among others by conservation in which ecosystem utilization is carried out by considering the overall presence function. Forest management that only considers one function alone will lead to the destruction of forests (Ismaini *et al.*, 2015). Undergrowth is one part of the forest functions (Prihantoro, 2013).

Most research on tropical forests vegetation has tended to focus on the tree component, and very few toward the understory vegetation (Lü *et al.*, 2011). Herbaceous community in tropical rainforests is so little known, and it is less research on the structure quantitatively. On the other hand, underground herbaceous plants is a group of rich plant, comprising 14 to 40% of the species of the total species found in tropical rainforests. (Costa, 2004).

Understory vegetation layer is important for nutrient cycling (Hart & Chen, 2006), carbon cycling at an ecosystem (Larashati, 2010), and towards the plants of upper canopy layers (Hart & Chen, 2006). This layer is able to inhibit runoff on the forest floor to prevent erosion (Kinho, 2011), often used as indicators of soil fertility and litter producers in improving soil fertility (Hilwa & Masyrifina, 2015). One of the forest components that use by the society around the forest is an understory vegetation (Suharti, 2015), as food, medicinal plants, and as an alternative energy source (Hilwa & Masyrifina, 2015).

Studies on herbaceous vegetation understory, especially in lower mountain zone on Mount Salak still need to be increased, especially the ecology of understory vegetation considering the importance of this vegetation layer, beside that, it is also still rarely conducted. The comprehension on understory vegetation ecology is important for conservation activity of the MHSNP and also as a source of baseline data. This study aimed to determine the groups of herbaceous vegetation stands and how the floristic composition and structure of herbaceous vegetation on the group stands at tropical lower mountain forest of Mount Salak, Bogor, West Java.

METHODS

Study site

This research was conducted at the lower mountain forest of Mount Salak, at the MHSNP. Location of the study can be climbed from several places, and in this study through the village of Gunung Bunder Dua (S6°41'484 "-E106°42'234") and the village of Gunung Sari (Kawah Ratu) (S6°41'.786"-E106°42'.006") subdistrict of Pamijahan, Bogor Regency. The research area is located from 1033 m asl - 1362 m asl.

Based on soil maps of West Java, the soil type in this area consists of the brown andosol association and brown regosol, the brown latosol association and yellowish brown latosol, brown latosol reddish and brown latosol, the brown latosol reddish association and laterite, complex brown latosol reddish and lithosol, brown latosol association and gray regosols (LP Tanah, 1966 *in* Larashati, 2010). According to the Schmidt & Ferguson's classification (1951), the climate at the MHSNP region including type A, with an annual rainfall of 4000-6000 mm.

Butterfly at Mount Salak mostly came from the Nymphalidae family. Butterfly species with the highest number of individuals is *Melanitis leda* (Tapadebu *et al.*, 2008). Among others of Lucanid beetle that can be found in this mount are *Cyclommatus canaliculatus*, *Dorcus taurus*, *Prosopocoilus astocoides*, *Odontolabis bellicosa*, and *Prosopocoilus zebra* (Koneri *et al.*, 2010). The endemic bird of Sumatra, Java, and Bali, the Walik Purple Head (*Ptilinopus porphyreus*) can be found on Mount Salak (van Balen & Nijman, 2004).

Sampling Method

Sampling was carried out in four places, namely on slopes facing toward the North, South, East, and West so that ecologically the entire region could be represented. At each location, we put 15 transects. The transect was placed cut perpendicular to the direction of elevation topography. Each transect has a size of 200m x 5m, and we considered it as 1 stand. The total number of stands were 60.

We made 10 observation plots with each has a size of 5m x 5m at every transect and the distance between the plot was 15m. Herbaceous plants in this study is a plant that does not have woody body parts above the soil surface. Fern was included in this study if it has a height at maturity ≤ 50 cm. Data was obtained by estimating the percentage of the herbaceous plants crown cover in each observation plot. Assessments carried out with reference to the Braun-Blanquet scale (Barbour *et al.*, 1987).

Data Analysis

We determined species Importance Value Index (IVI) at each stand, where the IVI was obtained through the sum of the relative frequency with the relative dominance of each species (Cox, 1978; Hardjosuwarno, 1990; and Kusmana, 1997). We calculated the Shannon-Wiener's diversity index (Lalchhuanawma, 2008),

the Pielou's evenness index, species richness (Omoro & Luukkanen, 2011), and the Simpson's dominance index (Barbour *et al.*, 1987).

Data matrix of IVI was imported into R using `read.csv()` function, then transformed so that the column part of the data matrix became the line in order to be processed with the `vegan` library (Oksanen *et al.*, 2016). Hierarchical cluster analysis method was used for grouping herbaceous vegetation stands by IVI and calculated it using `hclust()` function. The distance between stands was determined using Sorensen index (Kent & Cooker, 1992), and calculated with `designdist()` function, which was part of the `vegan` library. Dendrogram graph is displayed to indicate the SG.

The most dominant species (tmds) in each SG was the species that have the most number of highest IVI of first and second order. If there are two or more species that have the highest IVI with the same number of stands, tmds will be decided based on the highest second ordered number of IVI from those species. The number of dominant species in SG were 3, where the first order of tmds was written first followed by the second and the third order of tmds in the next sequence.

Non-parametric test Kruskal-Wallis rank sum was conducted to determine whether there was a difference in vegetation structure among SGs (Lukiastuti & Hamdani, 2012). Data analysis and statistical tests performed by use the R version 3.2.2 (R Core Team, 2015).

RESULTS

Stand Groups

There were four SGs at 80% distance of the Sorensen's index. Every SG was composed with the different number of stands. The SG 1 was consisted with 17 stands which were stands 1, 4, 5, 17, 18, 19, 20, 24, 27, 30, 33, 40, 41, 46, 51, 52, and 60. The SG 2 consisted with 19 stands and was the highest number compared to other SG. This SG was made up of stands 2, 6, 8, 12, 13, 16, 21, 23, 31, 35, 36, 39, 45, 47, 49, 50, 54, 56, and 58.




Figure 1. The Stand groups of herbaceous vegetation at lower mountain zone of Mount Salak. Each stand groups was confined within a box. Number 1,2,3, and 4 indicating stand group. T: Stand. Tegakan: Stand.

The SG 3 consisted with 14 stands, which were stands 3, 7, 11, 14, 22, 25, 29, 34, 37, 38, 43, 48, 55, and 59. The SG 4 was composed of 10 stands, which were stands 9, 10, 15, 26, 28, 32, 42, 44, 53, and 57. This SG has the lowest number of stands among the other stands (Figure 1).

The composition of species in Stand Groups

The most dominant species on the SG 1 is: *Nephrolepis exaltata* (family: Lomariopsidaceae), *Etilingera megalochelios* (family: Zingiberaceae), and *Scleria purpurascens* (family: Cyperaceae); on SG 2 is: *N. exaltata*, *Isachne globosa* (family: Poaceae), and *E. megalochelios*; on SG 3 is *N. exaltata*, *E. megalochelios* and *I. globosa*, and in the SG 4 is *E. megalochelios*, *N. exaltata* and *Dinochloa scandens* (family: Poaceae). The *N. exaltata* was found as tmds in 16 stands (Table 1), in addition, the *N. exaltata*, *S. purpurascens*, and *Selaginella plana* (Family: Selaginellaceae) were found in all stands in the whole SGs.

Table 1. Species with highest Important Value Index

Stand Group	Rank I		Rank II		Rank III	
	Species	Number of stands	Species	Number of stands	Species	Number of stands
1	<i>Nephrolepis exaltata</i>	4	<i>Etilingera megalochelios</i>	3	<i>Scleria purpurascens</i>	2
2	<i>Nephrolepis exaltata</i>	6	<i>Isachne globosa</i>	4	<i>Etilingera megalochelios</i>	3
3	<i>Nephrolepis exaltata</i>	6	<i>Etilingera megalochelios</i>	3	<i>Isachne globosa</i>	2
4	<i>Etilingera megalochelios</i>	3	<i>Nephrolepis exaltata</i>	2	<i>Dinochloa scandens</i>	1

The most different of SG in terms of species with the highest IVI and the most was SG 4. In this group, *N. exaltata* was not the first order of tmds, but *E. megalochelios* and the third order of tmds was *D. scandens*. The first order of tmds in the other three groups was *N. exaltata*. The next group was SG 1, where of the third order of tmds was *S. purpurascens*, and this species was not tmds in the other SGs (Table 1).

At SG 2, *E. megalochelios* and *E. punicea* were the third order of tmds. Both species have the highest IVI where each in three stands, however *E. megalochelios* has more of the second order of tmds on more stands than *E. punicea*. The most dominant species on SG 2 and SG 3 were same but different in terms of the second and the third order of tmds, which for SG 2, those species were *I. globosa* and *E. megalochelios*, while for SG 3 was the opposite (Table 1).

The species that only found in SG 2 was *Phragmites karka* (family: Poacea) and *Mussaenda frondosa* (Family: Rubiaceae), while in SG 4 was *Dipteris conjugata* (family: dipteridaceae). At SG 1 and SG 3, there was no species that can only be found in those of SGs. The total number of species found in this study was 58. Species *N. exaltata* is a plant with fern life form, *E. megalochelios* is a plant with a ginger life form, *I. globosa* is a plant with grass life form, *S. purpurascens* is a plant with a grass life form and *D. scandens* is a plant with a climbing bamboo life form.

Species Diversity structure in stand groups

The mean range number of herbaceous vegetation species on each SG was 24.41 at SG 1 to 25.316 at SG 2 (Figure 1). The Kruskal-Wallis rank sum test showed that there was no different number of species in each SG (Kruskal-Wallis $\chi^2 = 0.66999$, df = 3, P = 0.8802).

Shannon-Wiener's diversity index on each SG has a mean value that ranges from 2,910 at SG 1 which was the lowest value, while the other value was 2,959 at SG 4 which is the highest value (Figure 1). There were no differences of Shannon Weiner's diversity index among the four SGs (Kruskal-Wallis $\chi^2 = 0.226$, df = 3, p = 0.973).

Pielou's evenness index in four SGs has a mean value that ranges from 0,910 at SG 1 which was the lowest value and 0.927 at SG 4 which was the highest value (Figure 1). The Pielou's evenness index among the four SG has no different (Kruskal-Wallis $\chi^2 = 1,652$, df = 3, P = 0.6477). This species evenness values was closed to 1, which indicated that each species composing herbaceous vegetation in each SG has relatively equal of the individual number.




Figure 2. Mean and Standard Deviation of (a) Shannon Wiener's Diversity Index, (b) Number of Species, (c) Pielou's Evenness index, and (d) Simpson's index on four SG herbaceous vegetation. The y axis represents mean and standard deviation. Note: SG: Stand group.

The mean value of dominance species based on Simpson's index range from 0.063 at SG 4, which was the lowest value to 0,070 at SG 1, which was the highest value (Figure 1). There were no difference of Simpson's Index among the SGs (Kruskal-Wallis $\chi^2 = 0.29953$, $df = 3$, $p = 0.9601$). These values were very small compared to 1, indicating that there were no species that relatively dominant in each SG.

The significant differences in terms of the number of species, diversity, evenness and dominant species do not appear in that of the four SGs. However, it appears that there is a tendency of SG 1 and SG 4 always has a value that is relatively different in terms of Shannon-Wiener's diversity index, Pielou's evenness index, and Simpson's index. SG 1 also appears to have the lowest mean value of species number (Figure 2). This probably indicating that SG 4 is more diverse than SG 1.

Discussion

The difference between the SGs is not very visible through the vegetation structure, but rather towards floristic composition. Indriyanto (2009) said that the species of lower plants depend on the species' ability adapting to a growing site and associated with other plants. According to Whitten *et al.*, (1996), the demarcation between the lowland and the mountain zones at 1000 m asl at Jawa Island determined mostly determined floristically. Furthermore, Larashati (2010) said that, the grouping of stand vegetation on Mount Salak especially related with canopy cover, which comprised with the area of closed canopy, open canopy, and very open canopy.

The species number, high diversity, high Evenness, and low Dominance Index probably because geographical position of Mount Salak, which is located at high altitude areas that support high precipitation. Most of the dominant species in the studied sites are members of fern life form. The abundance presences of fern species is an indicator of high precipitation. The similarity in terms of the structure of the vegetation may be caused by the location of the stands which at the zone of the tropical lower mountain forest at Mount Salak.

Espinosa *et al.*, (2011) in his research showed that the species composition of Tumbesian dry forests at the southern part of Aquador mainly determined by climate, topography, and soil. Water availability at that area significantly determined the species assemblage. Bai *et al.*, (2011) said that important factor in explaining the spatiotemporal trends within the vegetation were climatic factors.

CONCLUSION

The number of stands that make up the SG was different. The difference among the SG was determined by the difference in floristic composition. Vegetation structure obtained through diversity, evenness, species richness and dominance of species did not differ significantly between the SG.

REFERENCES

1. Bai, F., W. Sang., J. C. Axmacher. 2011. Forest vegetation responses to climate and environmental change: A case study from Changbai Mountain, NE China. *Forest Ecology and Management*. vol. 262 (11): p. 2052–2060.
2. Barbour, M.G., J.H. Burk., & W.P. Pitts. 1987. *Terrestrial Plant Ecology*. The Benjamin/Cumming Publishing Company Inc. Menlo Park, Reading, California, Massachusetts, Singapore.
3. Costa, F.R.C. 2004. Structure and composition of the ground-herb community in a terra-firme Central Amazonian forest. *Acta Amaz.* vol.34 (1): p. 53 - 59.
4. Cox, G.W. 1978. *Laboratory Manual of General Ecology*. W.M.C. Brown Company Publisher, Dubuque, Iowa.
5. Ekayani, M., Nuva, R. Yasmin, F. Sinaga, La Ode M. Maaruf., 2014. Wisata Alam Taman Nasional Gunung Halimun Salak: Solusi Kepentingan Ekologi dan Ekonomi (Natural tourism at Gunung Halimun Salak National Park: A solution for ecological and economic interest). *Jurnal Ilmu Pertanian Indonesia (JIPI)*. vol. 19 (1): p. 29 37.
6. Espinosa C, I., O. Cabrera, A. L. Luzuriaga., & A.Escudero. 2011. What factors affect diversity and composition of endengared tumbesian dry forests in Southern Ecuador. *Biotropica*. vol. 43(1): p. 15 - 22.
7. Galudra, G., M. Sirait., N. Ramdhaniaty., F. Soenarto., & B. Nurzaman., 2005. History of Land-Use Policies and Designation of Mount Halimun-Salak National Park (Sejarah Kebijakan Tata Ruang dan Penetapan Kawasan Taman Nasional Gunung Halimun-Salak). *Jurnal Manajemen Hutan Tropika*. vol. XI (1): p.1-13
8. GHSNPMP-JICA. 2009. *Ecological study Halimun-Salak Corridor Mount Halimun-Salak National Park*. GHSNPMP-JICA, Bogor
9. Hardjosuwarno, S. 1990. *Dasar-dasar Ekologi Tumbuhan*. Fakultas Biologi, Universitas Gadjah Mada, Jogjakarta.
10. Hart S. A., & Chen H. Y. H. 2006 Understory vegetation dynamics of North American boreal forests. *Critical Reviews in Plant Sciences*. Vol. 25(4): p. 381-397.
11. Indriyanto. 2009. *Komposisi Jenis Dan Pola Penyebaran Tumbuhan Bawah Pada Komunitas Hutan Yang Dikelola Petani di Register 19 Provinsi Lampung*. Seminar Hasil Penelitian & Pengabdian Kepada Masyarakat, Unila.
12. Ismaini, L., M. Lailati., Rustandi., & D. Sunandar. 2015. Analisis komposisi dan keanekaragaman tumbuhan di Gunung Dempo, Sumatera Selatan. *Pros Sem Nas Masy Biodiv Indon*. vol.1 (6): p. 1397-1402
13. Hilwa, I., & I. Masyrifina. 2015. Keanekaragaman jenis tumbuhan bawah di Gunung Papandayan bagian timur, Garut, Jawa Barat. *Jurnal Silvikultur Tropika*. vol. 6(2): p. 119-125
14. Kent, M. & P. Cooker. 1992. *Vegetation Description and Analysis*. ACRC Press, Belhaven Press, London.
15. Kinho, J. 2011. Karakteristik Morfologi Zingiberaceae Di Cagar Alam Gunung Ambang Sulawesi Utara. *Info BPK Manado*. vol.1(1): p. 35-49.
16. Koneri, R., D. D. Solihin., D. Buchori., & R. Tarumingkeng. 2010. Keanekaragaman Kumbang Lucanid (Coleoptera:Lucanidae) Pada Berbagai Ketinggian Tempat di Hutan Konsensi Unocal Gunung Salak, Jawa Barat. *Jurnal Matematika dan Sains*. vol.15 (2): p. 77-84.
17. Kusmana, 1997. *Metode Survey Vegetasi*. PT. Penerbit Institut Pertanian Bogor, Bogor
18. Larashati, I. 2010. *Penelitian Ekologi Jenis Tumbuhan Sebagai Dasar Pengelolaan Dan Pengembangan Taman Nasional Gunung Halimun-Salak*. Laporan Akhir Program Insentif Peneliti dan Perakayasa LIPI. Pusat Penelitian Biologi - LIPI.
19. Lalchhuanawma. 2008. *Ecological Studies on Plant Diversity and Productivity of Herbaceous Species in Mizoram University Campus at Tanhril, Aizawl, Mizoram (N.E. India)*. Department of Forest Ecology, Biodiversity and Environmental Sciences Mizoram University, Aizawl.

20. Lü, X.T., J. X. Yin., & J. W. Tang. 2011. Diversity and composition of understory vegetation in the tropical seasonal rain forest of Xishuangbanna, SW China. *Rev. Biol. Trop.* vol. 59 (1): p. 455-463,
21. Lukiastuti, F., & M. Hamdani. 2012. *Statistika Non Parametris. Aplikasinya dalam Bidang Ekonomi dan Bisnis.* CAPS, Yogyakarta.
22. Oksanen, J., F. G. Blanchet., R. Kindt., P. Legendre., P. R. Minchin., R. B. O'Hara, G. L. Simpson., P. Solymos., M. Henry H., Stevens & H. Wagner 2016. *vegan: Community Ecology Package.* R package version 2.3-5. <http://CRAN.R-project.org/package=vegan>
23. Omoro, L.M.A. & O. Luukkanen. 2011. *Native Tree Species Regeneration and Diversity in the Mountain Cloud Forests of East Africa, Biodiversity Loss in a Changing Planet*, PhD. Oscar Grillo (Ed.), InTech, DOI: 10.5772/23108. Available from: <http://www.intechopen.com/books/biodiversity-loss-in-a-changing-planet/native-tree-species-regeneration-and-diversity-in-the-mountain-cloud-forests-of-east-africa>
24. Prihantoro, D. E. 2013. Komposisi Keanekaragaman Jenis Tumbuhan Bawah Pasca Erupsi Merapi Di Petak 58 Rph Kaliurang, Pakem, Sleman, Yogyakarta. Tugas Akhir. Program Diploma III Sekolah Vokasi, Universitas Gadjah Mada, Yogyakarta
25. R Core Team .2015. *R: A language and environment for statistical computing.* R Foundation for Statistical Computing. Vienna, Austria. URL <http://www.R-project.org/>.
26. Schmidt & JHA Ferguson. 1951. Rainfall types based on wet and dry period ratios for Indonesia with WesternNew Guinea. Kementrian Perhubungan, Djawatan Meteorologi dan Geofisic, Jakarta. Verhandelingen, No.42.
27. Simon, H. 1993. *Hutan Jati dan Kemakmuran*, Aditya Media, Yogyakarta.
28. Suharti, S. 2015. Pemanfaatan tumbuhan bawah di zona pemanfaatan Taman Nasional Gunung Merapi oleh masyarakat sekitar hutan. PROS SEM NAS MASY BIODIV INDON vol. 1(6), ISSN: 2407-8050. P. 1411-1415.
29. Tabadepu, H., D. Buchori., & B. Sahari., 2008. Butterfly Record from Salak Mountain, Indonesia. *J. Entomol. Indon.* vol.5 (1): p.10-16
30. van Balen & Nijman, 2004. Biology and conservation of Pink-headed Fruit-dove *Ptilinopus porphyreus*. *Bird Conservation International.* 14: p. 139–152
31. Whitten, T., R.E. Soeriaatmadja., & S.A. Afitt. 1996. *The Ecology of Java and Bali.* The Ecology of Indonesia Series. vol. II. Periplus Editions, Singapore.