

## PAPER NAME

**1.C.1\_1.1\_2. IOP Muh Abdy -1028-1-012  
123-2018-The Application of Algebraic  
Methods in Balanced Incom**

---

## WORD COUNT

**1900 Words**

## CHARACTER COUNT

**8511 Characters**

## PAGE COUNT

**5 Pages**

## FILE SIZE

**933.3KB**

## SUBMISSION DATE

**Jun 20, 2023 10:01 AM GMT+8**

## REPORT DATE

**Jun 20, 2023 10:02 AM GMT+8**

---

● **19% Overall Similarity**

The combined total of all matches, including overlapping sources, for each database.

- 10% Internet database
- 13% Publications database
- Crossref database
- Crossref Posted Content database
- 15% Submitted Works database

● **Excluded from Similarity Report**

- Bibliographic material
- Manually excluded sources
- Manually excluded text blocks

PAPER • OPEN ACCESS

## The Application of Algebraic Methods in Balanced Incomplete Block Design

To cite this article: Muhammad Abdy and Wahida Sanusi 2018 *J. Phys.: Conf. Ser.* **1028** 012123

View the [article online](#) for updates and enhancements.

### Related content

- [Algebraic method for finding equivalence groups](#)  
Alexander Bihlo, Elsa Dos Santos Cardoso-Bihlo and Roman O Popovych
- [Comparison of the PIC model and the Lie algebraic method in the simulation of intense continuous beam transport](#)  
Zhao Xiao-Song and Lu Jian-Qin
- [A Generalized Variable-Coefficient Algebraic Method Exactly Solving \(3+1\)-Dimensional Kadomtsev–Petviashvili Equation\\*](#)  
Bai Cheng-Lin, Bai Cheng-Jie and Zhao Hong



**IOP | ebooks™**

Bringing you innovative digital publishing with leading voices to create your essential collection of books in STEM research.

Start exploring the collection - download the first chapter of every title for free.

# The Application of Algebraic Methods in Balanced Incomplete Block Design

Muhammad Abdy\* and Wahida Sanusi

Department of Mathematics, FMIPA Universitas Negeri Makassar, Indonesia

muh.abdy@unm.ac.id

**Abstract.** A balanced incomplete block design contains finite set  $P$  with  $v$  elements called varieties and a collection of set  $B$  consisting of  $b$  sets called blocks to qualify: (i). each block contains  $k$  varieties, (ii). Each variety occurs in every  $r$  blocks, (iii). Each pair of varieties occurs in exactly  $\lambda$  block. In this paper we discuss the use of algebraic method to construct such designs.

**Keywords**— Algebraic, BIBD

## 1. Introduction

In certain experiments using randomized block design (RBD), we may not be able to run all treatment combinations in each block. Situations like this usually occur because of shortages experimental equipment or facilities or the physical size of the block. To solve this problem, we can use a randomized block design where each treatment does not need to exist in every block. The design is known as balanced incomplete block design (BIBD). If all treatment comparisons are equally important, the treatment combinations used in each block should be selected in a balanced manner, so that any pair of treatments occur together the same number of times as any other pair. Thus, a BIBD is an incomplete block design in which any two treatments appear together an equal number of times. Suppose that there are  $a$  treatments and each block can hold exactly  $k$  ( $k < a$ ) treatments, then a BIBD may be constructed by taking  $\binom{a}{k}$  block and assigning a different combinations of treatments to each block. In this paper, we will apply the methods of modern algebra to construct a BIBD. Detailed discussion of statistical analysis of BIBD will not be discussed in this paper. Likewise, the algebraic method. We will only discuss the use of the methods of modern algebra at BIBD. For more details on the methods of modern algebra, see [1] and [2] and for a discussion of BIBD and its statistical analysis, see [3] and [4]

## 2. Algebraic Method of BIBD

### Definition 1

A BIBD with parameters  $(v, b, r, k, \lambda)$  is a pair  $(P, B)$  with the following properties:

- (i).  $P$  is a set with  $v$  elements.
- (ii).  $B = (B_1, \dots, B_b)$  is a subset of the power set  $\wp(P)$  with  $b$  elements.
- (iii). each  $B_i$  has exactly  $k$  elements, where  $k < v$ .



(iv). each unordered pair  $(p, q)$  with  $p, q \in P, p \neq q$ , occurs in exactly  $\lambda$  elements of  $B$ .

The sets  $B_1, \dots, B_b$  are called the block of BIBD. Each  $a \in P$  occurs in exactly  $r$  sets of  $B$ . Such a BIBD is also called a  $(v, b, r, k, \lambda)$  configuration or  $2$ - $2$ - $(v, k, \lambda)$  tactical configuration or design. The term "balanced" indicates that each pair of elements appears in exactly the same number of blocks, while the term "incomplete" means that each block contains less than  $v$  elements. A BIBD is symmetric if  $v = b$ .

### Definition 2

The incidence matrix of matrix  $a$   $(v, b, r, k, \lambda)$  configuration is the  $v \times b$  matrix  $A = (a_{ij})$ , where

$$a_{ij} = \begin{cases} 1 & \text{if } i \in B_j \\ 0 & \text{otherwise} \end{cases}$$

Here  $i$  denotes the  $i$ th element of the configuration.

Some properties of incidence matrices:

### Theorem 1

Let  $A$  is the incidence matrix of a  $(v, b, r, k, \lambda)$  configuration, then:

$$(i). AA^T = (r - \lambda)I_v + \lambda J_{vv}$$

$$(ii). \det(AA^T) = [r + (v - 1)\lambda](r - \lambda)^{v-1}$$

$$(iii). AJ_{bb} = rJ_{rb}$$

$$(iv). J_{vv}A = kJ_{vb}$$

where  $J_{mn}$  is the  $m \times n$  matrix with all entries equal to 1 and  $I_n$  is the  $n \times n$  identity matrix.

### Theorem 2

The following conditions are necessary for the existence of a BIBD with parameters  $v, b, r, k, \lambda$ :

$$(i). bk = rv$$

$$(ii). r(k - 1) = \lambda(v - 1)$$

$$(iii). b \geq v$$

Proof:

(i). we count of the number of ordered pairs consisting of a block and an element of the block as  $bk$ . Each of  $P$  must appear in  $r$  blocks  $B$ . Therefore, there are  $vr$  ordered pairs of an element and a block containing it. The numbers  $bk$  and  $vr$  count the number of elements of the same set, so  $bk = vr$ .

(ii). Follows by a similar argument

(iii). If  $v > \lambda$  then  $A^T$  has rank  $v$ , (by Theorem 1 (ii)). On the other hand, since the rank of  $A$  can be at most  $b$ , and since  $\text{rank } A \geq \text{rank } AA^T = v$ , then  $b \geq v$ . ■

### Example 1

Let  $\{1, 2, 3, 4, 5, 6, 7\}$  be the set of elements (varieties) and  $\{\{1, 2, 3\}, \{1, 5, 6\}, \{3, 4, 5\}, \{1, 4, 7\}, \{3, 6, 7\}, \{2, 5, 7\}, \{2, 4, 6\}\}$  is the set of the blocks. It is obtained a  $(7, 7, 3, 3, 1)$  configuration. This is the so called Fano geometry, the simplest example of a finite projective plane over  $F_2$ . The blocks are the set of points on lines and the element are 7 points of the plane.

There is a special type of algebraic structure, near rings, which can be used to construct block design.

### Definition 3

A set  $N$  with operations  $+$  and  $\cdot$  is near-ring provided that:

(i).  $(N, +)$  is a group (not necessarily commutative)

(ii).  $(N, \cdot)$  is a semigroup.

(iii).  $\forall n, n', n'' \in N: (n + n') \cdot n'' = n \cdot n'' + n' \cdot n''$ .

Each ring is a near-ring. Two ring axioms does not exist in near-ring, which is commutative of addition and distributive properties.

### Definition 4

Let  $N$  be a near-ring and  $n_1, n_2 \in N$ ;

$$n_1 \equiv n_2 : \Leftrightarrow \forall n \in N, n \cdot n_1 = n \cdot n_2$$

### Definition 5

The near-ring  $N$  is called planar if  $|N| \geq 3$  and if all the equations  $a \bullet x = x \bullet b + c$ , ( $a, b, c \in N$ ,  $a$  not mod of  $b$ ) have exactly one solution  $x \in N$ .

Near-ring planar produce BIBD's with parameters  $v, b, r, k, \lambda$  and efficiency  $E = \lambda v / rk$ . This  $E$  is a number between 0 and 1 and it estimate, for the statistical analysis of experiments, the "quality" of the design: .BIBD's with efficiency  $E \geq 0.75$  are usually considered to be "good" ones.

One get BIBD's from planar near rings in the following way: We define for  $a \in N$ ,  $N$  a planar near-ring,  $ga : N \rightarrow N$ ,  $n \mapsto n \bullet a$  and from  $G = \{ga \mid a \in N\}$ . Call  $a \in N$  as a "group-forming" if  $a \bullet N$  is a subgroup of  $(N, +)$ . Let us also call all sets  $a \bullet N + b$ , ( $a \in N^*$ ,  $b \in N$ ) as a block. Then these blocks together with  $N$  as a set of points from a tactical configuration with parameters:

$$(v, b, r, k, \lambda) = \left( v, \frac{\alpha_1 v}{|G|} + \alpha_2 v, \alpha_1 + \alpha_2 |G|, |G|, \lambda \right) \text{ where } v = |N| \text{ and } \alpha_1(\alpha_2) \text{ denote the number of}$$

orbit of  $F$  under the group- $G \setminus \{0\}$  which consists entirely of group forming (nongroup forming, respectively). This tactical configuration is a BIBD if and only if all either all elements are group-forming or just 0 is group forming.

**Example 2**

We consider the problem of testing combinations of six out of ten fertilizers, each on three fields.. By Definition 1, we have to look for a BIBD with parameters  $(v, 10, 6, 3, \lambda)$ . By Theorem 2 (i) and (ii) we obtain  $v = 5$  and  $\lambda = 3$ . Hence we are searching for a planar near- ring of order 5 and the near-ring  $N$ .

We construct the blocks  $a \bullet N + b$ , ( $a \neq 0$ ):

- $1 \bullet N + 0 = 4 \bullet N + 0 = \{0, 1, 4\} := B_1$
- $1 \bullet N + 1 = 4 \bullet N + 1 = \{1, 2, 0\} := B_2$
- $1 \bullet N + 2 = 4 \bullet N + 2 = \{2, 3, 1\} := B_3$
- $1 \bullet N + 3 = 4 \bullet N + 3 = \{3, 4, 2\} := B_4$
- $1 \bullet N + 4 = 4 \bullet N + 4 = \{4, 0, 3\} := B_5$
- $2 \bullet N + 0 = 3 \bullet N + 0 = \{0, 2, 3\} := B_6$
- $2 \bullet N + 1 = 3 \bullet N + 1 = \{1, 3, 4\} := B_7$
- $2 \bullet N + 2 = 3 \bullet N + 2 = \{2, 4, 0\} := B_8$
- $2 \bullet N + 3 = 3 \bullet N + 3 = \{3, 0, 1\} := B_9$
- $2 \bullet N + 4 = 3 \bullet N + 4 = \{4, 1, 2\} := B_{10}$

It is clear that only 0-is group-forming. Parameters can also be obtained directly from:

- (i)  $v = 5$  points ( $ie: 0, 1, 2, 3, 4$ )
- (ii)  $b = 10$  blocks
- (iii) Each point lies in exactly  $r = 6$  blocks
- (iv) Each block contains precisely  $k = 3$  elements
- (v) Each pair of different points appears in  $\lambda = 3$  blocks

To solve our fertilizer problem, we divide whole field into five parts which we call 0, 1, 2, 3, 4. Then we use fertilizers  $F_i$  on every field of the block  $B_i$  ( $i = 1, 2, \dots, 10$ ).

Field 0	Field 1	Field 2	Field 3	Field 4
$F_1$	$F_1$			$F_1$
$F_2$	$F_2$	$F_2$		
	$F_3$	$F_3$	$F_3$	

		$F_4$	$F_4$	$F_4$
$F_5$			$F_5$	$F_5$
$F_6$		$F_6$	$F_6$	
	$F_7$		$F_7$	$F_7$
$F_8$		$F_8$		$F_8$
$F_9$	$F_9$		$F_9$	
	$F_{10}$	$F_{10}$		$F_{10}$

Then we obtain: Every field contains exactly 6 fertilizers, and every fertilizer is applied on three fields. Every pair of different fields have three fertilizers in common. The efficiency  $E$  of this BIBD is  $E = (\lambda v/rk) = 0.83$

### 3. Conclusion

The method used to construct the BIBD is still possible if the number of elements of  $P$  is not too much. The problems will arise if in constructing the BIBD there are elements of  $P$  that is too large. It will be very difficult to do. For situation which  $P$  is too large, we can use graph theory as in [Budayasa, 2004].

### References

- [1] Durbin J R 2009 *Modern Algebra 6<sup>th</sup> eds.* (New York: John Wiley and Sons)
- [2] Choudhary P 2008 *Abstract Algebra* (Delhi: Oxford Book Company)
- [3] Silva M B (ed.) 2013 *Design of Experiments: Applications* (Croatia: InTeOp)
- [4] Montgomery D C 2012 *Design and Analysis of Experiments 12<sup>nd</sup> eds.* (New York: John Wiley and Sons)
- [5] Lidl R and Pilz G 1984 *Applied Abstract Algebra* (New York: Springer-Verlag)
- [6] Budayasa K 2004 Rancangan Blok Simetris dengan Varitas Relatif Besar *Makalah Seminar Nasional Matematika*, ITS Surabaya.

● **19% Overall Similarity**

Top sources found in the following databases:

- 10% Internet database
- 13% Publications database
- Crossref database
- Crossref Posted Content database
- 15% Submitted Works database

TOP SOURCES

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

<b>1</b>	<b>D. G. Kabe, A. K. Gupta. "Experimental Designs: Exercises and Solution...</b>	<b>3%</b>
	Crossref	
<b>2</b>	<b>Higher Education Commission Pakistan on 2022-02-15</b>	<b>2%</b>
	Submitted works	
<b>3</b>	<b>neliti.com</b>	<b>2%</b>
	Internet	
<b>4</b>	<b>Prasad, . "Fundamentals of Near Rings", Near Rings Fuzzy Ideals and G...</b>	<b>2%</b>
	Crossref	
<b>5</b>	<b>spada.uns.ac.id</b>	<b>2%</b>
	Internet	
<b>6</b>	<b>The University of Manchester on 2020-05-14</b>	<b>1%</b>
	Submitted works	
<b>7</b>	<b>"Computational Complexity", Springer Nature, 2012</b>	<b>1%</b>
	Crossref	
<b>8</b>	<b>North West University on 2015-05-01</b>	<b>1%</b>
	Submitted works	

- 9

**R.C. BOSE. "Characterization Problems of Combinatorial Graph Theory...**

Crossref

<1%
- 10

**d-nb.info**

Internet

<1%
- 11

**Indian Institute of Technology, Bombay on 2016-11-30**

Submitted works

<1%
- 12

**Tariq Alraqad. "New families of non-embeddable quasi-derived design...**

Crossref

<1%
- 13

**mafiadoc.com**

Internet

<1%
- 14

**Asma Ali, Inzamam ul Huque. "Chapter 4 Structure ofPrime Near Rings ...**

Crossref

<1%
- 15

**N.J. Groenewald, P.C. Potgieter. "A generalisation of regularities in nea...**

Crossref

<1%
- 16

**Higher Education Commission Pakistan on 2009-12-24**

Submitted works

<1%



● Excluded from Similarity Report

- Bibliographic material
- Manually excluded text blocks
- Manually excluded sources

---

EXCLUDED SOURCES

**Muhammad Abdy, Wahida Sanusi. "The Application of Algebraic Methods in B... 77%**

Crossref

---

**vdoc.pub 50%**

Internet

---

**Rudolf Lidl, Günter Pilz. "Applied Abstract Algebra", Springer Science and Busi... 45%**

Crossref

---

**Rudolf Lidl, Günter Pilz. "Applied Abstract Algebra", Springer Nature, 1984 45%**

Crossref

---

**Rudolf Lidl, Günter Pilz. "Chapter 5 Further Applications of Fields and Groups",... 45%**

Crossref

---

**Rudolf Lidl, Günter Pilz. "Applied Abstract Algebra", Springer Science and Busi... 25%**

Crossref

---

**Undergraduate Texts in Mathematics, 1998. 25%**

Crossref

---

**Rudolf Lidl, Günter Pilz. "Applied Abstract Algebra", Springer Nature, 1998 25%**

Crossref

---

**Rudolf Lidl, Günter Pilz. "Chapter 7 Further Applications of Algebra", Springer ... 24%**

Crossref

---

**Lidl, Rudolf, and Günter Pilz. "Further Applications of Algebra", Undergraduate ... 24%**

Crossref

<b>fs.unm.edu</b>	<b>21%</b>
Internet	
<b>ar.scribd.com</b>	<b>21%</b>
Internet	
<b>gallup.unm.edu</b>	<b>20%</b>
Internet	
<b>preprints.readingroo.ms</b>	<b>20%</b>
Internet	
<b>pt.scribd.com</b>	<b>20%</b>
Internet	
<b>readbag.com</b>	<b>20%</b>
Internet	
<b>fr.scribd.com</b>	<b>20%</b>
Internet	
<b>archive.org</b>	<b>19%</b>
Internet	
<b>baixardoc.com</b>	<b>19%</b>
Internet	
<b>uap.unnes.ac.id</b>	<b>19%</b>
Internet	
<b>ia801307.us.archive.org</b>	<b>19%</b>
Internet	
<b>docshare.tips</b>	<b>18%</b>
Internet	

<b>fs.gallup.unm.edu</b>	<b>18%</b>
Internet	
<b>id.scribd.com</b>	<b>18%</b>
Internet	
<b>docplayer.net</b>	<b>18%</b>
Internet	
<b>docstoc.com</b>	<b>17%</b>
Internet	
<b>coursehero.com</b>	<b>14%</b>
Internet	
<b>epdf.pub</b>	<b>11%</b>
Internet	
<b>IAN F. BLAKE, RONALD C. MULLIN. "Combinatorial Constructions and Coding"...</b>	<b>9%</b>
Crossref	
<b>IAN F. BLAKE, RONALD C. MULLIN. "Combinatorial Constructions and Coding"...</b>	<b>7%</b>
Crossref	
<b>ebookreading.net</b>	<b>7%</b>
Internet	
<b>Sim University on 2018-03-19</b>	<b>7%</b>
Submitted works	
<b>vibdoc.com</b>	<b>7%</b>
Internet	
<b>moam.info</b>	<b>7%</b>
Internet	

<b>edoc.tips</b>	7%
Internet	
<b>mohsenmehdipour.profile.semnan.ac.ir</b>	7%
Internet	
<b>Roger Louis Goodwin. "Class Notes in Operations Research, Statistics, and Pr...</b>	7%
Internet	
<b>pdfcoffee.com</b>	7%
Internet	
<b>Sim University on 2019-07-10</b>	7%
Submitted works	
<b>Günter E. Pilz. "Near-rings and Near-fields", Elsevier BV, 1996</b>	6%
Crossref	
<b>Alexander V. Mikhalev, Günter F. Pilz. "The Concise Handbook of Algebra", Sp...</b>	6%
Crossref	
<b>Alexander V. Mikhalev, Günter F. Pilz. "The Concise Handbook of Algebra", Sp...</b>	6%
Crossref	
<b>repositories.lib.utexas.edu</b>	5%
Internet	
<b>University of Dayton on 2018-03-27</b>	5%
Submitted works	
<b>Higher Education Commission Pakistan on 2016-08-09</b>	5%
Submitted works	
<b>Higher Education Commission Pakistan on 2014-01-17</b>	5%
Submitted works	

<b>Higher Education Commission Pakistan on 2013-11-22</b>	<b>5%</b>
Submitted works	
<hr/>	
<b>econwpa.repec.org</b>	<b>5%</b>
Internet	
<hr/>	
<b>econwpa.ub.uni-muenchen.de</b>	<b>5%</b>
Internet	
<hr/>	
<b>researchgate.net</b>	<b>4%</b>
Internet	
<hr/>	
<b>seminar.fmipa.unp.ac.id</b>	<b>4%</b>
Internet	
<hr/>	
<b>doczz.net</b>	<b>4%</b>
Internet	
<hr/>	
<b>china.iopscience.iop.org</b>	<b>4%</b>
Internet	
<hr/>	
<b>University of Dayton on 2017-04-01</b>	<b>3%</b>
Submitted works	
<hr/>	
<b>vtechworks.lib.vt.edu</b>	<b>3%</b>
Internet	
<hr/>	
<b>hdl.handle.net</b>	<b>2%</b>
Internet	
<hr/>	
<b>txspace.tamu.edu</b>	<b>2%</b>
Internet	
<hr/>	
<b>repository.tamu.edu</b>	<b>2%</b>
Internet	

## EXCLUDED TEXT BLOCKS

**Journal of Physics: Conference Series PAPER • OPEN ACCESS Related content The**

Universitas Hasanuddin on 2018-12-11

**Muhammad Abdy and Wahida Sanusi 2018 J. Phys.: Conf. Ser. 1028 012123**

china.iopscience.iop.org

**View the article online for updates and enhancements**

eprints.unm.ac.id

**Algebraic method for finding equivalence groups**

Alexander Bihlo, Elsa Dos Santos Cardoso-Bihlo, Roman O Popovych. "Algebraic method for finding equivalence...

**Comparison of the PIC model and the Lie**

china.iopscience.iop.org

**algebraic method in the simulation of intense continuous beam transport Zhao Xiao...**

china.iopscience.iop.org

**A Generalized Variable-Coefficient Algebraic Method Exactly Solving (3+1**

Huang Ding-Jiang, Zhang Hong-Qing. "Variable-Coefficient Hyperbola Function Method and Its Application t...

**Dimensional Kadomtsev-Petviashvili Equation\* Bai Cheng-Lin, Bai Cheng-Jie and Zh...**

Huang Ding-Jiang, Zhang Hong-Qing. "Variable-Coefficient Hyperbola Function Method and Its Application t...

**This content was downloaded from IP address 202.67**

Universitas Hasanuddin on 2018-12-11

**2nd International Conference on Statistics, Mathematics, Teaching, and Research...**

Universitas Hasanuddin on 2018-12-11

**The Application of Algebraic Methods in Balanced Incomplete Block Design Muham...**

doczz.net

**Department of Mathematics, FMIPA Universitas Negeri Makassar, Indonesia**

M Abdy, S Zenin, Irwan. "An Over view on intuitionistic fuzzy topological spaces", Journal of Physics: Confer...

---

**unm.ac.idAbstract**

Muhammad Basri, Andi Anto Patak. "Exploring Indonesian students' perception on Mendeley Reference Man..."

---

**Content from this work may be used under the terms of the Creative Commons Att...**

core.ac.uk

---

**Field 0Field 1Field2FieldField 4**

Günter E. Pilz. "Near-rings and Near-fields", Elsevier BV, 1996

---

**3F1F1F1F2F2F2F3F3F3**

www.ikp.uni-koeln.de

---

**F4F4F5F5**

University of Sydney on 2021-05-31

---

**N + 1**

econwpa.ub.uni-muenchen.de

---

**2 = {2, 3, 1**

Chester College of Higher Education on 2018-05-01

---

**B4**

H. Hassani, J. A. Tenreiro Machado, Z. Avazzadeh, E. Naraghirad, S. Mehrabi. "Optimal solution of the fracti..."

---

**N + 1**

mohsenmehdipour.profile.semnan.ac.ir

---

**2**

mohsenmehdipour.profile.semnan.ac.ir

---