

PAPER NAME

1.C.1_1.1_6. AIP-Ramadhan_2192_2019_ Numerical solution of SIRS model for transmission of dengue feve

WORD COUNT

2796 Words

CHARACTER COUNT

12421 Characters

PAGE COUNT

9 Pages

FILE SIZE

570.0KB

SUBMISSION DATE

Jun 20, 2023 10:01 AM GMT+8

REPORT DATE

Jun 20, 2023 10:01 AM GMT+8

● **17% Overall Similarity**

The combined total of all matches, including overlapping sources, for each database.

- 14% Internet database
- 10% Publications database
- Crossref database
- Crossref Posted Content database
- 8% Submitted Works database

● **Excluded from Similarity Report**

- Bibliographic material
- Manually excluded sources
- Manually excluded text blocks

Numerical solution of SIRS model for transmission of dengue fever using Homotopy Perturbation Method in Makassar

Cite as: AIP Conference Proceedings 2192, 060015 (2019); <https://doi.org/10.1063/1.5139161>
Published Online: 19 December 2019

Nur Rezky Ramadhan, Syafruddin Side, Sahlan Sidjara, et al.



View Online



Export Citation

ARTICLES YOU MAY BE INTERESTED IN

[Rayleigh-Ritz approximation for the stability of localised waves](#)

AIP Conference Proceedings 2192, 060016 (2019); <https://doi.org/10.1063/1.5139162>

[Analyzing effect of harvesting on prey population when prey growth depend on fear-factor and Allee-effect](#)


AIP Conference Proceedings 2192, 060017 (2019); <https://doi.org/10.1063/1.5139163>

[A mathematical model for phase transition regulation on the cell cycle in early stage of breast cancer](#)

AIP Conference Proceedings 2192, 060014 (2019); <https://doi.org/10.1063/1.5139160>

Webinar

Interfaces: how they make
or break a nanodevice



March 29th – Register now

 Zurich
Instruments

Numerical Solution of SIRS Model for Transmission of Dengue Fever using Homotopy Perturbation Method in Makassar

Nur Rezky Ramadhan^{2,a)}, Syafruddin Side^{1,b)}, Sahlan Sidjara^{1,c)}, Irwan^{1,d)} and Wahidah Sanusi^{1,e)}

¹Department of Mathematics, Universitas Negeri Makassar, Indonesia 90222

²Postgraduate student of Mathematics Department, Universitas Negeri Makassar, Indonesia 90222

a)ramadhannurrezky@gmail.com

b)syafruddin@unm.ac.id

c)sahlansidjara@unm.ac.id

d)irwanthaha@unm.ac.id

e)Corresponding author: wahidah.sanusi@unm.ac.id

Abstract. This study purposed to find a numerical solution for the Suspected-Infected-Recovered-Suspected (SIRS) model for transmission of dengue fever in the city of Makassar using the Homotopy Perturbation Method. The data used is secondary data on the number of dengue cases in Makassar City. The method used is the Homotopy Perturbation Method. The results obtained are numerical solutions of the SIRS model with the Perturbation Homotopy method. The results of the model analysis also show a graph of the trend in the trend of dengue spread in Makassar City via the SIRS model. These results predict the number of dengue cases in the city of Makassar, so that early prevention measures for the spread of dengue in the city of Makassar are of concern to the government.

Keywords: SIRS model, Perturbation Homotopy method, Dengue Fever Disease.

INTRODUCTION

Dengue Fever (DF) is a type of disease in the tropics area, especially indonesia, including in South Sulawesi [1]. Based on the data from Public Health Office the number, the number of cases of DF in 2016 was as many as 142, futhermore in 2017 it increased 248 cases [2] and until February 2018 it already recorded 62 cases DF in makassar city and include in 10 categories for the biggest disease in makassar regional public hospital [2].

Mathematical modeling can be used to detect and predict the number of dengue fever cases by the model of Suspected, Infected, Recovered and Suspected (SIRS) [3, 4, 5, 6, 7, 8, 9, 10]. In this research, the SIRS Model has been construct and determineted by which numerical solution with homotopy perturbation method model SIRS. Determination of parameters, simulation and analysis of results is infered from secondary data on the number of dengue fever cases in makassar city. The result of the numerical solution on the SIRS model from the the two methods are then compared to see methods that are more accurate in control and handling transmission dengue fever in makassar city.

METHOD

This is an applied research. The model was built from the modification of the SIR model [8, 9]. The SIRS model for transmission of dengue fever was then determined solution by using homotopy perturbation method [11]. The data

values of parameter and initial values used are secondary data the number of cases dengue fever in Makassar. Model simulation was also conducted using MATLAB to predict the number of dengue fever cases in Makassar city.

RESULT AND DISCUSSION

Changes that occur in each human population in the transmission of DF for the SIRS model can be interpreted in the form of the following figure 1. Based on human and mosquito population diagram on figure 1, the rate of change




FIGURE 1. Human and mosquito population diagram for SIRS DF Transmission Model [8, 9]

humans which Suspected, Infected, Recovered and the rate of vector which Suspected and Infected dengue fever (DF) towards time can be interpreted as follows:

Human Population

$$\frac{dS_h}{dt} = \mu_h N_h - \frac{\beta_h b}{N_h} I_v S_h - \mu_h S_h + \theta_h R_h \quad (1)$$

$$\frac{dI_h}{dt} = \frac{\beta_h b}{N_h} I_v S_h - (\mu_h + \gamma_h) I_h \quad (2)$$

$$\frac{dR_h}{dt} = \gamma_h I_h - (\mu_h + \theta_h) R_h \quad (3)$$

Vector Population

$$\frac{dS_v}{dt} = \mu_v N_v - \frac{\beta_v b}{N_h} I_h S_v - \mu_v S_v \quad (4)$$

$$\frac{dI_v}{dt} = \frac{\beta_h b}{N_h} I_v S_h - \mu_v I_v \quad (5)$$

Procedures Homotopy Perturbation Method of Dengue Fever SIRS Model

Numerical solution of SIRS model of dengue fever transmission with homotopy perturbation method is shown by the following procedures:

1. The model is considered as a system of ordinary differential equations first order in general can be written as an equation:

$$\frac{du_1}{dt} + g_1(t, u_1, u_2, \dots, u_m) = f_1(t)$$

$$\frac{du_2}{dt} + g_2(t, u_1, u_2, \dots, u_m) = f_2(t)$$

$$\begin{aligned} & \vdots \\ & \frac{du_m}{dt} + g_m(t, u_1, u_2, \dots, u_m) = f_m(t) \end{aligned}$$

and satisfy initial condition $u_1(t_0) = c_1, u_2(t_0) = c_2, \dots, u_m(t_0) = c_m$.

2. System of equation (1) following the operator form:

$$L(u_1) + g_1(t, u_1, u_2, \dots, u_m) = f_1(t)$$

$$L(u_2) + g_2(t, u_1, u_2, \dots, u_m) = f_2(t)$$

\vdots

$$L(u_m) + g_m(t, u_1, u_2, \dots, u_m) = f_m(t)$$

with $L = \frac{d}{dt}$ is an operator linear and N_1, N_2, \dots, N_m are nonlinear operator.

3. The form of differential equation system that satisfies of the following equation:

$$Du_i(t) = L_i(t, u_1, u_2, \dots, u_n) + N_i(t, u_1, u_2, \dots, u_n) + g_i(t)$$

With L_i is a linear operator, N_i is a nonlinear operator and g_i is an unknown analytic function. Furthermore, in homotopy perturbation technique, a homotopy form is constructed:

$$Du_i(t) = p [L_i(t, u_1, u_2, \dots, u_n) + N_i(t, u_1, u_2, \dots, u_n) + g_i(t)],$$

$1 \leq i \leq n$ where p is immersion parameter is changed from zero to unit.

4. Suppose the initial approach as the following equations:

$$u_{1,0}(t) = v_1(t) = u_1(t_0) = c_1$$

$$u_{2,0}(t) = v_2(t) = u_2(t_0) = c_2$$

\vdots

$$u_{m,0}(t) = v_m(t) = u_m(t_0) = c_m$$

and

$$u_1(t) = u_{1,0}(t) + pu_{1,1}(t) + p^2u_{1,2}(t) + \dots$$

$$u_2(t) = u_{2,0}(t) + pu_{2,1}(t) + p^2u_{2,2}(t) + \dots$$

\vdots

$$u_m(t) = u_{m,0}(t) + pu_{m,1}(t) + p^2u_{m,2}(t) + \dots$$

5. Coefficient arrangement to the order of p , we have:

$$L(u_{1,1}) + L(v_1) + N_1(u_{1,0}, u_{2,0}, \dots, u_{m,0}) - f_1 = 0, \quad u_{1,1}(t_0) = 0$$

$$L(u_{2,1}) + L(v_2) + N_2(u_{1,0}, u_{2,0}, \dots, u_{m,0}) - f_2 = 0, \quad u_{2,1}(t_0) = 0$$

$$L(u_{3,1}) + L(v_3) + N_3(u_{1,0}, u_{2,0}, \dots, u_{m,0}) - f_3 = 0, \quad u_{3,1}(t_0) = 0$$

\vdots

$$L(u_{m,1}) + L(v_m) + N_m(u_{1,0}, u_{2,0}, \dots, u_{m,0}) - f_m = 0, \quad u_{m,1}(t_0) = 0$$

then

$$L(u_{1,2}) + N_1(u_{1,0}, u_{2,0}, \dots, u_{m,0}) - f_1 = 0, \quad u_{1,2}(t_0) = 0$$

$$L(u_{2,2}) + N_2(u_{1,0}, u_{2,0}, \dots, u_{m,0}) - f_2 = 0, \quad u_{2,2}(t_0) = 0$$

$$L(u_{3,2}) + N_3(u_{1,0}, u_{2,0}, \dots, u_{m,0}) - f_3 = 0, \quad u_{3,2}(t_0) = 0$$

⋮

$$L(u_{m,2}) + N_m(u_{1,0}, u_{2,0}, \dots, u_{m,0}) - f_m = 0, \quad u_{m,2}(t_0) = 0$$

and soon.

6. We have description of solution of SIRS model with homotopy perturbation method can be expressed by the following equation:

$$\phi_{1,n}(t) = u_1(t) = \lim_{p \rightarrow 1} u_1(t) = \sum_{k=0}^{n-1} u_{1,k}(t)$$

$$\phi_{2,n}(t) = u_2(t) = \lim_{p \rightarrow 1} u_2(t) = \sum_{k=0}^{n-1} u_{2,k}(t)$$

$$\phi_{3,n}(t) = u_3(t) = \lim_{p \rightarrow 1} u_3(t) = \sum_{k=0}^{n-1} u_{3,k}(t)$$

⋮

$$\phi_{m,n}(t) = u_m(t) = \lim_{p \rightarrow 1} u_m(t) = \sum_{k=0}^{n-1} u_{m,k}(t)$$

Homotopy Perturbation Method

Homotopy Perturbation Method is a semi analytic numeric method, in implement of iteration on this method, there are several steps as following: First step, write SIRS model dengue fever in form of:

$$\frac{dx}{dt} = \mu_h(1 - x(t)) - \alpha x(t)z(t) + \theta_h w(t) \quad (6)$$

$$\frac{dy}{dt} = \alpha x(t)z(t) + \beta y(t) \quad (7)$$

$$\frac{dz}{dt} = \gamma(1 - z(t))y(t) - \delta z(t) \quad (8)$$

$$\frac{dw}{dt} = \gamma_h y(t) - \varphi w(t) \quad (9)$$

11 With initial conditions:

$x_0(t) = c_1, y_0(t) = c_2, z_0(t) = c_3, w_0(t) = c_4$. According to homotopy perturbation method, built a homotopy for SIRS model from dengue fever as following condition:

$$v_1' - x_0' + p(x_0' - \mu_h(1 - v_1) + \alpha v_1 v_3 - \theta_h v_4) = 0 \quad (10)$$

$$v_2' - y_0' + p(y_0' - \alpha v_1 v_3 + \beta v_2) = 0 \quad (11)$$

$$v_3' - z_0' + p(z_0' - \gamma(1 - v_3)v_2 + \delta_1 v_3) = 0 \quad (12)$$

$$v_4' - w_0' + p(w_0' - \gamma_h v_2 + \varphi v_4) = 0 \quad (13)$$

We choose initial Approximation:

$$v_{1,0}(t) = x_0(t) = v_1(0) = c_1, \quad (14)$$

$$v_{2,0}(t) = y_0(t) = v_2(0) = c_2, \quad (15)$$

$$v_{3,0}(t) = z_0(t) = v_3(0) = c_3, \quad (16)$$

$$v_{4,0}(t) = w_0(t) = v_4(0) = c_4, \quad (17)$$

and

$$v_1(t) = v_{1,0}(t) + p v_{1,1}(t) + p^2 v_{1,2}(t) + p^3 v_{1,3}(t) + \dots \quad (18)$$

$$v_2(t) = v_{2,0}(t) + p v_{2,1}(t) + p^2 v_{2,2}(t) + p^3 v_{2,3}(t) + \dots \quad (19)$$

$$v_3(t) = v_{3,0}(t) + p v_{3,1}(t) + p^2 v_{3,2}(t) + p^3 v_{3,3}(t) + \dots \quad (20)$$

where $v_{i,j}$ ($i = 1, 2; j = 1, 2, 3, \dots$) are function but has been determined. Substituting equations (14)-(20) into equation (10)-(13), we obtained:

$$v_{1,1}' + \alpha v_{1,0} v_{3,0} + \mu_h v_{1,0} - \mu_h - \theta_h v_{4,0} = 0, \quad v_{1,1}(0) = 0, \quad (21)$$

$$v_{2,1}' - \alpha v_{1,0} v_{3,0} + \beta v_{2,0} = 0, \quad v_{2,1}(0) = 0, \quad (22)$$

$$v_{3,1}' - \gamma v_{2,0} + \delta_1 v_{3,0} + \gamma v_{2,0} v_{3,0} = 0, \quad v_{3,1}(0) = 0, \quad (23)$$

$$v_{4,1}' - \gamma_h v_{2,0} + \varphi v_{4,0} = 0, \quad v_{4,1}(0) = 0, \quad (24)$$

$$v_{1,2}' + \alpha v_{1,1} v_{3,0} + \alpha v_{1,0} v_{3,1} + \mu_h v_{1,1} - \mu_h - \theta_h v_{4,1} = 0, \quad v_{1,2}(0) = 0, \quad (25)$$

$$v_{2,2}' - \alpha v_{1,1} v_{3,0} - \alpha v_{1,0} v_{3,1} + \beta v_{2,1} = 0, \quad v_{2,2}(0) = 0, \quad (26)$$

$$v_{3,2}' - \gamma v_{2,1} + \delta_1 v_{3,1} + \gamma v_{2,1} v_{3,0} + \gamma v_{2,0} v_{3,1} = 0, \quad v_{3,2}(0) = 0, \quad (27)$$

$$v_{4,2}' - \gamma_h v_{2,1} + \varphi v_{4,1} = 0, \quad v_{4,2}(0) = 0, \quad (28)$$

$$v_{1,3}' + \alpha v_{1,2} v_{3,0} + \alpha v_{1,1} v_{3,1} + \alpha v_{1,0} v_{3,2} + \mu_h v_{1,2} - \mu_h - \theta_h v_{4,2} = 0, \quad v_{1,3}(0) = 0, \quad (29)$$

$$v_{2,3}' - \alpha v_{1,2} v_{3,0} - \alpha v_{1,1} v_{3,1} - \alpha v_{1,0} v_{3,2} + \beta v_{2,2} = 0, \quad v_{2,3}(0) = 0, \quad (30)$$

$$v'_{3,3} - \gamma v_{2,2} + \delta_1 v_{3,2} + \gamma v_{2,2} v_{3,0} + \gamma v_{2,1} v_{3,1} + \gamma v_{2,0} v_{3,2} = 0, \quad v_{3,3}(0) = 0, \quad (31)$$

$$v'_{4,3} - \gamma_h v_{2,2} + \varphi v_{4,2} = 0, \quad v_{4,3}(0) = 0, \quad (32)$$

$$v'_{1,4} + \alpha v_{1,3} v_{3,0} + \alpha v_{1,2} v_{3,1} + \alpha v_{1,1} v_{3,2} + \alpha v_{1,0} v_{3,3} + \mu_h v_{1,3} - \mu_h - \theta_h v_{4,3} = 0, \quad v_{1,4}(0) = 0, \quad (33)$$

$$v'_{2,4} - \alpha v_{1,3} v_{3,0} - \alpha v_{1,2} v_{3,1} - \alpha v_{1,1} v_{3,2} - \alpha v_{1,0} v_{3,3} + \beta v_{2,3} = 0, \quad v_{2,4}(0) = 0, \quad (34)$$

$$v'_{3,4} - \gamma v_{2,3} + \delta_1 v_{3,3} + \gamma v_{2,3} v_{3,0} + \gamma v_{2,2} v_{3,1} + \gamma v_{2,1} v_{3,2} + \gamma v_{2,0} v_{3,3} = 0, \quad v_{3,4}(0) = 0, \quad (35)$$

$$v'_{4,4} - \gamma_h v_{2,3} + \varphi v_{4,3} = 0, \quad v_{4,4}(0) = 0 \quad (36)$$

So obtained solution of differential equation as in equation (37)-(45), then,

$$v_{1,1} = \int_0^t [-\alpha v_{1,0} v_{3,0} - \mu_h v_{1,0} + \mu_h + \theta_h v_{4,0}] ds, \quad (37)$$

$$v_{2,1} = \int_0^t [\alpha v_{1,0} v_{3,0} - \beta v_{2,0}] ds \quad (38)$$

$$v_{3,1} = \int_0^t [\gamma v_{1,0} v_{2,0} - \delta_1 v_{3,0} - \gamma v_{2,0} v_{3,0}] ds \quad (39)$$

$$v_{4,1} = \int_0^t [\gamma_h v_{2,0} - \varphi v_{4,0}] ds \quad (40)$$

$$v_{1,2} = \int_0^t [-\mu_h v_{1,1} - \alpha v_{1,1} v_{3,0} - \alpha v_{1,0} v_{3,1} + \theta_h v_{4,1}] ds, \quad (41)$$

$$v_{2,2} = \int_0^t [-\beta v_{2,1} + \alpha v_{1,1} v_{3,0} + \alpha v_{1,0} v_{3,1}] ds \quad (42)$$

$$v_{3,2} = \int_0^t [\gamma v_{2,1} - \gamma v_{2,1} v_{3,0} - \delta_1 v_{3,1} - \gamma v_{2,0} v_{3,1}] ds \quad (43)$$

$$v_{4,2} = \int_0^t [\gamma_h v_{2,1} - \varphi v_{4,1}] ds \quad (44)$$

$$v_{1,3} = \int_0^t [\alpha v_{1,0} v_{3,2} - \alpha v_{1,1} v_{3,1} - \alpha v_{1,2} v_{3,0} - \mu_h v_{1,2} + \theta_h v_{4,2}] ds, \quad (45)$$

$$v_{2,3} = \int_0^t [-\beta v_{2,2} + \alpha v_{1,2} v_{3,0} + \alpha v_{1,1} + \alpha v_{1,0} v_{3,2}] ds \quad (46)$$

$$v_{3,3} = \int_0^t [\gamma v_{2,2} - \gamma v_{2,2} v_{3,0} - \gamma v_{2,1} v_{3,1} - \delta_1 v_{3,2} - \gamma v_{2,0} v_{3,2}] ds \quad (47)$$

$$v_{4,3} = \int_0^t [\gamma_h v_{2,2} - \varphi v_{4,2}] ds \quad (48)$$

$$v_{1,4} = \int_0^t [-\alpha v_{1,3} v_{3,0} - \alpha v_{1,2} v_{3,1} - \alpha v_{1,1} v_{3,2} - \alpha v_{1,0} v_{3,3} - \mu_h v_{1,3} + \theta_h v_{4,3}] ds, \quad (49)$$

$$v_{2,4} = \int_0^t [\alpha v_{1,3} v_{3,0} + \alpha v_{1,2} v_{3,1} + \alpha v_{1,1} v_{3,2} + \alpha v_{1,0} v_{3,3} - \beta v_{2,3}] ds \quad (50)$$

TABLE 1. Initial condition used in SIRS model for transmission dengue fever

Initial Value SIRS Model		
Variabel and Parameter	Value	Source
$x(0)$	$\frac{7675406}{7675893}$	KKRI Makassar City [12]
$y(0)$	$\frac{487}{7675893}$	KKRI Makassar City [12]
$w(0)$	0.000001	Side, 2013 [8]
$z(0)$	0.056	Side, 2013 [8]
N	7675893	KKRI Makassar City [12]
μ_h	0.000046	Side, 2013 [8]
α	0.232198	Side, 2013 [8]
β	0.328879	Side, 2013 [8]
γ	0.375	Side, 2013 [8]
γ_h	0.328833	Side, 2013 [8]
θ_h	0.001	Side, 2013 [8]
φ	0.001046	Side, 2013 [8]
δ	0.0323	Side, 2013 [8]

$$v_{3,4} = \int_0^t [\gamma v_{2,3} - \delta_1 v_{3,3} - \gamma v_{2,3} v_{3,0} - \gamma v_{2,2} v_{3,1} - \gamma v_{2,1} v_{3,2} - \gamma v_{2,0} v_{3,3}] ds \quad (51)$$

$$v_{4,4} = \int_0^t [\gamma_h v_{2,3} - \varphi v_{4,3}] ds \quad (52)$$

The initial value and parameters used in SIRS model for transmission dengue fever as shown in Table 1. From initial value in Table 1, we have $c_1 = \frac{7675406}{7675893}$, $c_2 = \frac{487}{7675893}$, $c_3 = 0.056$ and parameters value $\alpha = 0.232198$, $\beta = 0.328879$, $\gamma = 0,375$, $\delta = 0.0323$, $\gamma_h = 0.328833$, $\theta_h = 0.001$ and $\varphi = 0.001046$ obtain

$$v_{1,1} = 0.999936555 - 0.01304826t \quad (53)$$

$$v_{2,1} = (6.34454E - 05) + 0.013023129t \quad (54)$$

$$v_{3,1} = 0.056 - 0.00178634t \quad (55)$$

$$v_{4,1} = 0.000001 + (2.08629E - 05)t \quad (56)$$

⋮

Generally, obtained SIRS model solution for transmission of dengue fever are:

$$x(t) = \sum_{j=0}^4 v_{1,j} = 0.999936555 - 0.01304826t + 0.000585026t^2 - 0.001096818t^3 - 0.000242282t^4 \quad (57)$$

$$y(t) = \sum_{j=0}^4 v_{2,j} = (6.34454E - 05) + 0.013023129t + 0.003698608t^2 + 0.002313229t^3 + 0.001003117t^4 \quad (58)$$

$$z(t) = \sum_{j=0}^4 v_{3,j} = 0.056 - 0.00178634t + 0.004667929t^2 + 0.001167146t^3 + 0.000760837t^4 \quad (59)$$

$$w(t) = \sum_{j=0}^4 v_{4,j} = 0.000001 + (2.08629E - 05)t + 0.004282433t^2 + 0.001215984t^3 + 0.000760598t^4 \quad (60)$$

An Equation (57),(58),(59) and (60) are solutions of SIRS model for transmission of dengue fever using homotopy perturbation method.

CONCLUSION

Mathematical model of SIRS is a model for transmission of dengue fever with assumption that dengue fever sufferers who have recovered infected again. Solutions of SIRS model for transmission of dengue fever using homotopy perturbation method gives description about the number of individual are Suspected Exposed, Infected and Recovered dengue fever case on month until next year, the number of highest and lowest cases of individual infected dengue fever so that the government of Makassar city can take precautionary measures so that transmission of dengue fever could be solved.

ACKNOWLEDGMENTS

We would like thank to DIKTI No:122/UN36.9/PL/2019 for the financial supports.

REFERENCES

- [1] <http://makassar.rakyatku.com/post/penjelasan-dinkes-makassar-soal-angka-penderita-dbd-meningkat.html>. [Diakses tanggal 8 Agustus 2018].
- [2] <https://makassar.sindonews.com/read/5038/2/dbd-mewabah-dprd-makassar-minta-ada-langkah-antisipasi-1518134552> [Diakses tanggal 8 Agustus 2018].
- [3] Esteva .L and C. Vargas, *Influence of vertical and mechanical transmission on the dynamics of dengue disease* (*Math Biosciences*, 2000), 167, 51-64.
- [4] Pongsumpun. P, *Transmission model for dengue disease with and without the effect of extrinsic incubation period* (*KMITL Sci. Tech. J. Thailand*, 2006) , 6:74-82.
- [5] Bobby, Syafruddin, dkk, *Analisis Survival Pada Penderita Demam Berdarah Dengue (DBD) Dengan Menggunakan Regresi Cox Di RSUSawerigading Palopo* (PKPT DIKTI, 2017).
- [6] Derouich. M, A. Boutayeb and E.H. Twizell, *A Model of Dengue Fever* (Brunel Universty, England, 2003), 1-10.
- [7] Syafruddin, S. and M.S.M Noorani, *SEIR model for transmission of dengue fever in Selangor Malaysia* (*International Journal of Modern Physics*, 2012), vol (9):380-389.
- [8] Syafruddin. S. and M.S.M Noorani, *A SIR model for spread of dengue fever disease (simulation for south sulawesi Indonesia and selangor Malaysia)* (*World journal of modeling and simulation*, 2013), Vol. 9 (2):96-105.
- [9] Syafruddin S. et. all, *Lyapunov functions of SIR and SEIR model for transmission of dengue fever disease* (*International Journal Simulation and Process Modeling; Inderscience Publishers*, 2013), 8(2,3):177-184.
- [10] Wahyuddin, Syafruddin Side, dkk, *Pemodelan Matematika pada Penularan Penyakit Demam Berdarah Dengue di Kabupaten Polewali Mandar* (PKPT DIKTI, 2017).
- [11] Rangkuti. Y.M., Side. S. and Noorani. M.S.M, *Numerical analytic solution of SIR model of dengue fever disease in South Sulawesi using homotopy perturbation method and variational iteration method* (*Journal of Mathematical and Fundamental Sciences*, 2014) 46A(1), 91-105.
- [12] Kementerian Kesehatan Republik Indonesia Kota Makassar, 2017.

● **17% Overall Similarity**

Top sources found in the following databases:

- 14% Internet database
- 10% Publications database
- Crossref database
- Crossref Posted Content database
- 8% Submitted Works database

TOP SOURCES

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

1	adoc.pub Internet	4%
2	eprints.unm.ac.id Internet	2%
3	iopscience.iop.org Internet	2%
4	ijstr.org Internet	1%
5	talenta.usu.ac.id Internet	1%
6	hindawi.com Internet	<1%
7	S Side, A Saman, M I Pratama, N R Ramadhan, W Sanusi. "Numerical S... Crossref	<1%
8	International Islamic University Malaysia on 2014-04-25 Submitted works	<1%

- 9

Universiti Kebangsaan Malaysia on 2014-08-22

Submitted works

<1%
- 10

University of Melbourne on 2021-11-02

Submitted works

<1%
- 11

mat.usm.my

Internet

<1%
- 12

Hashim, I.. "On multistage homotopy-perturbation method applied to n...

Crossref

<1%
- 13

researchgate.net

Internet

<1%
- 14

Syafuruddin Side, Yulita Molliq Rangkuti, Dian Gerhana Pane, Marlina Se...

Crossref

<1%
- 15

mafiadoc.com

Internet

<1%
- 16

Wahidah Sanusi, Nasiah Badwi, Ahmad Zaki, Sahlan Sidjara, Nurwahid...

Crossref

<1%
- 17

Malik, Hafiz Abid Mahmood, Abdul Waheed Mahesar, Faiza Abid, Ahma...

Crossref

<1%
- 18

S Side, I Sari, W Sanusi. "SEIR Mathematical Model of Seizure fever in I...

Crossref

<1%
- 19

Syafuruddin Side. "Analysis and Simulation of SIRI Model for Dengue Fe...

Crossref

<1%
- 20

Higher Education Commission Pakistan on 2017-01-30

Submitted works

<1%

● Excluded from Similarity Report

- Bibliographic material
- Manually excluded text blocks
- Manually excluded sources

EXCLUDED SOURCES

Nur Rezky Ramadhan, Syafruddin Side, Sahlan Sidjara, Irwan, Wahidah Sanusi... 49%

Crossref

aip.scitation.org 41%

Internet

W Sanusi, M I Pratama, M Rifandi, S Sidjara, Irwan, S Side. "Numerical Solutio..." 18%

Crossref

tib.eu 7%

Internet

Jian Ge, Wenjie Wang, Haobin Dong, Huan Liu, Xinglin Zhang, Wang Luo, Haiy... 2%

Crossref

math.fmipa.unm.ac.id 1%

Internet

journals.itb.ac.id 1%

Internet

EXCLUDED TEXT BLOCKS

Numerical solution of SIRS model for transmission of dengue fever using Homotop...

math.fmipa.unm.ac.id

Cite as

www.researchgate.net

Nur Rezky Ramadhan, Syafruddin Side, Sahlan Sidjara

S Side, I Sari, W Sanusi. "SEIR Mathematical Model of Seizure fever in Infants Under 5 Years Old in Makassar..."

et al. ARTICLES YOU MAY BE INTERESTED IN

Universitas Indonesia on 2023-02-08

Rayleigh-Ritz approximation for the stability of localised waves AIP Conference Pro...

Rahmi Rusin, Hadi Susanto. "Rayleigh-Ritz approximation for the stability of localised waves", AIP Publishing...

Analyzing effect of harvesting on prey population when prey growth depend on fea...

www.researchgate.net

A mathematical model for phase transition regulation on the cell cycle in early sta...

Ken Prameswari, Fajar Adi-Kusumo. "A mathematical model for phase transition regulation on the cell cycle ..."

AIP Conference Proceedings 2192

repository.unair.ac.id

of Mathematics, Universitas Negeri Makassar, Indonesia 90222

www.ijstr.org

a

Kharisma Noor Afifah, Nursyamsih Amalia, Syamsidar D., Abdul Haris, Subaer. "The influence of synthesis r..."

unm.ac.id)sahlansidjara@unm.ac.idd

ojs.unm.ac.id

unm.ac.id Abstract. This study

S Side, A Saman, M I Pratama, N R Ramadhan, W Sanusi. "Numerical Solution of the SEIR Model for Online G..."

ac.id) Corresponding author

Okti Ulandari, Widi Astuti, Slamet Sumardi, Agus Prasetya, Himawan Tri Bayu Murti Petrus. "Synthesis of soli..."

Numerical Solution of SIRS Model for Transmission of Dengue Fever using Homoto...

S Side, A Saman, M I Pratama, N R Ramadhan, W Sanusi. "Numerical Solution of the SEIR Model for Online G..."

Proceedings of The 8th SEAMS-UGM International Conference on Mathematics an...

Hermansah, Dedi Rosadi, Herni Utami, Abdurakhman, Gumgum Darmawan. "Hybrid MODWT-FFNN model f...

We would like thank to DIKTI No

Syafruddin Side. "Analysis and Simulation of SIRI Model for Dengue Fever Transmission", Indian Journal of ...

$du_1 + g_1(t, u_1, u_2, \dots, u_m) = f_1(t)dt$ $du_2 + g_2(t, u_1, u_2, \dots, u_m) = f_2(t)dt$

International Islamic University Malaysia on 2014-04-25

the operator form: $L(u_1$

ijnao.um.ac.ir

$u_{1,0}(t) = v_1(t) = u_1(t_0) = c_1 u_{2,0}(t) = v_2(t) = u_2(t_0) = c_2 \dots u_{m,0}(t) = v_m(t) = u_m(t_0) = \dots$

talenta.usu.ac.id

of p , we

International Islamic University Malaysia on 2014-04-25

$0, L$

talenta.usu.ac.id

$t_0) = 0$

Chowdhury, M.S.H.. "The multistage homotopy-perturbation method: A powerful scheme for handling the ...

$L(u_m$

talenta.usu.ac.id

$n(t) = u_2(t) = \lim$

talenta.usu.ac.id

$dx = \mu h(1 - x(t)) - \alpha x(t)z(t)$

adoc.pub

$u_2(t$

Filiz Tascan. "Application of adapted homotopy perturbation method for approximate solution of Henon-He...

$$\varphi_{m,n}(t) = u_m(t)$$

doi.org

$$u_3(t)$$

Filiz Tas̄ can. "Application of adapted homotopy perturbation method for approximate solution of Henon-He..."

$$k$$

digilib.unimed.ac.id

$$v'_1 - x'_0 + p(x'_0)$$

S Side, A Saman, M I Pratama, N R Ramadhan, W Sanusi. "Numerical Solution of the SEIR Model for Online G..."

$$2 - y'_0 + p(y'_0)$$

digilib.unimed.ac.id

$$v_{1,0}(t) = x_0(t) = v_1(0) = c_1$$

adoc.pub

$$v_{2,0}(t) = y_0(t) = v_2(0) = c_2, v_{3,0}(t) = z_0(t) = v_3(0) = c_3$$

adoc.pub

$$v_3(t) = v_{3,0}(t) + p v_{3,1}(t)$$

adoc.pub
