

## PAPER NAME

1.C.1\_1.1\_7. IOP Sanusi\_1752\_012004\_2  
021\_Numerical Solution of SIRS model f  
or Dengue Fever Transmiss

---

## WORD COUNT

**3640 Words**

## CHARACTER COUNT

**16679 Characters**

## PAGE COUNT

**12 Pages**

## FILE SIZE

**1.5MB**

## SUBMISSION DATE

**Jun 20, 2023 10:01 AM GMT+8**

## REPORT DATE

**Jun 20, 2023 10:01 AM GMT+8**

---

● **19% Overall Similarity**

The combined total of all matches, including overlapping sources, for each database.

- 8% Internet database
- 13% Publications database
- Crossref database
- Crossref Posted Content database
- 12% Submitted Works database

● **Excluded from Similarity Report**

- Bibliographic material
- Manually excluded sources
- Manually excluded text blocks

PAPER • OPEN ACCESS

## 32 Numerical Solution of SIRS model for Dengue Fever Transmission in Makassar City with Runge Kutta Method

To cite this article: W Sanusi *et al* 2021 *J. Phys.: Conf. Ser.* **1752** 012004

View the [article online](#) for updates and enhancements.

### You may also like

- [Analysis of dengue fever disease transmission using suspected-infected-recovered \(SIR\) model](#)  
Endro Setyo Cahyono, Alfensi Faruk and Bambang Suprihatin
- [Dengue transmission risk in a changing climate: Bangladesh is likely to experience a longer dengue fever season in the future](#)  
Kishor K Paul, Ian Macadam, Donna Green et al.
- [Autoregressive Moving Average Model and Improved LSTM Neural Network Applied in Epidemic Prediction in Zhejiang Province](#)  
Lingjie Kong



### 244<sup>th</sup> Electrochemical Society Meeting

October 8 – 12, 2023 • Gothenburg, Sweden

50 symposia in electrochemistry & solid state science

▶ Deadline Extended!  
**Last chance to submit!**

New deadline:  
April 21  
**submit your abstract!**

# Numerical Solution of SIRS model for Dengue Fever Transmission in Makassar City with Runge Kutta Method

W Sanusi<sup>1</sup>, M I Pratama<sup>2</sup>, M Rifandi<sup>3</sup>, S Sidjara<sup>1</sup>, Irwan<sup>1</sup> and S Side<sup>1\*</sup>

<sup>1</sup>Department of Mathematics, Universitas Negeri Makassar, Indonesia 90222

<sup>2</sup>Department of Actuarial Science, Universitas Muhammadiyah Bulukumba, Indonesia 92561

<sup>3</sup>Postgraduate student of Mathematics Department, Universitas Negeri Makassar, Indonesia 90222

\*e-mail: syafruddin@unm.ac.id

**Abstract.** The aim of this study was to obtain a SIRS model solution for the spread of dengue fever (DF) using the 4th order Runge-Kutta method. The data used was data on the number of dengue cases in Makassar city. The method used is the 4th order Runge Kutta Method. The results of this study are numerical solutions of the SIRS model on the spread of DF in Makassar city using the Runge Kutta method. Model simulation analysis shows that the estimated number of dengue cases in Makassar city, so that the government can take steps to prevent the spread of dengue in Makassar city.

**Keywords:** Numerical solutions, SIRS model, runge kutta Method

## 1. Introduction

Dengue fever (DF) is a type of disease in the tropics area, especially Indonesia, including in South Sulawesi [1]. Based on the data from Public Health Office, the number cases of DF in 2016 was 142 cases, furthermore in 2017 it increased to 248 cases [2] and until February 2018 62 cases had been recorded DF which was included in 10 category the biggest disease in Makassar regional public hospital [2].

Mathematical modeling can be used to detect and predict the number of dengue fever (DF) cases were done using the model of Suspected, Infected, Recovered and Suspected (SIRS) [3;4;5;6;7;8;9;10;11;12;13]. In this research, the SIRS model that has been constructed is determined by which numerical solution with Runge-Kutta Method. Determination of parameters, simulation and analysis of result using secondary data on the number of dengue fever cases in Makassar city. The result of numerical solution on the SIRS model with Runge-Kutta method to show result of accurate in control and handling transmission dengue fever in Makassar city.


## 2. Method

This research is applied research. The model was built from the modification of the SIR model [8,9]. The SIRS model for transmission of dengue fever was then determined solution by using the 4th order Runge Kutta Method [14]. The data values of the parameter and initial values used are secondary data the number of cases dengue fever in Makassar. The model simulation was also conducted using MAPLE 17 to predict the number of dengue fever cases in Makassar city.



### 3. Results and Discussion

#### 5.1. SIRS Model for dengue fever



**Figure 1.** Human and mosquito population diagram for SIRS DF Transmission Model [8,9]

Based on human and mosquito population diagram on Figure 1, the rate of change humans which Suspected, Infected, Recovered and the rate of vector which Suspected and Infected dengue fever (DHF) towards time can be interpreted as follows:

##### 3.1.1. Human Population.

$$\frac{dS_h}{dt} = \mu_h N_h - \frac{\beta_h b}{N_h} I_v S_h - \mu_h S_h + \theta_h R_h \tag{1}$$

$$\frac{dI_h}{dt} = \frac{\beta_h b}{N_h} I_v S_h - (\mu_h + \gamma_h) I_h \tag{2}$$

$$\frac{dR_h}{dt} = \gamma_h I_h - (\mu_h + \theta_h) R_h \tag{3}$$

##### 3.1.2. Vector Population

$$\frac{dS_v}{dt} = \mu_v N_v - \frac{\beta_v b}{N_h} I_h S_v - \mu_v S_v \tag{4}$$

$$\frac{dI_v}{dt} = \frac{\beta_v b}{N_h} I_h S_v - \mu_v I_v \tag{5}$$

#### 5.2. Runge-Kutta Method

The thought of Runge-Kutta method is to maintain Taylor approximation. However in solving ordinary differential equation with the Taylor method not practical because the method requires derivative estimation  $f(x,y)$ . Furthermore, not all a function are easy to calculate its derivatives, especially for difficult functions. The higher order of the Taylor method, its derivative must be calculated is also high [14].

Runge-Kutta Method is another alternative to the Taylor series method that doesn't require derivative. This method is trying to obtain a high degree of accuracy, and also to avoid looking for higher derivatives by evaluating functions  $f(x,y)$  with a fixed point on each interval. In general, the Runge-Kutta method has three characteristics [14]:

- This method is a one-step method, that its to obtain  $y_{n+1}$  only requires information available on the previous point  $(x_0, y_0)$ .
- This method is corresponding with the taylor series until  $h^p$  term, with  $p$  is different for a different method.
- This method doesn't require evaluation of each derivative  $f(x,y)$ , but the function itself.

2.2.1. The 4th Order Runge-Kutta Method.

The 4th order Runge-Kutta is the most accurate method compared to the previous order Runge-Kutta Method. Therefore, The 4th Runge-Kutta is often used to solve a differential equation. This method is obtained from the 2nd Runger-Kutta method second order with  $n=4$ , which has a form of an equation [13].

$$y_{r+1} = y_r + \frac{1}{6}(k_1 + 2k_2 + 2k_3 + k_4) \tag{6}$$

with

$$\begin{aligned}
 k_1 &= hf(x_r, y_r) \\
 k_2 &= hf(x_r + \frac{1}{2}h, y_r + \frac{1}{2}k_1) \\
 k_3 &= hf(x_r + \frac{1}{2}h, y_r + \frac{1}{2}k_2) \\
 k_4 &= hf(x_r + h, y_r + k_3)
 \end{aligned}
 \tag{7}$$

The 4th order Runge-Kutta method is having a high accuracy level of the solution reatherthan the previous order Runge-Kutta method, this method is also easy, stable and its have a small cutting and rounding error.

An equations (6) and (7) are resolved using the 4th order Runger-Kutta method have obtained degression from the general form the 4th orderRunge-Kutta method with  $n = 4$ . The general form of the 4th order Runge-Kutta as in equation (8).

$$y_{r+1} = y_r + (a_1k_1 + a_2k_2 + a_3k_3 + a_4k_4)h \tag{8}$$

with

$$k_1 = f(x_r, y_r) \tag{9}$$

$$k_2 = f(x_r + p_1h, y_r + q_{11}k_1h) \tag{10}$$

$$k_3 = f(x_r + p_2h, y_r + q_{21}k_1h + q_{22}k_2h) \tag{11}$$

$$k_4 = f(x_r + p_3h, y_r + q_{31}k_1h + q_{32}k_2h + q_{33}k_3h) \tag{12}$$

To obtain solution with using Runge-Kutta method fourth order, required Taylor series fourth order, such that by following general form of Taylor series equation is obtained equation (13).

$$y_{r+1} = y_r + f(x_r, y_r)h + f'(x_r, y_r)\frac{h^2}{2!} + f''(x_r, y_r)\frac{h^3}{3!} + f'''(x_r, y_r)\frac{h^4}{4!} + O(h^5) \tag{13}$$

With

$$f'(x_r, y_r) = \frac{\partial f}{\partial x} + f(x_r, y_r)\frac{\partial f}{\partial y} \tag{14}$$

$$\begin{aligned}
 f''(x_r, y_r) &= \left[ \frac{\partial^2 f}{\partial x^2} + 2f(x_r, y_r) \frac{\partial^2 f}{\partial x \partial y} + (f(x_r, y_r))^2 \frac{\partial^2 f}{\partial y^2} \right] + \left[ \left( \frac{\partial f}{\partial x} + f(x_r, y_r) \frac{\partial f}{\partial y} \right) \frac{\partial f}{\partial y} \right] \\
 f'''(x_r, y_r) &= \left[ \frac{\partial^3 f}{\partial x^3} + 3f(x_r, y_r) \frac{\partial^3 f}{\partial x^2 \partial y} + 3(f(x_r, y_r))^2 \frac{\partial^3 f}{\partial x \partial y^2} + \frac{\partial^3 f}{\partial y^3} \left( \frac{dy}{dx} \right)^3 \right] + \\
 &\quad \left[ 3 \left( \frac{\partial f}{\partial x} + f(x_r, y_r) \frac{\partial f}{\partial y} \right) \left( \frac{\partial^2 f}{\partial x \partial y} + f(x_r, y_r) \frac{\partial^2 f}{\partial y^2} \right) \right] + \\
 &\quad \left[ \frac{\partial^2 f}{\partial x^2} + 2f(x_r, y_r) \frac{\partial^2 f}{\partial x \partial y} + (f(x_r, y_r))^2 \frac{\partial^2 f}{\partial y^2} \right] + \left[ \left( \frac{\partial f}{\partial x} + f(x_r, y_r) \frac{\partial f}{\partial y} \right) \left( \frac{\partial f}{\partial y} \right)^2 \right]
 \end{aligned}
 \tag{15}$$

Then we obtained Taylor series fourth order in equation (16).

$$\begin{aligned}
 y_{r+1} &= y_r + f(x_r, y_r)h + \left[ \frac{\partial f}{\partial x} + f(x_r, y_r) \frac{\partial f}{\partial y} \right] \frac{h^2}{2} + \\
 &\quad \left[ \frac{\partial^2 f}{\partial x^2} + 2f(x_r, y_r) \frac{\partial^2 f}{\partial x \partial y} + (f(x_r, y_r))^2 \frac{\partial^2 f}{\partial y^2} \right] \frac{h^3}{6} + \left[ \frac{\partial f}{\partial x} \frac{\partial f}{\partial y} + f(x_r, y_r) \left( \frac{\partial f}{\partial y} \right)^2 \right] \frac{h^3}{6} + \\
 &\quad \left[ \frac{\partial^3 f}{\partial x^3} + 3f(x_r, y_r) \frac{\partial^3 f}{\partial x^2 \partial y} + 3(f(x_r, y_r))^2 \frac{\partial^3 f}{\partial x \partial y^2} + \frac{\partial^3 f}{\partial y^3} \left( \frac{dy}{dx} \right)^3 \right] \frac{h^4}{24} + \\
 &\quad \left[ 3 \left( \frac{\partial f}{\partial x} \frac{\partial^2 f}{\partial x \partial y} + f(x_r, y_r) \frac{\partial f}{\partial x} \frac{\partial^2 f}{\partial y^2} + f(x_r, y_r) \frac{\partial f}{\partial y} \frac{\partial^2 f}{\partial x \partial y} + (f(x_r, y_r))^2 \frac{\partial f}{\partial y} \frac{\partial^2 f}{\partial y^2} \right) \right] \frac{h^4}{24} + \\
 &\quad \left[ \frac{\partial^2 f}{\partial x^2} + 2f(x_r, y_r) \frac{\partial^2 f}{\partial x \partial y} + (f(x_r, y_r))^2 \frac{\partial^2 f}{\partial y^2} \right] \frac{h^4}{24} + \\
 &\quad \left[ \frac{\partial f}{\partial x} \left( \frac{\partial f}{\partial y} \right)^2 + f(x_r, y_r) \left( \frac{\partial f}{\partial y} \right)^3 \right] \frac{h^4}{24}
 \end{aligned}
 \tag{16}$$

After obtaining Taylor series fourth order in equation (16), its equation identified with equations (10)-(12) so we have a values  $a_1, a_2, a_3, a_4, p_1, p_2, p_3, q_{11}, q_{21}, q_{31}, q_{22}, q_{32},$  and  $q_{33}$ , where:

$$k_1 = f(x_r, y_r) \tag{17}$$

$$k_2 = f(x_r + p_1 h, y_r + q_{11} k_1 h) = f(x_r, y_r) + p_1 h \frac{\partial f}{\partial x} + q_{11} h f(x_r, y_r) \frac{\partial f}{\partial y} \tag{18}$$

$$k_3 = f(x_r + p_2 h, y_r + q_{21} k_1 h + q_{22} k_2 h)$$

$$k_3 = f(x_r, y_r) + p_2 h \frac{\partial f}{\partial x} + q_{21} h f(x_r, y_r) \frac{\partial f}{\partial y} + q_{22} h f(x_r, y_r) \frac{\partial f}{\partial y} + p_1 q_{22} h^2 \frac{\partial f}{\partial x} \frac{\partial f}{\partial y} + \tag{19}$$

$$q_{11} q_{22} h^2 f(x_r, y_r) \left( \frac{\partial f}{\partial y} \right)^2$$

$$\begin{aligned}
 k_4 &= f(x_r + p_3h, y_r + q_{31}k_1h + q_{32}k_2h + q_{33}k_3h) \\
 &= f(x_r, y_r) + p_3h \frac{\partial f}{\partial x} + q_{31}hf(x_r, y_r) \frac{\partial f}{\partial y} + q_{32}hf(x_r, y_r) \frac{\partial f}{\partial y} + p_1q_{32}h^2 \frac{\partial^2 f}{\partial x \partial y} + \\
 &\quad q_{11}q_{32}f(x_r, y_r)h^2 \left(\frac{\partial f}{\partial y}\right)^2 + q_{33}f(x_r, y_r)h \frac{\partial f}{\partial y} + p_2q_{33}h^2 \frac{\partial f}{\partial x} \frac{\partial f}{\partial y} + \\
 &\quad q_{21}q_{33}f(x_r, y_r)h^2 \left(\frac{\partial f}{\partial y}\right)^2 + q_{22}q_{33}f(x_r, y_r)h^2 \left(\frac{\partial f}{\partial y}\right)^2 + p_1q_{22}q_{33}h^3 \frac{\partial^2 f}{\partial x} \left(\frac{\partial f}{\partial y}\right)^2 + \\
 &\quad q_{11}q_{22}q_{33}f(x_r, y_r)h^3 \left(\frac{\partial f}{\partial y}\right)^3
 \end{aligned} \tag{20}$$

Substituting the values  $k_1, k_2, k_3,$  and  $k_4$  to equation (8).

$$\begin{aligned}
 y_{r+1} &= y_r + (a_1k_1 + a_2k_2 + a_3k_3 + a_4k_4)h = y_r + a_1k_1h + a_2k_2h + a_3k_3h + a_4k_4h \\
 y_{r+1} &= y_r + [a_1f(x_r, y_r) + a_2f(x_r, y_r) + a_3f(x_r, y_r) + a_4f(x_r, y_r)]h + \\
 &\quad \left[ a_2p_1 \frac{\partial f}{\partial x} + a_3p_2 \frac{\partial f}{\partial x} + a_4p_3 \frac{\partial f}{\partial x} + a_2q_{11}f(x_r, y_r) \frac{\partial f}{\partial y} + \right. \\
 &\quad \left. a_3q_{21}f(x_r, y_r) \frac{\partial f}{\partial y} + a_3q_{22}f(x_r, y_r) \frac{\partial f}{\partial y} + a_4q_{31}f(x_r, y_r) \frac{\partial f}{\partial y} + \right. \\
 &\quad \left. a_4q_{32}f(x_r, y_r) \frac{\partial f}{\partial y} + a_4q_{33}f(x_r, y_r) \frac{\partial f}{\partial y} \right] h^2 + \\
 &\quad \left[ a_3p_1q_{22} \frac{\partial^2 f}{\partial x \partial y} + a_4p_1q_{32} \frac{\partial^2 f}{\partial x \partial y} + a_4p_2q_{33} \frac{\partial^2 f}{\partial x \partial y} + \right. \\
 &\quad \left. a_3q_{11}q_{22}f(x_r, y_r) \left(\frac{\partial f}{\partial y}\right)^2 + a_4q_{11}q_{32}f(x_r, y_r) \left(\frac{\partial f}{\partial y}\right)^2 \right] h^3 + \\
 &\quad \left[ a_4q_{21}q_{33}f(x_r, y_r) \left(\frac{\partial f}{\partial y}\right)^2 + a_4q_{22}q_{33}f(x_r, y_r) \left(\frac{\partial f}{\partial y}\right)^2 \right] h^3 + \\
 &\quad \left[ a_4p_1q_{22}q_{33} \frac{\partial^2 f}{\partial x} \left(\frac{\partial f}{\partial y}\right)^2 + a_4q_{11}q_{22}q_{33}f(x_r, y_r) \left(\frac{\partial f}{\partial y}\right)^3 \right] h^4
 \end{aligned} \tag{21}$$

According to the comparison with two equations, we obtaining equation (22)

$$\begin{aligned}
 p_1 &= q_{11}, \\
 p_2 &= q_{21} + q_{22}, \\
 p_3 &= q_{31} + q_{32} + q_{33}, \\
 a_1 + a_2 + a_3 + a_4 &= 1,
 \end{aligned} \tag{22}$$

$$a_2 p_1 + a_3 p_2 + a_4 p_3 = \frac{1}{2},$$

$$a_3 p_1 q_{22} + a_4 (p_1 q_{32} + p_2 q_{33}) = \frac{1}{6},$$

$$a_1 p_1 q_{22} q_{33} = \frac{1}{24}.$$

So that from equation (23) we suppose that:

$$q_{11} = \frac{1}{2}, q_{21} = 0, q_{22} = \frac{1}{2}, q_{31} = 0, q_{32} = 0, \text{ dan } q_{33} = 1,$$

Then the solution with remaining parameters are:

$$p_1 = \frac{1}{2}, p_2 = \frac{1}{2}, p_3 = 1, \tag{23}$$

$$a_1 = \frac{1}{6}, a_2 = \frac{1}{3}, a_3 = \frac{1}{3}, a_4 = \frac{1}{6}. \tag{24}$$

The values of parameters are substituted to equation (13), so we have standard form Runge-Kutta method:

$$y_{r+1} = y_r + \frac{1}{6} (k_1 + 2k_2 + 2k_3 + k_4) \tag{25}$$

$$k_1 = hf(x_r, y_r) \tag{26}$$

$$k_2 = hf(x_r + \frac{1}{2}h, y_r + \frac{1}{2}k_1) \tag{27}$$

$$k_3 = hf(x_r + \frac{1}{2}h, y_r + \frac{1}{2}k_2) \tag{28}$$

with  $k_4 = hf(x_r + h, y_r + k_3) \tag{29}$

Futhermore, by substituting mathematical model for dengue fever transmission in equation (13) then, we have an equations:

$$x_{t+1} = x_t + \frac{1}{6} (k_1 + 2k_2 + 2k_3 + k_4)h \tag{30}$$

$$y_{t+1} = y_t + \frac{1}{6} (l_1 + 2l_2 + 2l_3 + l_4)h \tag{31}$$

$$z_{t+1} = z_t + \frac{1}{6} (m_1 + 2m_2 + 2m_3 + m_4)h \tag{32}$$

$$w_{t+1} = w_t + \frac{1}{6} (n_1 + 2n_2 + 2n_3 + n_4)h \tag{33}$$

with

$$k_1 = \mu_h(1 - x_t) - \alpha x_t z_t + \theta_h w_t \tag{34}$$

$$l_1 = \alpha x_t z_t - \beta y_t \tag{35}$$

$$m_1 = \gamma(1 - z_t)y_t - \delta z_t \tag{36}$$

$$n_1 = \gamma_h y_t - \varphi w_t \tag{37}$$



$$k_2 = \mu_h \left( 1 - (x_t + k_1 \frac{h}{2}) \right) - \alpha(x_t + k_1 \frac{h}{2})(z_t + m_1 \frac{h}{2}) + \theta_h(w_t + n_1 \frac{h}{2}) \tag{38}$$

$$l_2 = \alpha(x_t + k_1 \frac{h}{2})(z_t + m_1 \frac{h}{2}) - \beta(y_t + l_1 \frac{h}{2}) \tag{39}$$

$$m_2 = \gamma \left( 1 - (z_t + m_1 \frac{h}{2}) \right) (y_t + l_1 \frac{h}{2}) - \delta(z_t + m_1 \frac{h}{2}) \tag{40}$$

$$n_2 = \gamma_h(y_t + l_1 \frac{h}{2}) - \varphi(w_t + n_1 \frac{h}{2}) \tag{41}$$

$$k_3 = \mu_h \left( 1 - (x_t + k_2 \frac{h}{2}) \right) - \alpha(x_t + k_2 \frac{h}{2})(z_t + m_2 \frac{h}{2}) + \theta_h(w_t + n_2 \frac{h}{2}) \tag{42}$$

$$l_3 = \alpha(x_t + k_2 \frac{h}{2})(z_t + m_2 \frac{h}{2}) - \beta(y_t + l_2 \frac{h}{2}) \tag{43}$$

$$m_3 = \gamma \left( 1 - (z_t + m_2 \frac{h}{2}) \right) (y_t + l_2 \frac{h}{2}) - \delta(z_t + m_2 \frac{h}{2}) \tag{44}$$

$$n_3 = \gamma_h(y_t + l_2 \frac{h}{2}) - \varphi(w_t + n_2 \frac{h}{2}) \tag{45}$$

$$k_4 = \mu_h(1 - (x_t + k_3h)) - \alpha(x_t + k_3h)(z_t + m_3h) + \theta_h(w_t + n_3h) \tag{46}$$

$$l_4 = \alpha(x_t + k_3h)(z_t + m_3h) - \beta(y_t + l_3h) \tag{47}$$

$$m_4 = \gamma(1 - (z_t + m_3h))(y_t + l_3h) - \delta(z_t + m_3h) \tag{48}$$

$$n_4 = \gamma_h(y_t + l_3h) - \varphi(w_t + n_3h) \tag{49}$$

The initial value and parameters used in SIRS model for dengue fever transmission shown in table 1:

Table 1. Initial condition used in SIRS model for transmission dengue fever

Initial Value SIRS Model		
Variabel and Parameter	Value	Source
$x(0)$	$\frac{7675406}{7675893}$	[15]
$y(0)$	$\frac{487}{7675893}$	[15]
$w(0)$	0.000001	[10]
$z(0)$	0.056	[10]
$N$	7675893	[15]
$\mu_h$	0,000046	[10]
$\alpha$	0,232198	[10]
$\beta$	0,328879	[10]
$\gamma$	0,375	[10]
$\gamma_h$	0.328833	[10]
$\theta_h$	0.001	[10]
$\varphi$	0.001046	[10]
$\delta$	0,0323	[10]

By using a time interval  $0 \leq t \leq 100$  month, with  $h = 0,01$ , where initial condition and initial value in the form of parameter value in table 1, so that we obtained:

$$x_{0+1} = x_0 + \frac{1}{6}(\kappa_1 + 2k_2 + 2k_3 + k_4)h \tag{50}$$

$$y_{0+1} = y_0 + \frac{1}{6}(l_1 + 2l_2 + 2l_3 + l_4)h \tag{51}$$

$$z_{0+1} = z_0 + \frac{1}{6}(m_1 + 2m_2 + 2m_3 + m_4)h \tag{52}$$

$$w_{0+1} = w_0 + \frac{1}{6}(n_1 + 2n_2 + 2n_3 + n_4)h \tag{53}$$

with

$$k_1 = 0,000046 \left( 1 - \frac{7675406}{7675893} \right) - 0,232198 \left( \frac{7675406}{7675893} \right) 0,056 + (0,001)(0,000001) \tag{54}$$

$$l_1 = 0,232198 \left( \frac{7675406}{7675893} \right) 0,056 - 0,328879 \left( \frac{487}{7675893} \right) \tag{55}$$

$$m_1 = 0,375(1 - 0,056) \left( \frac{487}{7675893} \right) - 0,0323(0,056) \tag{56}$$

$$n_1 = (0,328833) \left( \frac{487}{7675893} \right) - (0,001046)(0,000001) \tag{57}$$

$$k_2 = (0,000046) \left( 1 - \left( \left( \frac{7675406}{7675893} \right) + k_1 \frac{0,01}{2} \right) \right) - (0,232198) \left( \frac{7675406}{7675893} \right) \tag{58}$$

$$+ k_1 \frac{0,01}{2} (0,056 + m_1 \frac{0,01}{2}) + (0,001)(0,000001 + n_1 \frac{0,01}{2})$$

$$l_2 = (0,232198) \left( \left( \frac{7675406}{7675893} \right) + k_1 \frac{0,01}{2} \right) (0,056 + m_1 \frac{0,01}{2}) - (0,328879) \left( \frac{487}{7675893} \right) \tag{59}$$

$$+ l_1 \frac{0,01}{2}$$

$$m_2 = (0,375) \left( 1 - \left( 0,056 + m_1 \frac{0,01}{2} \right) \right) \left( \frac{487}{7675893} \right) + l_1 \frac{0,01}{2} - (0,0323)(0,056) \tag{60}$$

$$+ m_1 \frac{0,01}{2}$$

$$n_2 = (0,328833) \left( \frac{487}{7675893} \right) + l_1 \frac{0,01}{2} - (0,001046)(0,000001 + n_1 \frac{0,01}{2}) \tag{61}$$

$$k_3 = (0,000046) \left( 1 - \left( \left( \frac{7675406}{7675893} \right) + k_2 \frac{0,01}{2} \right) \right) - (0,232198) \left( \frac{7675406}{7675893} \right) \tag{62}$$

$$+ k_2 \frac{0,01}{2} (0,056 + m_2 \frac{0,01}{2}) + (0,001)(0,000001 + n_2 \frac{0,01}{2})$$

$$l_3 = (0,232198) \left( \left( \frac{7675406}{7675893} \right) + k_2 \frac{0,01}{2} \right) (0,056 + m_2 \frac{0,01}{2}) \tag{63}$$

$$- (0,328879) \left( \frac{487}{7675893} \right) + l_2 \frac{0,01}{2}$$

$$m_3 = (0,375) \left( 1 - \left( 0,056 + m_2 \frac{0,01}{2} \right) \right) \left( \frac{487}{7675893} \right) + l_2 \frac{0,01}{2} - (0,0323)(0,056) \tag{64}$$

$$+ m_2 \frac{0,01}{2}$$

$$n_3 = (0,328833) \left( \frac{487}{7675893} \right) + l_2 \frac{0,01}{2} - (0,001046)(0,000001 + n_2 \frac{0,01}{2}) \tag{65}$$

$$k_4 = (0,000046) \left( 1 - \left( \left( \frac{7675406}{7675893} \right) + k_3(0,01) \right) \right) - (0,232198) \left( \frac{7675406}{7675893} \right) \tag{66}$$


$$l_4 = (0,232198) \left( \left( \frac{7675406}{7675893} \right) + k_3(0,01) \right) (0,056 + m_3(0,01)) + k_3(0,01) (0,056 + m_3(0,01)) + (0,001) (0,000001 + n_3(0,01)) \tag{67}$$

$$m_4 = (0,375) \left( 1 - (0,056 + m_3(0,01)) \right) \left( \left( \frac{487}{7675893} \right) + l_3(0,01) \right) - (0,328879) \left( \left( \frac{487}{7675893} \right) + l_3(0,01) \right) \tag{68}$$

$$n_4 = (0,328833) \left( \left( \frac{487}{7675893} \right) + l_3(0,01) \right) - (0,001046) (0,000001 + n_3(0,01)) \tag{69}$$

3.3. SIRS model simulation for dengue fever transmission

Based on exiting data and parameters, simulation model was completed with using MAPLE 17. The initial condition with used to model of this simulation are based on reported by Ministry of Health Republic Indonesia Makassar city 2017, as shown in table 1 and the result of simulation model are given by Figure 2 to Figure 5 is below



3 Figure 2. Prediction of the number of suspected dengue fever transmission 2






Figure 3. Prediction of the number of infected dengue fever transmission 2



**Figure 4.** Prediction of the number of infected vectors dengue fever transmission



**Figure 5.** Prediction of the number of recovered vectors dengue fever transmission

Figure 2 shown that the number of initial value suspect is 0,9999365 and capable to make a description about the next suspect number, and which suspects reaches the minimum on the month 60-th (5 years), this cause of the number of host population that switch status become infected and also at the same time the number of host population recovers, but must be remembered that those population still have the chance to get infected again.

The result of the SIRS model in Figure 3 shown that the number of dengue fever (DF) cases, in other words, the number of infected people is 14 to 18 months to reach the highest stage (12% from the human population) and would decrease to zero in the 45-th months. As for, the number of mosquitoes infected suitable in Figure 4, only takes time less than 30 months, more precisely the 28-th month to reach the highest number, i.e approach 60% from the mosquito population for Makassar city takes time 160 months (13 years) to approach zero. Figure 5 shown of a most significant increase in the population recovering from dengue fever, the positive trend would attain top in the 4 years (50-th month) and afterward, but this graph would slowly decline because in the SIRS, model the recovered population can be contracting again. Which could also be seen in the suspected graph, where its graph would slowly increasing in the 50-th month.

#### 4. Conclusion

By using the 4th order Runge-Kutta would be obtained an illustration the number of suspected, exposed, infected and recovered in next month and year, the number of increase and decrease of dengue fever cases in Makassar city. So, the government of makassar city can take precautionary measures so that the transmission of dengue fever could be solved. The 4th order Runge-Kutta method also shown the rate of development of the number of infected vectors.

#### 5. Acknowledgements

We would like thank to DIKTI No:122/UN36.9/PL/2019 for the financial supports.

#### References

- [1] <https://makassar.sindonews.com/read/5038/2/dbd-mewabah-dprd-makassar-minta-ada-langkah-antisipasi-1518134552>. [accessed, 14 July, 2019]
- [2] <http://makassar.rakyatku.com/post/penjelasan-dinkes-makassar-soal-angka-penderita-dbd-meningkat.html>. [accessed, 14 July, 2019]
- [3] Bobby, Syafruddin, dkk. 2017. Analisis Survival PadaPenderitaDemamBerdarah Dengue (DBD) DenganMenggunakanRegresi Cox Di RSUSawerigadingPalopo. PKPT DIKTI 2017.
- [4] Derouich, M, A Boutayeb& EH Twizell. 2003. A Model of Dengue Fever. *Brunel Universty*, England:1-10.
- [5] Edy Soewono dan Supriatna., A.K. : 2005. A two-dimensional model for the transmission of dengue fever disease.*Bulletin of the Malaysian Mathematical Sciences Society*: 49-57.
- [6] Esteva L,& C. Vargas. 2000. Influence of vertical and mechanical transmission on the dynamics of dengue disease, *Math Biosciences* 167:51-64.
- [7] Pongsumpun, P. 2006. Transmission model for dengue disease with and without the effect of extrinsic incubation period.*KMITL Sci. Tech. J. Thailand* 6:74-82.
- [8] Rangkuti, Y.M., Side, S., Noorani, M.S.M., 2014. Numerical analytic solution of SIR model of dengue fever diseasein South Sulawesi using homotopy perturbation method and variational iteration method. *Journal of Mathematical and Fundamental Sciences*.46A(1),91-105.
- [9] Syafruddin, S. and M.S.M Noorani. 2012. SEIR model for transmission of dengue fever in Selangor Malaysia. *International Journal of Modern Physics*: vol.9 (2012):380-389.
- [10] Syafruddin S. & M.S.M Noorani. 2013. A SIR model for spread of dengue fever disease (simulation for South Sulawesi Indonesia and Selangor Malaysia); *World Journal of Modeling and Simulation* 9 (2):96-105.
- [11] Syafruddin S. et. all. 2013. Lyapunov functions of SIR and SEIR model for transmission of dengue fever disease. *International Journal Simulation and Process Modeling; Inderscience Publishers* 8(2,3):177-184.
- [12] Syafruddin S., 2015. A Susceptible-Infected-Recovered Model and Simulation for Transmission of Tuberculosis. *Advanced Science Letters*. 21 (2):137-139.
- [13] Annas, S, Pratama, M. I, Rifandi, M., Sanusi, W, Side, S: Stability analysis and numerical simulation of SEIR model for pandemic COVID-19 spread in Indonesia, *Chaos, Solitons & Fractals*, 139, 110072 (2020)
- [14] Putri, P. P. 2013. Analisis Solusi Numerik Model Predator-Prey dengan Metode Runge-Kutta Orde Empat dan Gill. *Skripsi*. Fakultas MIPA Universitas Jember.
- [15] Ministry of Health of the Republic of Indonesia Makassar City, 2017.

● **19% Overall Similarity**

Top sources found in the following databases:

- 8% Internet database
- 13% Publications database
- Crossref database
- Crossref Posted Content database
- 12% Submitted Works database

TOP SOURCES

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

1	<b>S Side, A M Utami, Sukarna, M I Pratama. "Numerical solution of SIR m...</b>	2%
	Crossref	
2	<b>Syafruddin Side. "Analysis and Simulation of SIRI Model for Dengue Fe...</b>	2%
	Crossref	
3	<b>S Side, N Badwi, M Abdy, W Sanusi. "SEIR Model and Simulation for Ty...</b>	2%
	Crossref	
4	<b>Syafruddin Side, Gustman Putra Astari, Muh Isbar Pratama, Irwan, Wa...</b>	1%
	Crossref	
5	<b>ijstr.org</b>	1%
	Internet	
6	<b>University of Birmingham on 2016-03-25</b>	1%
	Submitted works	
7	<b>hindawi.com</b>	1%
	Internet	
8	<b>Central Queensland University on 2014-05-21</b>	<1%
	Submitted works	

- 9

**F. Lekien. "Tricubic interpolation in three dimensions", International Jo...**

Crossref

<1%
- 10

**journal.uin-alauddin.ac.id**

Internet

<1%
- 11

**coursehero.com**

Internet

<1%
- 12

**SYAFRUDDIN S., M. S. M. NOORANI. "SEIR MODEL FOR TRANSMISSIO...**

Crossref

<1%
- 13

**University of Lincoln on 2020-04-27**

Submitted works

<1%
- 14

**Universitas Pendidikan Indonesia on 2022-01-21**

Submitted works

<1%
- 15

**University of Warwick on 2012-05-16**

Submitted works

<1%
- 16

**Wahidah Sanusi, Nasiah Badwi, Ahmad Zaki, Sahlan Sidjara, Nurwahid...**

Crossref

<1%
- 17

**Li Ying-Le. "Distribution characteristic of scattering field for an ellipsoi...**

Crossref

<1%
- 18

**State Islamic University of Alauddin Makassar on 2021-03-22**

Submitted works

<1%
- 19

**repository.uin-suska.ac.id**

Internet

<1%
- 20

**Langlois, M.H.. "Protonation equilibrium and lipophilicity of moxifloxaci...**

Crossref

<1%

21	Suwardi Annas, Muh. Isbar Pratama, Muh. Rifandi, Wahidah Sanusi, Sy...	<1%
	Crossref	
22	digilib.unimed.ac.id	<1%
	Internet	
23	ict.csiro.au	<1%
	Internet	
24	Al Hussein bin Talal University on 2023-01-18	<1%
	Submitted works	
25	Aristotle University of Thessaloniki on 2021-02-27	<1%
	Submitted works	
26	Universitas Jember on 2015-11-18	<1%
	Submitted works	
27	University of Utah on 2016-05-02	<1%
	Submitted works	
28	Central Queensland University on 2013-05-20	<1%
	Submitted works	
29	Prentice Hall on 2002-02-18	<1%
	Submitted works	
30	publications.polymtl.ca	<1%
	Internet	
31	Huan-nan Shi. "Schur-Convex Functions and Inequalities", Walter de Gr...	<1%
	Crossref	
32	Wahidah Sanusi, Suwardi Annas, Muh. Isbar Pratama, Muh. Rifandi, Ir...	<1%
	Crossref	



33	<b>University of Southampton on 2007-02-01</b> Submitted works	<1%
34	<b>University of Wales Swansea on 2016-03-17</b> Submitted works	<1%
35	<b>vsip.info</b> Internet	<1%

● Excluded from Similarity Report

- Bibliographic material
- Manually excluded text blocks
- Manually excluded sources

---

EXCLUDED SOURCES

<b>W Sanusi, M I Pratama, M Rifandi, S Sidjara, Irwan, S Side. "Numerical Solutio...</b>	<b>49%</b>
Crossref	
<b>Nur Rezky Ramadhan, Syafruddin Side, Sahlan Sidjara, Irwan, Wahidah Sanusi...</b>	<b>12%</b>
Crossref	
<b>aip.scitation.org</b>	<b>11%</b>
Internet	
<b>researchgate.net</b>	<b>6%</b>
Internet	
<b>iopscience.iop.org</b>	<b>5%</b>
Internet	
<b>eprints.unm.ac.id</b>	<b>4%</b>
Internet	
<b>Nurhaeda, S Anas, S Side. "Analysis and simulation of mathematical model fo...</b>	<b>2%</b>
Crossref	
<b>academic-accelerator.com</b>	<b>1%</b>
Internet	
<b>semanticscholar.org</b>	<b>1%</b>
Internet	
<b>journal.universitاسbumigora.ac.id</b>	<b>&lt;1%</b>
Internet	

## EXCLUDED TEXT BLOCKS

**Journal of Physics: Conference Series PAPER • OPEN ACCESS** You may also like

Birkbeck College on 2023-03-03

**Analysis**

nlist.inflibnet.ac.in

**recovered (SIR) model** Endro Setyo Cahyono, Alfensi Faruk and Bambang Suprihatin

R Sriningsih, B W Otok, Sutikno. "Factors affecting the number of dengue fever cases in West Sumatra provi...

**Dengue transmission risk in a changing climate: Bangladesh is likely to experience...**

Kishor K Paul, Ian Macadam, Donna Green, David G Regan, Richard T Gray. "Dengue transmission risk in a ch...

**Kishor K Paul, Ian Macadam, Donna Green et al**

Md A H A Askary, T B Hai, S S K Showmik, Md A. Kader, Md Tauhidur Rahman. "The prospect of green energ...

**Autoregressive Moving Average Model and Improved LSTM Neural Network Applie...**

Hong Fang, Lan Zhang. "Analysis and Prediction of Air Quality based on LSTM Neural Network — Take Beij...

**W Sanusi**

iopscience.iop.org

**ICSMTR 2019 Journal of Physics: Conference Series IOP Publishing 1752 (2021**

School of Business and Management ITB on 2023-05-05

**114.125**

digilib.iainkendari.ac.id

**Numerical Solution of SIRS model for Dengue Fever Transmission in Makassar City...**

www.e-journal.my.id

**of Mathematics, Universitas Negeri Makassar, Indonesia 90222**

aip.scitation.org

## Science

M Asri, S Sidjara, W Sanusi, S Side, M I Pratama. "Analysis and Solution of The SEIRS Model for The Rubella ...

student of Mathematics Department, Universitas Negeri Makassar, Indonesia 90222

aip.scitation.org

**e-mail: syafuruddin@unm.ac.id** Abstract. The

S Side, N Badwi, M Abdy, W Sanusi. "SEIR Model and Simulation for Typus Disease", Journal of Physics: Con...

**Content from this work may be used under the terms of the Creative Commons Att...**

zagan.unizar.es

**We would like thank to DIKTI No:122/UN36.9/PL/2019 for the financial supports**

aip.scitation.org

## Numerical Solution

aip.scitation.org

**$yr?1 ? yr ? (k1 ? 2k2 ? 2k3 ? k4 )1(6$**

University of Wales Swansea on 2016-03-17

**$k1 ? hf (xr , yr )11$**

www.coursehero.com

**$22k4 ? hf$**

University of Newcastle on 2015-10-19

**$k1 ? f (xr , yr$**

journal.uin-alauddin.ac.id

**$f (xr , yr )h ? f '(xr , yr$**

University of Southampton on 2007-02-01

**$f '(xr , yr$**

journal.uin-alauddin.ac.id

**f (xr , yr**

journal.uin-alauddin.ac.id

---

**3 f**

F. Lekien. "Tricubic interpolation in three dimensions", International Journal for Numerical Methods in Engin...

---

**x?y**

Huan-nan Shi. "Schur-Convex Functions and Inequalities", Walter de Gruyter GmbH, 2019

---

**f (xr , yr ) ?f**

Huan-nan Shi. "Schur-Convex Functions and Inequalities", Walter de Gruyter GmbH, 2019

---

**2 f ??? ??x? f**

Huan-nan Shi. "Schur-Convex Functions and Inequalities", Walter de Gruyter GmbH, 2019

---

**f (xr , yr ) ?f**

Huan-nan Shi. "Schur-Convex Functions and Inequalities", Walter de Gruyter GmbH, 2019

---

**yr?1 ? yr ? f(xr,yr**

www.coursehero.com

---

**h244????3???? ??fx ??x2**

State Islamic University of Alauddin Makassar on 2021-08-20

---

**f? ?f ?2**

Rumen L. Mishkov. "Generalization of the formula of Faa di Bruno for a composite function with a vector arg...

---

**6? ??? ? 3 f?? ?x 3**

Rumen L. Mishkov. "Generalization of the formula of Faa di Bruno for a composite function with a vector arg...

---

**xr , yr**

Anadolu University on 2020-11-22

---

**a1, a2 , a3 , a4 , p1, p2 , p3 , q11, q21, q31, q22, q32**

repository.uin-suska.ac.id

---

**k1 ? f (xr , yr**

journal.uin-alauddin.ac.id

---

**k3 ? f (xr ? p2h, yr ? q21k1h ? q22k2h**

journal.uin-alauddin.ac.id

---

**k4 ? f (xr ? p3h, yr ? q31k1h ? q32k2h ? q33k3h**

journal.uin-alauddin.ac.id

---

**f (xr , yr**

journal.uin-alauddin.ac.id

---

**a1k1 ? a2k2 ? a3k3 ? a4k4)h**

State Islamic University of Alauddin Makassar on 2021-03-22