

PAPER NAME

## OA-17-696-final-B.pdf

WORD COUNT CHARACTER COUNT

3397 Words 17924 Characters

PAGE COUNT FILE SIZE

8 Pages 193.9KB

SUBMISSION DATE REPORT DATE

May 1, 2023 1:15 PM GMT+8 May 1, 2023 1:15 PM GMT+8

# 3% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.

- 3% Internet database
- Crossref database
- 0% Submitted Works database

- 0% Publications database
- Crossref Posted Content database

# Excluded from Similarity Report

- · Bibliographic material
- · Cited material
- · Manually excluded sources

- Quoted material
- Small Matches (Less then 10 words)

Article type: Research Article

ISSN: 0976-8610 Year: 2017 Volume: 8 Issue: 4

Article numbers: OA-17-696

## Used Freon Influential against Balance Life at the Earth

## Soetyono Iskandar\*, Mardi Syahir, Badaruddin Anwar

Department of Mechanical Engineering, Universitas Negeri Makassar, Makassar, South Celebes, Indonesia

#### **ABSTRACT**

One of material that have considered as source pollution at the atmosphere there are Freon, as Chloro-fluoro Carbon (CFC), or as Hydro Chloro-fluoro Carbon (HCFC). This CFC could do process photo condensation with radiation ultra violet that high energy at stratosphere and the next would reaction with ozone.

**Keywords:** Earth degradation, Pollution, Cancer, Atmosphere, Photosynthesis

#### INTRODUCTION

Empty ozone event at atmosphere caused radiation ultraviolet that from sun have not held at the atmosphere but through to the earth. Affected against life at the earth that as caused cancer diseases, happened from secondary material pollution as photochemical *smog* and heated earth that added caused at the quality life at the earth degradation (Figure 1).

Nevertheless, special for HCFC there are that not damaged ozone, although at the atmosphere could make from green house caused.

#### Atmosphere

Atmosphere is atmosphere layer blanketing equi amplitude surface of earth divided to troposphere, stratosphere, mesosphere and thermosphere. Layer troposphere (ozone layer) consisted of gathering of gas and there are two the numbers gas types big that is: oxygen gas (O<sub>2</sub>) counted 20,95%, Nitrogen (N<sub>2</sub>) 78,08%, the numbers medium that is: Argon counted 0,934%, carbon dioxide (CO<sub>2</sub>) 0,035%, vapor (H<sub>2</sub>O) 0,1-5%, medium of other gas (including inert gas besides Argon) there is in number traces (< 10<sup>-3</sup>%). Stratosphere layer only be consisted of ozone gas (O<sub>3</sub>), mesosphere radical Ozone and Nitrogen Oxide, medium of layer thermosphere consisted of ozone radical, oxygen and oxide nitrogen.

Atmosphere has many inter as utilities: be source of carbon dioxide (useful for fotosintesa plant sees figure 3) and oxygen (for exhalation of mortal), also sebgai important component in hydrology cycle (condenser), canopy for mortal (permeates absorbent infrared ray by the sun and transmitted again by earth). Photosynthesis usually done by plant containing leaf green matter (chlorophyll), algae and some bacterium types (Figure 2).

## Photosynthesis

Photosynthesis is way of carbon assimilation because in carbon photosynthesis free of  $CO_2$  is bound (fixation) becomes sugar for dissociation energy of diatomic reservoir molecule. Other way done by organism in carbon assimilation is through chemosynthesis that is usually done by sulfur bacterium (Figure 3).

At equi amplitude surface of the temperature soil; land; earth around  $15^{\circ}$ C, at elevation of the 15 Km temperature reduces to  $-56^{\circ}$ C, 50 Km rising temperature returns until  $-2^{\circ}$ C, 85 Km Temperature went down again until  $-92^{\circ}$ C, hereinafter at above elevation is 85 Km temperature returns to increasing so that at elevation of 500 Km from equi amplitude surface of the temperature ground reaching  $1200^{\circ}$ C (Figure 4).

District troposphere has derivation characteristic of temperature with increasing of elevation, because distance from radiation of ground increases, at temperature stratosphere district then increases as result of existence of permeable ozone of ultraviolet radiation, in district mesosphere temperature returns to declines because the lessen of permeable material of radiation coming from the sun, medium at district thermosphere the temperature returns to increasing because in this district happened absorption process of radiation having gumption height so that the temperature can be reach 1200 °C. Figure condition of atmosphere as explained to be visible above at above figure 1 and figure 2 following (Figure 5).

Besides correlation between temperatures, atmospheric pressure and elevation from sea level can be formulated with equation of continuity following:

$$P_h = P_o \; e^{\; -Mgh/RT}$$

Where  $P_o$  it is atmospheric pressure at elevation of zero (sea level/surface sea water), M is molecule weight average of atmosphere (28,97 g/mol at troposphere), g is gravitation (981 cm det<sup>-2</sup>), h it is elevation (cm), and R is gas constant (8,314.x107 erg det<sup>-1</sup> mol<sup>-1</sup>).

If equation of continuity above logarithm and by assuming pressure at sea surface equal to 1 atm is obtained by equation of continuity:

$$LogP_h = \frac{Mgh.10^5}{2,303RT} - - - - - - - - (1)$$

With grooving between logarithms  $P_h$ , temperature to elevation, visible figure 5. From above figure 5 seen that at elevation around 15 Km Atmosphere temperature comes near - 60  $^{0}$ C, 50 Km becomes - 2  $^{\circ}$ C, 85 Km equal - 92  $^{\circ}$ C and hereinafter temperature increases again, however atmospheric pressure more and more low if distance from earth more and more far.

#### **Geothermic Equilibrium**

Solar energy which up to earth atmosphere 1340 watt/m<sup>2</sup> and some of this dissociation energy of diatomic permeated on the way to to equi amplitude surface of earth by atmosphere, cloud and by the particle in atmosphere so that reaching equi amplitude surface of the dissociation energy of diatomic earth only 636 watt/m<sup>2</sup>. Dissociation energy of diatomic which up to equi amplitude surface of this earth partly are bounced again to atmosphere in the form of infrared temperature and radiation to balance temperature in earth.

Equi amplitude surface temperature of earth taken care of to shifts  $15^{\circ}$ C caused by vapor and gas  $CO_2$  where permeable  $CO_2$  of radiation which in transmitting by earth and hereinafter he radiation some of earth equi amplitude surfaces. If this thing is not happened equi amplitude surface temperature of earth will become -  $18^{\circ}$ C. Equi amplitude surface temperature of earth will go up when grade  $CO_2$  in atmosphere increases and this in mentioning greenhouse effect. This thing might possibly happened caused by baking activity of increasing forest fuel, so that equilibrium of  $CO_2$  in atmosphere annoyed and reaction of photosynthesis in earth decreases.

$$H_2O$$
 +  $CO_2$  +  $hv$   $\longrightarrow$   $(CH_2O)_n$  +  $O_2$ 

Besides, ozone in atmosphere is also temperature ballast in ground because ozone stands permeates ultraviolet radiation stepping into atmosphere at wavelength 220 - 330 nm based on reaction:

$$O_3 + hv (220 - 330 nm)$$
  $\longrightarrow$   $O_2 + O^*$ 

If this ultraviolet radiation not in permeated hence temperature which up to equi amplitude surface of earth around 1000°C and if this happened hence difficult there are life in this earth. Ultraviolet [light/ray] which up to earth can cause the happening of skin neoplastic disease and in also on the way towards to earth can generate pollution of secondary in the form of photochemical smog in Troposphere (Figure 6).

That equilibrium of ozone remains to happen  $O_2$  and free O in atmosphere must be enough and required also other molecule like  $N_2$  and  $O_2$  which can function to permeate excess of energy reaction of formation of ozone

$$O + O_2 + M \longrightarrow O_3 + M$$
 (the energy increases.)

Oxygen there is at atmosphere, lithosphere, hydrosphere and biosphere as seen at figure 4 and 5 and oxygen at district troposphere plays a part important in process of happened in grounds, especially at process happened atmosphere, orthogonal transformation geochemistry and in process of life. Permeable oxygen atom of visible light at wavelength 636, 630 and 538 nm and hereinafter yields oxygen ion and electron transmitting radiation electromagnet especially at infrared spectrum district.

$$O + hv \longrightarrow O^+ + e^-$$

The Atom and ion and oxygen molecule in atmosphere can react with other molecule like  $N_2$ ,  $NO_x$ ,  $H_2S_2$ ,  $SO_x$ , Cl O, hydrocarbon compound, etc.

Compound Chlorofluorocarbon (CFC), Hydrofluorocarbon (HFC), Hydroclorofluorocarbon (HCFC) and Perfluorocarbon (PFC).

Compound Chlorofluorocarbon like for example Dichlorodifluoromethane, common is named by Freon and has is volatile. This nontoxic and stable compound and some last decades exploited as coolant. Some compounds produced on a large scale inter as

<sup>2</sup> 
$$Cl_3F$$
 (CFC – 11, bp 24 °C),  $CCl_2F_2$  (CFC – 12, bp -28 °C)  $C_2Cl_3F_3$  (CFC – 113),  $C_2Cl_2F_4$  (CFC – 114) dan  $C_2ClF_5$  (CFC – 115). Dupon, 1996.

Halon is the same compound and contains bromine and in general is applied as component of fire company. Compound halon which has been commercialized for example  $CBrF_3$  (Halon - 1301) and  $C2Br_2F_4$  (Halon - 2402). This very effective compound stops fire because breaking hydrogenic atom cause of burning, based on reaction of following:

Because in character which can break ozone layer, hence usage of this halon limited. Besides in the form of halofluorocarbon, has been made integer in the form of the hydrogen compound as Hydroclorofluorocarbon (HCFC) and Hydro fluorocarbon (HFC) like CH<sub>2</sub>FCF<sub>2</sub> (HFC - usual 134a mixed at CFC - 12 at equipments of coolant from carriage and refrigerator), CHCL<sub>2</sub>CF<sub>3</sub> (HCFC - 123 generally is mixed with CFC - 11 needs of plastic foam blowing), CH<sub>3</sub>CCL<sub>2</sub>F (HCFC - 141b, also is mixture CFC - 11 for plastic foam food container), Also commonly use HCFC - 124 (CHCLFCF<sub>3</sub>), HCFC - 142b (CH<sub>3</sub>CCLF<sub>2</sub>) and HFC - 152a (CH<sub>3</sub>CHF<sub>2</sub>).

CFC although has property that is reactive not, but because has been produced is mass that is one half a million m<sup>3</sup> per year and later in the year 1974 compounds chlorofluoromethane earns breaks ozone layer in stratosphere, hence since then started looked for substitution material of friendly other coolant of area.

Although in this case producer CFC - 12 by US-EFA until the year 1995 looking for other alternative of substitution CFC.

Why Freon is assumed by dangerous in process of ruining of ozone is because in reacting it is entangling reaction to relate to where one molecules CFC can destroy many ozone molecules mellaui reaction of fotode compostioni following:

$$Cl_2CF_2$$
 + hv  $\longrightarrow$   $Cl$  +  $ClCF_2$ 

Radical Clor formed will react with ozone to yield radical ClO and hereinafter this radical reacts with free oxygen atom yielded from result of resolution of ozone is stratosphere by base reaction of following:

$$O_3$$
 + hv  $\longrightarrow$   $O_2$  +  $O$ 
 $ClO$  +  $O$   $\longrightarrow$   $Cl$  +  $O_2$ 
 $Cl$  +  $O_3$   $\longrightarrow$   $ClO$  +  $O_2$ 

Besides radical CIO earns also reacts with the nitrogen oxide in atmosphere yields radical Color and this radical hereinafter will react with ozone in relates to.

ClO + NO 
$$\longrightarrow$$
 Cl + NO<sub>2</sub>
Cl + O<sub>3</sub>  $\longrightarrow$  ClO + O<sub>2</sub>

Compound CFC has lifetime at stratosphere layer in order hundreds of year and because the reaction property relates to this causes incidence of aperture in ozone recognized by the name "Antarctic Ozone Hole." Compound hydrofluorocarbon (HFC) and hydroclorofluorocarbon (HCFC) there is also doesn't endanger ozone layer like: HFC - 134a, CF<sub>3</sub>CH<sub>2</sub>F, because although reacted with hydroxyl radical however the reaction only takes place at district troposfir:

$$CF_3CH_2F + HO \longrightarrow CF_3CHF + H_2O$$

Radical formed at above reaction will react with oxygen to yield peroxide radical and hereinafter will react with okssida nitrogen then decomposes forming radical CF<sub>3</sub>, where its(the reaction takes place as follows:

$$CF_3CH_2F + HO \longrightarrow CF_3CHF + H_2O$$
 $CF_3CHF + O_2 + M \longrightarrow CF_3CHFO_2 + M$ 
 $CF_3CHFO_2 + NO \longrightarrow CF_3CHFO + NO_2$ 
 $CF_3CHFO \longrightarrow CF_3 + HC(O)F$ 

Or reacts with oxygen molecule yielding radical perocsil:

$$CF_3CHFO + O2 \longrightarrow CF_3C(O)F + HO_2$$

Other reaction that is formation of radical CFO is earning also reacts with ozone in relates to, but the reaction the end when this radical meets with nitrogen oxide and or **methane gas.** 

$$CF_3O + O_3 \longrightarrow CF_3OO + O_2$$
 $CF_3OO + O_3 \longrightarrow CF_3O + O_2$ 
 $CF_3O + NO \longrightarrow CF_3O + FNO$ 
 $CF_3O + CH_4 \longrightarrow CF_3OH + CH_3$ 

Radical  $CF_3$  is not met in stratosphere layer to show HFC and HCFC something still can be exploited as coolant and or **aerosol.** 

Special for compound Perfluorocarbon (PFC) like Carbontetrafluoride (CF<sub>4</sub>) and Heksafluoroetana (C<sub>2</sub>F<sub>6</sub>) be compound having the character of non-toxic, doesn't react with hydroxyl radical, the other ozone and substance in atmosphere. But this compound earned as its cause hot improvement in earth because earning permeable of infrared radiation ably some thousand bigger times compared to with carbon dioxide.

Performance of compound reacts with ozone hardly determined by molecular weight, number of atoms hydrogen, klor, and its (the fluor is and potential deflesi (PD) its to ozone. Between compound CFC and HCFC which can destroy ozone layer is CFC - 11 (PD 1,0), HCFC - 22 (PD 0,030), HCFC - 123 (PD 0,013), HCFC - 141b (PD 0,10), HCFC - 124 (PD 0,035) dan HCFC - 142B (PD 0,038). (Tuazon, E., and R. Atkinson. 1993.

More and more low potential value of its deflesi, lifetime radical yielded by CFC or HCFC at layer troposfir dwarf, so that has fallen to pieces at layer troposfir before migration to stratosphere layer.

### **CONCLUSION**

Compound that is most dominance destroys ozone layer is all types CFC and HCFC which is not contained  $CF_3$ . Alternative which is usable permanent from compound fluorocarbon is HCFC is containing  $CF_3$  and perfluorocarbon (PFC). 1 molecule = 224 litres (Ozone).

Our earth are indium actually clean and lapped over is regular like at figure, only deed of human being so that impure. Because freon rest of from workshop that is is not is accommodated, because wishing quickly is discharged into the air free which. Very dangerous freon for human life, so that we try writes thing like this. We urge to all men to earn consciousness that this very dangerous Freon.

Earth has experienced transformation of big area like height of Cosentration greenhouse gas because man activity that it is said can generate transformation of climate, like height of content CFCs in atmosphere destroying ozone layer. Where also happened area damage, like forest damage, erosion, annihilate various floral species and fauna (Zoer'aini DI: 2005). In forest area Indonesia degradation has reached wide 59,17 Ha, while area of stall is outside by forest area is noted with a width of 41,47 Ha, this data taken up to the year 2004 (Department of Forestry: 2006). Environmental damage is caused by increase of resident that is is not controlled and incommensurate to improvement of quality or performance in managing this natural resources. Beside damage of area which caused by man hand, also happened damage naturally or event of nature there is in the universe.

Because generally environmental damage happened as result of man hand who is irresponsible, therefore to improve; repair damage of area which already happened, and lessen and prevents damage which can happened, hence early on man is given by education of environment like: what interaction with nature, knows cause and as result of area damage, teachs form of activities to preserve nature. Besides man as governor is earth by this hence claimed also to protect and preserves earth that the benefit is always can awake. According to newest research pickings concludes, that intellectual development happened very fast in the early of child life.

At the age of four years a child has formed 50% intelligence which will be owned it after adult, 30% again at the age of eight years, and 20% the rest at mid of final a decay second, therefore perceivable when age four first years in development of child of conceived of golden age (Jalal Fasli: 2002). Mean at the age of the child of starting taught by education of environment in the form of knowledge and understanding to nature area, to ecosystem and awareness conservation of nature (Lexy JM: 2004). Learning man, springs up and grows from experience that is first of all obtaining of through family life (Conny Semiawan: 1997) together with parent fellow, hence as a stripper which will supply the child requiring spiritual and understanding of adequate area. Parent fellow is learning is first for his own children, teaches environmental education like protecting and preserves its area, comprehends cause and effect of environmental damage happened like its breakdown is environmental around forest, ozone layer, flora and fauna. Parent fellow also place of questioning for the

children, answers given and old fellow caring to the chlid is yesterday, today and tomorrow determines to the future of ground which has experienced damage. Then who is old fellow that is very stands for education of chlid? Mother

At present modern age, function of family as place of educate, takes care of and takes care of chlid to start shifting because the increasing of old fellow especially Mother. Working outdoors (Depdiknas: 2003). Besides, not all old fellows now has good understanding to area, and awareness to comprehend education of environment to the chlid. Hence joining child of in its environments is a real many the knowledges to chlid.

God said (Greek logos). Said "logos" means life uniting the universe. Helai dau, harmonious of season, the stars in the sky in the form of perfect because of logos. Philosopher Heracleitus, Plutarch, Philo and Pluto, and so it is with the philosophers "Stoic" writes thing which about "logos".

What makes everlasting life promises is important is, although all makluk life in this earth died, but only man which can understand thinking of everlasting life. Only man us which can understand difference between brief lives in this earth and expectation of everlasting life in coming. And we also can understand how death to earn at any times snatchs us because stuffing up vein, a drunken power steering, or virus HIV which never last.

What a respects it expectation which we have, understands that death is not end of anything; but, that only when sleep for just momentary for us, and happened the next of we know that we would together with Jesus forever and ever.

#### REFERENCES

- D. Pasek, N. P. Tandian, A. Suwono, T. Hardianto, T. A. F. Soelaiman, "Indonesian Experience in Retrofitting of Some Milk Cooler Systems, Seminar on ODS Phase Out: Solution for the Refrigerantion Sector, Kuta, Mei 05 – 07, 1999.
- 2. A. Suwono, A. D. Pasek dan, N. P. Tandian, "Application Expansion of Propane Mixture /n-Butana as Refrigerant as a change of CFC, *Seminar on Altenative Refrigerants*, Bandung, 5 juni 1996.
  - A. Suwono, N. P. Tandian, A. D. Pasek, T. Hardianto, T. A. F. Soelaiman, "Indonesian Experience on Hydrocarbon Application as Refrigeran Substituting the Phase Out CFC", *Natural Working Fluid '98 IIR-Gustav Lorentzen Conference*, 557 566, 1998.
- 3. Suwono, N. P. Tandian, A. D. Pasek, T. Hardianto, T. A. F. Soelaiman, N. Damitri, P. Atmadi, C, Soekardi, "Retrofit and Conversion of a R-22 Chiller to R-290 and its Performance", *Seminar on ODS Phase Out: Solution for the Refrigerantion Sector*, Kuta, Mei 05 07, 1999.
- 4. Suwono, A. D. Pasek dan, N. P. Tandian, "Looks For Friendliness Refrigerant Area of", *Republika*, hal.6, 5 juni 1996.
- 5. T. Wirakusumah, "The Government Policy Concerning the Prohibition to Produce and to trade Ozone Depleting Substance and Equipment Using Ozone Depleting Substance in Indonesia", *Seminar on ODS Phase out Solution for the Refrigeration Sector*, Kuta, Mei 05 07 1999.
- 6. Conny Semiawan. (1997) Perspective of Education of Talented Child.. Jakarta, Grasindo.
- 7. Departement of Forest (2006) Ohio Forests. Northen, Research Station.
- 8. Depdiknas. (2003) *Undang-Undang Sistem Pendidikan Nasional Tahun 2003*. Direktorat A. Pendidikan Nasional.
- 9. Jalal, Fasli, "Education of Child Of Early Age, Basic education", Bulletin Padu, Edisi Perdana, Direktorat Jendral Pendidikan Luar Sekolah, Depdiknas Jakarta, 2002.
- 10. N. Damitri, C, Soekardi, A. Suwono, N. P. Tandian, A. D. Pasek, T. Hardianto, T. A. F. Soelaiman, P. Atmadi, "Performance Comparison of CFC 12, HCR 12 and HCR 12<sup>+LFS</sup> as Refrigerats", *Seminar on ODS Phase Out: Solution for the Refrigerantion Sector*, Kuta, Mei 05 07, 1999.
- 11. N. P. Tandian, A. Suwono, C. A. Sylva, A. D. Pasek, "Application of Propana /n-Butane Mixture as Refrigerant in Milk Cooling Unit", *International Conference of Fluid and Thermal Energy Conversion* '97, Yogyakarta 21 24 Juli 1997.
- 12. N. P. Tandian, A. Suwono, C. A. Sylva, A. D. Pasek, T. Hardianto, T. A. F. Soelaiman, B. T. Prasetyo, Sarwono, Kafi'uddin, "Performance Testing og Indonesia HC Refrigerant", *Seminar on ODS Phase Out: Solution for the Refrigerantion Sector*, Kuta, Mei 05 07, 1999.
- 13. Nimitz, J. S., and S. R. Skaggs. 1992. "Estimating Tropospheric Lifetime and Ozone Depletion Potential of One and Two Carbon Hydrofluorocarbon and Hydrochlorofluorocarbons," Environ Sci. Technol., 26: 739-744.
- 14. K. Watanabe, J. V. Widiatmo, "Alternative Refrigerant and Their Thermophysical Properties Research", *Seminar on ODS Phase Out: Solution for the Refrigeration Sector*, Kuta, Mei 05 07, 1999.

- 15. Ravishankara, A. R. A. A. Turnipseed, and S. Salomon. 1993. "Do Hydrofluororcarbons Destroy Stratospheric Ozone?." Science, 263: 71 75.
- 16. S. Devotta, M. M. Kulkami, N. N. Sawant, P. A. Patil, N. K. Sane. "Compressor Life Test with HC Refrigerants", *Proceeding of Natural Working Fluid '98 IIR Gustav Lorentzen Conference*, Oslo, Norwegia 1998.
- 17. Tuazon, E., and R. Atkinson. 1993. "Tropospheric Degradation Production of CH2FCF3(HFC 134a)," J. Atmos. Chem., 16: 301 312.
- 18. Wazter, J. W., L. Froidevaux, and R. S. Harwood. 1993. "Stratospheric CIO and Ozone from the Microwave Limb Sounder on the Upper Atmosphere Research Satelite," Nature, 362: 597 602.
- 19. Zoer'aini Djamal. 2005. "Forest area Indonesia," P. Jakarta: Bumi Aksara. P. 179.
- 20. Zurer, P. 1993. "Perfluorocarbons Use, Emissions May Face Restriction," Chem.Eng. News, August 9: P.16
- 21. Zurer, P. 1994. "CFC Substituens Proven Save for Ozone Layer," Chem.Eng. News, Januari 10: P. 5-6.



# 3% Overall Similarity

Top sources found in the following databases:

- 3% Internet database
- Crossref database

Internet

• 0% Submitted Works database

- 0% Publications database
- Crossref Posted Content database

## **TOP SOURCES**

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.





# Excluded from Similarity Report

- Bibliographic material
- Cited material
- Manually excluded sources

- Quoted material
- Small Matches (Less then 10 words)

**EXCLUDED SOURCES** 

# primescholars.com

Internet

81%